

CUSPED HITCHIN REPRESENTATIONS AND ANOSOV REPRESENTATIONS OF GEOMETRICALLY FINITE FUCHSIAN GROUPS

RICHARD CANARY, TENGREN ZHANG, AND ANDREW ZIMMER

ABSTRACT. We develop a theory of Anosov representation of geometrically finite Fuchsian groups in $\mathrm{SL}(d, \mathbb{R})$ and show that cusped Hitchin representations are Borel Anosov in this sense. We establish analogues of many properties of traditional Anosov representations. In particular, we show that our Anosov representations are stable under type-preserving deformations and that their limit maps vary analytically. We also observe that our Anosov representations fit into the previous frameworks of relatively Anosov and relatively dominated representations developed by Kapovich-Leeb and Zhu.

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Anosov representations into $\mathrm{SL}(d, \mathbb{K})$	7
4. Basic properties of Anosov representations	11
5. Basic properties of cusp representations	15
6. A dynamical characterization of linear Anosov representations	18
7. Hitchin representations are Borel Anosov	23
8. Stability of Anosov representations	25
9. Positive representations in the sense of Fock-Goncharov	33
Appendix A. Constructing cusp representations	37
Appendix B. Anosov representations into semisimple Lie groups	37
References	43

1. INTRODUCTION

In this paper we develop a theory of Anosov representations for geometrically finite Fuchsian groups into $\mathrm{SL}(d, \mathbb{K})$ where \mathbb{K} is either the field of real numbers or the field of complex numbers (and more generally, into any semisimple Lie group G with finite center). This theory shares two crucial features with Anosov representations of convex cocompact Fuchsian groups. First, if $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite Fuchsian group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is Anosov, then there is a ρ -equivariant quasi-isometric embedding of an orbit of Γ in \mathbb{H}^2 into the Riemannian symmetric space $X_d(\mathbb{K})$ associated to $\mathrm{SL}(d, \mathbb{K})$. Moreover, there is a ρ -equivariant map of the limit set of Γ into the appropriate (partial) flag variety. Second, small deformations of Anosov representations which preserve the conjugacy class of the images of parabolic elements in Γ remain Anosov. We further show that these limit maps vary analytically in ρ . We observe that our representations are relatively Anosov representations in the sense of Kapovich and Leeb [18] and relatively dominated representations as defined by Zhu [34]. Our concrete setting allows for simpler proofs and more explicit results.

Date: March 11, 2022.

Canary was partially supported by grant DMS-1906441 from the National Science Foundation and grant 674990 from the Simons Foundation. Zhang was partially supported by the NUS-MOE grant R-146-000-270-133. Zimmer was partially supported by grants DMS-2105580 and DMS-2104381 from the National Science Foundation.

Our main motivation was to study the class of cusped Hitchin representations. A representation $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ of a geometrically finite Fuchsian group is said to be *cusped Hitchin* if there exists a continuous ρ -equivariant positive map from the limit set of Γ into the full flag variety \mathcal{F}_d . We show that cusped Hitchin representations are irreducible and Borel Anosov (i.e. they are P_k -Anosov for all $1 \leq k \leq d-1$). This generalizes results of Labourie [22], when Γ is cocompact, and Labourie-McShane [23] (see also Burelle-Treib [9]), when Γ is convex cocompact. (Recent results of Sambarino [30] also imply that cusped Hitchin representations are irreducible). Fock and Goncharov [14] introduced the theory of positive representations. They consider the case where Γ is not cocompact but has cofinite area, and they only require that the limit map be defined on fixed points of peripheral elements of Γ . We show that type-preserving positive representations, in their sense, are in fact cusped Hitchin representations. Other examples of cusped Anosov representations are provided by exterior powers of cusped Hitchin representations and cusped Barbot representations.

In turn, the motivation for studying cusped Hitchin representations arises from an intriguing potential analogy with the augmented Teichmüller space from classical Teichmüller theory. The augmented Teichmüller space of a closed orientable surface S is obtained by appending to the Teichmüller space of S points corresponding to (marked) finite area hyperbolic structures on the complement in S of any multicurve on S . Masur [28] showed that the augmented Teichmüller space is the metric completion of Teichmüller space with the Weil-Petersson metric. Loftin and Zhang [24] construct an analytic model for an augmented Hitchin component. In the near future, we plan to combine our work with thermodynamical results from a companion paper by Bray, Canary, Kao and Martone [5] to construct pressure metrics on deformation spaces of cusped Hitchin representations, generalizing work of Bridgeman, Canary, Labourie and Sambarino [7]. The hope is that this will allow us to investigate whether the augmented Hitchin component is the metric completion of the Hitchin component with the pressure metric.

We now turn to a more detailed discussion of our work. Let $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ be a geometrically finite group with limit set $\Lambda(\Gamma) \subset \partial\mathbb{H}^2$. Suppose that $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a representation and that we are given a continuous ρ -equivariant map

$$\xi_\rho = (\xi_\rho^k, \xi_\rho^{d-k}) : \Lambda(\Gamma) \rightarrow \mathrm{Gr}_k(\mathbb{K}^d) \times \mathrm{Gr}_{d-k}(\mathbb{K}^d)$$

into the Grassmanians of k -planes and $(d-k)$ -planes in \mathbb{K}^d . We require that ξ_ρ is transverse, i.e.

$$\xi_\rho^k(x) \oplus \xi_\rho^{d-k}(y) = \mathbb{K}^d \quad \text{if } x \neq y \in \Lambda(\Gamma).$$

We obtain an associated splitting

$$E_\rho = \mathrm{U}(\Gamma) \times \mathbb{K}^d = \Theta^k \oplus \Xi^{d-k} \quad \text{where } \Theta^k|_v = \xi_\rho^k(v^+) \quad \text{and} \quad \Xi^{d-k}|_v = \xi_\rho^{d-k}(v^-),$$

and $\mathrm{U}(\Gamma) \subset T^1\mathbb{H}^2$ is the set of tangent vectors which extend to geodesics both of whose endpoints lie in $\Lambda(\Gamma)$. This descends to a splitting

$$\widehat{E}_\rho = \Gamma \backslash E_\rho = \widehat{\Theta}_\rho^k \oplus \widehat{\Xi}_\rho^{d-k}$$

of the flat bundle associated to ρ over the non-wandering portion $\widehat{\mathrm{U}}(\Gamma) = \Gamma \backslash \mathrm{U}(\Gamma)$ of the geodesic flow. The geodesic flow on $\widehat{\mathrm{U}}(\Gamma)$ lifts naturally to a flow on \widehat{E}_ρ which preserves the splitting and is parallel to the flat connection. We say that ρ is P_k -Anosov if the associated flow on $\mathrm{Hom}(\widehat{\Xi}_\rho^{d-k}, \widehat{\Theta}_\rho^k)$ is uniformly contracting with respect to the (operator) norm arising from **some** family of continuous norms on the fibers of \widehat{E}_ρ (see Definition 3.1 for details). In this case, we call ξ_ρ the P_k -Anosov limit map of ρ . When Γ contains a parabolic element, we will sometimes refer to our P_k -Anosov representations as *cusped P_k -Anosov* representations to distinguish them from traditional Anosov representations.

We obtain generalizations of many of the classical properties of Anosov representations, see Labourie [22] or Guichard-Wienhard [16], in our setting. We say that a ρ -equivariant, continuous map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \mathrm{Gr}^k(\mathbb{K}^d) \times \mathrm{Gr}^{d-k}(\mathbb{K}^d)$ is *strongly dynamics preserving* if whenever $\{\gamma_n\}$ is a sequence in Γ , $\gamma_n(z) \rightarrow x \in \Lambda(\Gamma)$, and $\gamma_n^{-1}(z) \rightarrow y \in \Lambda(\Gamma)$ for some (any) $z \in \mathbb{H}^2$, then

$$\rho(\gamma_n)(V) \rightarrow \xi^k(x)$$

for any $V \in \text{Gr}(\mathbb{K}^d)$ which is transverse to $\xi^{d-k}(y)$. Given $g \in \text{SL}(d, \mathbb{K})$ let

$$\lambda_1(g) \geq \cdots \geq \lambda_d(g)$$

denote the absolute values of the (generalized) eigenvalues of g . Then g is P_k -proximal if $\lambda_k(g) > \lambda_{k+1}(g)$ and g is weakly unipotent if the (multiplicative) Jordan-Chevalley decomposition of g has elliptic semisimple part and non-trivial unipotent part.

Theorem 1.1. *If $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is P_k -Anosov, then*

(1) *For any $z_0 \in \mathbb{H}^2$, there exists $A, a > 1$ so that if $\gamma \in \Gamma$, then*

$$\frac{1}{A} \exp\left(\frac{1}{a} d_{\mathbb{H}^2}(z_0, \gamma(z_0))\right) \leq \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq A \exp(ad_{\mathbb{H}^2}(z_0, \gamma(z_0))).$$

(2) *There exists $B, b > 1$ so that if $\gamma \in \Gamma$, then*

$$\frac{1}{B} \exp\left(\frac{1}{b} \ell(\gamma)\right) \leq \frac{\lambda_k(\rho(\gamma))}{\lambda_{k+1}(\rho(\gamma))} \leq B \exp(b\ell(\gamma))$$

where $\ell(\gamma)$ is the translation length of γ on \mathbb{H}^2 .

(3) *The P_k -Anosov limit map ξ_ρ is strongly dynamics-preserving and unique. In particular, if $\alpha \in \Gamma$ is parabolic, then $\rho(\alpha)$ is weakly unipotent, while if $\gamma \in \Gamma$ is hyperbolic, then $\rho(\gamma)$ is P_k -proximal.*

(4) *If $z_0 \in \mathbb{H}^2$ and x_0 is a point in the symmetric space $X_d(\mathbb{K})$ associated to $\text{SL}(d, \mathbb{K})$, then the orbit map $\tau_\rho : \Gamma(z_0) \rightarrow X_d(\mathbb{K})$ given by $\tau_\rho(\gamma(z_0)) = \rho(\gamma)(x_0)$ is a quasi-isometric embedding.*

We also give a dynamical characterization of Anosov representations in the spirit of characterizations of traditional Anosov representations by G uritaud-Guichard-Kassel-Wienhard [15], Kapovich-Leeb-Porti [19] and Tsouvalas [33].

Theorem 1.2. *Suppose $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation. Then ρ is P_k -Anosov if and only if there exists a ρ -equivariant, transverse, continuous, strongly dynamics preserving map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$. Furthermore, ξ is the P_k -Anosov limit map.*

In general, being P_k -Anosov is not an open condition in the space of representations of a geometrically finite group. For instance, consider the case where $\Gamma = \langle g_1, g_2 \rangle \subset \text{PSL}(2, \mathbb{R})$ is a free group, g_2 is parabolic, and $\rho : \Gamma \rightarrow \text{SL}(4, \mathbb{R})$ is P_1 -Anosov with

$$\rho(g_2) = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Next define a family of representations ρ_t where $\rho_t(g_1) = \rho(g_1)$ and

$$\rho_t(g_2) = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & t \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Then ρ_t is not P_1 -Anosov for any $t \neq 0$ since the sequence $\{\rho_t(g_2)^n\}$ does not converge to a rank one element of $\mathbb{P}(\text{End}(\mathbb{R}^4))$. Bowditch [3, Sec. 5] gave an example of a sequence $\{\rho_n\}$ of indiscrete representations of a free group on two generators into $\text{SO}(4, 1)$ which converge to a geometrically finite representation ρ , so that $\rho_n(\alpha)$ is parabolic if and only if $\rho(\alpha)$ is parabolic. However, in his example $\rho_n(\alpha)$ is not conjugate to $\rho(\alpha)$ for any n . Bowditch [3, Thm. 1.5] also established a stability theorem for deformations of geometrically finite representations which preserve the structure of the Jordan decomposition of parabolic elements.

To account for these examples, we introduce the following subset of the representation variety. If $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation of a geometrically finite Fuchsian group, let

$$\text{Hom}_{\text{tp}}(\rho) \subset \text{Hom}(\Gamma, \text{SL}(d, \mathbb{K}))$$

be the space of representations $\sigma : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ so that if $\alpha \in \Gamma$ is parabolic, then $\sigma(\alpha)$ is conjugate to $\rho(\alpha)$. We obtain the following stability result for type-preserving deformations. (In their preprint, Kapovich and Leeb [18] suggest that such a stability result holds more generally.) Furthermore, we show that limit maps vary analytically. Combined with work of Bray-Canary-Kao-Martone [5] this will imply that entropy and pressure intersection vary analytically over the cusped Hitchin component (which will be used crucially in the construction of the pressure metric). The proof also allows us to see that the P_k -Anosov limit maps are uniformly Hölder in a neighborhood of a P_k -Anosov representation.

We say that $\{\rho_u\}_{u \in M}$ is a \mathbb{K} -analytic family of representations if M is a \mathbb{K} -analytic manifold and the map $u \rightarrow \rho_u$ is a \mathbb{K} -analytic map from M into $\mathrm{Hom}(\Gamma, \mathrm{SL}(d, \mathbb{K}))$.

Theorem 1.3. *If $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is geometrically finite and $\rho_0 : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov, then there exists an open neighborhood \mathcal{O} of ρ_0 in $\mathrm{Hom}_{\mathrm{tp}}(\rho_0)$, so that*

- (1) *If $\rho \in \mathcal{O}$, then ρ is P_k -Anosov.*
- (2) *There exists $\alpha > 0$ so that if $\rho \in \mathcal{O}$, then its P_k -Anosov limit map ξ_ρ is α -Hölder.*
- (3) *If $\{\rho_u\}_{u \in M}$ is a \mathbb{K} -analytic family of representations in \mathcal{O} and $z \in \Lambda(\Gamma)$ then the map from M to $\mathrm{Gr}_k(\mathbb{K}^d) \times \mathrm{Gr}_{d-k}(\mathbb{K}^d)$ given by $u \rightarrow \xi_{\rho_u}(z)$ is \mathbb{K} -analytic.*

When $\widehat{\Gamma}$ is compact, stability follows from standard arguments in hyperbolic dynamics and the Hölder regularity of the boundary maps is a consequence of standard results, e.g. [31, Cor. 5.19]. The non-compact case is more involved. Our key idea to prove stability is to observe that if $\rho_2 \in \mathrm{Hom}_{\mathrm{tp}}(\rho_1)$, then on each cusp there is a smooth conjugacy of the flows associated to ρ_1 and ρ_2 . This is made precise in Equation (21) below. This essentially means that the two flows differ by a compact perturbation and thus, essentially, reduces to the compact base case. Our key idea to prove Hölder regularity is to first introduce certain “canonical families of norms” on the flow spaces, see Section 3.1. Then we prove that if the flow is contracting with respect to any family of norms, then the flow is contracting with respect to any canonical family of norms (see Theorem 6.1 and Theorem 4.1). Finally, the canonical family of norms are well-behaved enough that we can adapt an argument from [36] to prove Hölder regularity of the boundary maps directly.

We now discuss the applications of our results to cusped Hitchin representations, which was the original goal of our work. Given an ordered basis \mathcal{B} for \mathbb{R}^d , we say that a unipotent $A \in \mathrm{SL}(d, \mathbb{R})$ is *totally positive* with respect to \mathcal{B} , if its matrix with respect to \mathcal{B} is upper triangular and all its minors (which are not forced to be 0 by the fact that the matrix is upper triangular) are strictly positive. The set $\mathcal{U}_{>0}(\mathcal{B})$ of unipotent, totally positive matrices with respect to \mathcal{B} is a semi-group (see Lusztig [25, Section 2.12]). Following Fock and Goncharov [14], we say that an ordered k -tuple (F_1, F_2, \dots, F_k) of distinct flags in \mathcal{F}_d is positive, if there exists an ordered basis $\mathcal{B} = (b_1, \dots, b_d)$ for \mathbb{R}^d so that $b_i \in F_1^i \cap F_k^{d-i+1}$ for all i , and there exists $u_2, \dots, u_{k-1} \in \mathcal{U}_{>0}(\mathcal{B})$ so that $F_i = u_{k-1} \cdots u_i F_k$ for all $i = 2, \dots, k-1$. If X is a subset of \mathbb{S}^1 then a map $\xi : X \rightarrow \mathcal{F}_d$ is *positive* if whenever (x_1, \dots, x_n) is a cyclically ordered subset of distinct points in X , then $(\xi(x_1), \dots, \xi(x_n))$ is a positive n -tuple of flags.

We say that a representation $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ of a geometrically finite Fuchsian group is a *Hitchin representation* if there exists a continuous positive ρ -equivariant map $\xi_\rho : \Lambda(\Gamma) \rightarrow \mathcal{F}_d$. If Γ is cocompact and torsion-free then these are exactly the Hitchin representations of closed surface groups introduced by Hitchin [17] and further studied by Labourie [22], while if Γ is convex cocompact they are the same as the Hitchin representations studied by Labourie and McShane [23]. We distinguish the case where Γ contains parabolic elements by calling these *cusped Hitchin representations*. If $d = 3$ and $S = \Gamma \backslash \mathbb{H}^2$ is a finite area hyperbolic surface, then cusped Hitchin representations of Γ are holonomy representations of strictly convex, finite area, projective structures on S (see Marquis [27]). Further it follows from [14, Thm. 3.3] that holonomy maps of geometrically finite projective surfaces in the sense of Crampon-Marquis [11] are also cusped Hitchin. Fock and Goncharov [14] studied the a priori more general class of representations which admit equivariant positive maps from the set $\Lambda_p(\Gamma)$ of fixed points of peripheral elements of Γ into \mathcal{F}_d when Γ is not cocompact but has cofinite area. We show that all such type-preserving representations are in fact cusped Hitchin representations. Our main result here is that cusped Hitchin representations are P_k -Anosov for all k .

Theorem 1.4. *If $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is geometrically finite and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ is a Hitchin representation, then ρ is irreducible and P_k -Anosov for all k . Moreover:*

- (1) *For all k the map $x \mapsto \xi_\rho^k(x)$ is the P_k -Anosov limit map.*
- (2) *If $\alpha \in \Gamma$ is parabolic, then $\rho(\alpha) = \pm u$ for some unipotent $u \in \mathrm{SL}(d, \mathbb{R})$ with a single Jordan block.*
- (3) *If $\gamma \in \Gamma$ is hyperbolic, then $\rho(\gamma)$ is loxodromic.*

If we let $\widehat{\mathcal{H}}_d(\Gamma)$ denote the space of all Hitchin representations of Γ into $\mathrm{SL}(d, \mathbb{R})$, it is easy to see that $\widehat{\mathcal{H}}_d(\Gamma)$ is a real analytic manifold. In fact, one may use results of Fock and Goncharov [14] to show that the space $\mathcal{H}_d(\Gamma)$ of conjugacy classes of Hitchin representations is diffeomorphic to \mathbb{R}^m (for some m). (Marquis [26] gives an explicit parametrization of $\mathcal{H}_3(\Gamma)$ as a topological cell when $S = \Gamma \backslash \mathbb{H}^2$ is a finite area hyperbolic surface.)

Comparison to other results: As mentioned above, Kapovich-Leeb [18] and Zhu [34] have previously developed theories of Anosov representations for relatively hyperbolic groups. Their work is based on extending characterizations of Anosov representations due to Kapovich-Leeb-Porti [19] and Bochi-Potrie-Sambarino [2] respectively. Theorem 1.1 immediately implies that a P_k -Anosov representation ρ is P_k -relatively dominated, in the sense of Zhu [34]. Theorem 9.4 in [34] then implies that ρ is P_k -relatively asymptotically embedded and Remark 9.10 in [34] implies that ρ is P_k -relatively uniform RCA in the sense of Kapovich-Leeb [18].

Corollary 1.5. *Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is geometrically finite and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a P_k -Anosov representation. Then ρ is P_k -relatively dominated, P_k -relatively asymptotically embedded and P_k -relatively uniform RCA.*

In a recent preprint, Zhu [35, Example 4.6] establishes that a cusped Hitchin representation of a geometrically finite Fuchsian group Γ is P_k -relatively dominated, with respect to its collection \mathcal{P} of parabolic subgroups, for all k . Kapovich and Leeb [18] mention that they can obtain a related result in their setting. Further, it follows from [35, Thm. C] and our Theorems 1.1 and 1.2, that a representation of a geometrically finite Fuchsian group Γ is P_k -Anosov if and only if it is P_k -relatively dominated with respect to \mathcal{P} .

In a subsequent preprint, Filip [12] introduces the class of log-Anosov representation of Fuchsian lattices, which also agrees with the class of cusped Anosov representations. He introduces adapted metrics, which correspond to our canonical norms. He shows that monodromy maps of certain variations of Hodge structures on finite area Riemann surfaces are log-Anosov and uses this result to study their properties. In particular many monodromy maps coming from hypergeometric differential equations can be analyzed in this manner, including those previously studied by Brav and Thomas [4] and by Filip and Fougeron [13].

Acknowledgements: The authors thank the referee for their careful reading of the original manuscript and helpful suggestions which improved the exposition. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930 while the first author participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2020 semester.

2. PRELIMINARIES

In this section, we recall some preliminary facts and introduce notation that will be used throughout this paper.

2.1. Hyperbolic 2-space. In this paper we will identify \mathbb{H}^2 with the Poincaré upper half plane model. For any $v \in T^1 \mathbb{H}^2$, let $r_v : \mathbb{R} \rightarrow \mathbb{H}^2$ denote the unit speed geodesic with $r'_v(0) = v$ and let

$$v^+ := \lim_{t \rightarrow +\infty} r_v(t) \in \partial \mathbb{H}^2 \quad \text{and} \quad v^- := \lim_{t \rightarrow -\infty} r_v(t) \in \partial \mathbb{H}^2$$

denote its limit points in $\partial \mathbb{H}^2 = \mathbb{R} \cup \{\infty\}$. We also let $\phi_t : T^1 \mathbb{H}^2 \rightarrow T^1 \mathbb{H}^2$ denote the geodesic flow, i.e.

$$\phi_t(v) = r'_v(t)$$

for all $v \in T^1 \mathbb{H}^2$ and $t \in \mathbb{R}$.

If $\{g_n\}$ is a sequence in $\mathrm{PSL}(2, \mathbb{R})$, we say that g_n converges to $x \in \partial \mathbb{H}^2$ if $g_n(z) \rightarrow x$ for some (any) $z \in \mathbb{H}^2$. We often simply write $g_n \rightarrow x$.

2.2. Geometrically finite Fuchsian groups: We say that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is *geometrically finite* if it is discrete, finitely generated and non-elementary (i.e. does not contain a cyclic subgroup of finite index). Let $\Lambda(\Gamma) \subset \partial \mathbb{H}^2$ denote its limit set. Then let $\mathbf{U}(\Gamma)$ denote the minimal, non-empty, closed, ϕ_t -invariant, Γ -invariant subset of $T^1 \mathbb{H}^2$, i.e.

$$\mathbf{U}(\Gamma) = \{v \in T^1 \mathbb{H}^2 : v^+, v^- \in \Lambda(\Gamma)\}.$$

Let $\widehat{\mathbf{U}}(\Gamma) := \Gamma \backslash \mathbf{U}(\Gamma)$, and note that ϕ_t descends to a flow on $\widehat{\mathbf{U}}(\Gamma)$, which we also denote by ϕ_t . If $S = \Gamma \backslash \mathbb{H}^2$, then $\widehat{\mathbf{U}}(\Gamma)$ is the non-wandering portion of $T^1 S$ and its orbits are the complete geodesics which remain entirely in the convex core of S .

Definition 2.1. (1) If $p \in \Lambda(\Gamma)$ is fixed by a parabolic element of Γ , let $\mathrm{st}_\Gamma(p) = \{\gamma \in \Gamma \mid \gamma(p) = p\}$. We call $\mathrm{st}_\Gamma(p)$ a *cuspidal subgroup* of Γ . An open horodisk H based at p is a *precisely invariant horodisk* for $\mathrm{st}_\Gamma(p)$ if whenever $\gamma \in \Gamma$ and $\gamma(H) \cap H$ is non-empty, then $\gamma \in \mathrm{st}_\Gamma(p)$. In this case, $C = \mathrm{st}_\Gamma(p) \backslash H$ is an *embedded cusp neighborhood*.
 (2) A set $\mathcal{C} = \{C_1, \dots, C_r\}$ of disjoint embedded cusp neighborhoods in $S = \Gamma \backslash \mathbb{H}^2$ is *full* if any curve on S which is represented by a parabolic element of Γ is freely homotopic into some C_i .

If $C = \mathrm{st}_\Gamma(p) \backslash H$ is an embedded cusp neighborhood, then we set

$$\mathbf{U}(\Gamma)_H = \{v \in \mathbf{U}(\Gamma) \mid r_v(0) \in H\} \quad \text{and} \quad \widehat{\mathbf{U}}(\Gamma)_C = \mathrm{st}_\Gamma(p) \backslash \mathbf{U}(\Gamma)_H.$$

If \mathcal{C} is a full collection of cusp neighborhoods for S , then we set

$$\widehat{\mathbf{U}}(\Gamma)_\mathcal{C} = \bigcup_{C \in \mathcal{C}} \widehat{\mathbf{U}}(\Gamma)_C.$$

Notice that its complement $\widehat{\mathbf{U}}(\Gamma)_\mathcal{C}^c$ is compact. We will sometimes informally refer to $\widehat{\mathbf{U}}(\Gamma)_\mathcal{C}$ as the thin part of the geodesic flow and its complement as the thick part.

2.3. The representation theory of $\mathrm{SL}(2, \mathbb{R})$: For $d \geq 1$, let $\tau_d : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ denote the standard irreducible representation. Explicitly, $\tau_1 \equiv 1$ and if $d \geq 2$, then $\tau_d = i_d \circ \bar{\tau}_d$, where

- $\bar{\tau}_d : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(\mathrm{Sym}^{d-1}(\mathbb{R}^2))$ is the representation induced by the linear $\mathrm{SL}(2, \mathbb{R})$ -action on $\mathrm{Sym}^{d-1}(\mathbb{R}^2)$ given by

$$\gamma \cdot (e_1^k e_2^{d-1-k}) \mapsto (\gamma \cdot e_1)^k (\gamma \cdot e_2)^{d-1-k}$$

for all $k = 0, \dots, d-1$, and

- $i_d : \mathrm{SL}(\mathrm{Sym}^{d-1}(\mathbb{R}^2)) \rightarrow \mathrm{SL}(d, \mathbb{R})$ is the isomorphism induced by the linear isomorphism $\mathrm{Sym}^{d-1}(\mathbb{R}^2) \simeq \mathbb{R}^d$ that identifies $e_1^k e_2^{d-1-k} \in \mathrm{Sym}^{d-1}(\mathbb{R}^2)$ with $e_{k+1} \in \mathbb{R}^d$ for all $k = 0, \dots, d-1$.

One may also regard τ_d as a representation into $\mathrm{SL}(d, \mathbb{C})$. Let $\xi_d : \partial \mathbb{H}^2 \simeq \mathbb{RP}^1 \rightarrow \mathcal{F}(\mathbb{K}^d)$ be the map defined by $[ae_1 + be_2] \mapsto F$, where $F^{(j)} = \mathrm{Span}_{\mathbb{K}}(f_1, \dots, f_j)$ and

$$f_j = \sum_{k=1}^{d+1-j} \binom{d-j}{k-1} a^{d+1-j-k} b^{k-1} e_k.$$

It is straightforward to verify that ξ_d is a continuous, τ_d -equivariant, strongly dynamics preserving map. We call ξ_d the *Veronese embedding* associated to τ_d .

One can compute that $\tau_d \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ is the upper triangular matrix in $\mathrm{SL}(d, \mathbb{R})$ given by

$$(1) \quad \left(\tau_d \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \right)_{k,j} = \binom{j-1}{k-1} a^{j-k}$$

if $1 \leq k \leq j \leq d$. Furthermore, it is well-known that the $d \times d$ upper triangular matrix given by (1) is unipotent and totally positive with respect to the standard basis of \mathbb{R}^d when $a > 0$. From this, it follows easily that ξ_d is a positive map.

2.4. Eigenvalues, singular values, and the (multiplicative) Jordan-Chevalley decomposition. Given an element $g \in \mathrm{SL}(d, \mathbb{K})$, let

$$\sigma_1(g) \geq \cdots \geq \sigma_d(g)$$

denote the singular values of g and let

$$\lambda_1(g) \geq \cdots \geq \lambda_d(g)$$

denote the absolute values of the (generalized) eigenvalues of g . Also, let $g = g_{ss}g_u = g_u g_{ss}$ denote the Jordan-Chevalley decomposition, that is g_{ss} is semisimple, g_u is unipotent, and g_{ss}, g_u commute. We say that g is *elliptic* if it is semisimple and $\lambda_j(g) = 1$ for all j . Notice that if g is elliptic, then the cyclic group $\langle g \rangle$ generated by g is relatively compact in $\mathrm{SL}(d, \mathbb{K})$.

If $g \in \mathrm{SL}(d, \mathbb{K})$ and $\sigma_k(g) > \sigma_{k+1}(g)$, let $U_k(g) \in \mathrm{Gr}_k(\mathbb{K}^d)$ denote the subspace spanned by the k major axes of the ellipse $g(\mathbb{S}^{d-1})$. The following lemma relates the singular values along a sequence in $\mathrm{SL}(d, \mathbb{K})$ to the action of this sequence on the associated Grassmannian. We omit the proof as it is standard.

Lemma 2.2. *Suppose $V_0 \in \mathrm{Gr}_k(\mathbb{K}^d)$ and $W_0 \in \mathrm{Gr}_{d-k}(\mathbb{K}^d)$. If $\{g_n\}_{n \geq 1}$ is a sequence in $\mathrm{SL}(d, \mathbb{K})$ and $k \in \{1, \dots, d-1\}$, then the following are equivalent:*

- (1) *There are open sets $\mathcal{O} \subset \mathrm{Gr}_k(\mathbb{K}^d)$ and $\mathcal{O}' \subset \mathrm{Gr}_{d-k}(\mathbb{K}^d)$, such that $g_n(V) \rightarrow V_0$ for all $V \in \mathcal{O}$ and $g_n^{-1}(W) \rightarrow W_0$ for all $W \in \mathcal{O}'$.*
- (2) *$g_n(V) \rightarrow V_0$ for all $V \in \mathrm{Gr}_k(\mathbb{K}^d)$ transverse to W_0 .*
- (3) *$\lim_{n \rightarrow \infty} \frac{\sigma_k(g_n)}{\sigma_{k+1}(g_n)} = \infty$, $\lim_{n \rightarrow \infty} U_k(g_n) = V_0$, and $\lim_{n \rightarrow \infty} U_{d-k}(g_n^{-1}) = W_0$.*

Moreover, if $g_n = g^n$ for all n , then g is P_k -proximal if $V_0 \oplus W_0 = \mathbb{K}^d$, and weakly unipotent if $V_0 \subset W_0$ or $W_0 \subset V_0$.

We will also make use of the following elementary calculations, which we recall without proof.

Lemma 2.3. (1) *If $u = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, then for all $d \geq 1$ there exists $c = c(d) > 0$ so that*

$$\frac{1}{c} \leq \frac{\sigma_k(\tau_d(u^n))}{n^{d+1-2k}} \leq c$$

for all $n \in \mathbb{N}$ and $k \in \{1, \dots, d\}$.

- (2) *If $A, B \in \mathrm{SL}(d, \mathbb{K})$ and $C = BAB^{-1}$, then*

$$\sigma_k(C) \leq \frac{\sigma_1(B)}{\sigma_d(B)} \sigma_k(A)$$

for all $k \in \{1, \dots, d\}$.

3. ANOSOV REPRESENTATIONS INTO $\mathrm{SL}(d, \mathbb{K})$

Following Labourie [22] and Guichard-Weinhard [16] we define Anosov representations for geometrically finite Fuchsian groups into $\mathrm{SL}(d, \mathbb{K})$.

In [Appendix B](#), we will extend our definition to the setting of all semi-simple Lie groups with finite center. If $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a representation, let

$$E_\rho = \mathrm{U}(\Gamma) \times \mathbb{K}^d \quad \text{and} \quad \widehat{E}_\rho = \Gamma \backslash \mathrm{U}(\Gamma) \times \mathbb{K}^d$$

where $\gamma \in \Gamma$ acts on the first factor by γ and on the second factor by $\rho(\gamma)$. The projection map $p : E_\rho \rightarrow \mathrm{U}(\Gamma)$ descends to a vector bundle

$$\widehat{p} : \widehat{E}_\rho \rightarrow \widehat{\mathrm{U}}(\Gamma)$$

which is called the *flat bundle associated to ρ* . The geodesic flow on $U(\Gamma)$ extends to a flow on $E_\rho = U(\Gamma) \times \mathbb{K}^d$ whose action is trivial on the second factor. This in turn descends to a flow on \widehat{E}_ρ which covers the geodesic flow on $\widehat{U}(\Gamma)$. We use ϕ_t to denote all four of these flows.

We say that a continuous map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$ is

- (1) ρ -equivariant if $\rho(\gamma) \circ \xi = \xi \circ \gamma$ for all $\gamma \in \Gamma$,
- (2) transverse if $\xi^k(x) \oplus \xi^{d-k}(y) = \mathbb{K}^d$ for all distinct $x, y \in \Lambda(\Gamma)$.

Such a map induces a continuous ϕ_t -invariant spitting

$$E_\rho = \Theta^k \oplus \Xi^{d-k} \quad \text{where} \quad \Theta^k|_v = \xi^k(v^+) \quad \text{and} \quad \Xi^{d-k}|_v = \xi^{d-k}(v^-)$$

which descends to a continuous ϕ_t -invariant splitting

$$\widehat{E}_\rho = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}.$$

Definition 3.1. If $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $k \in \{1, \dots, d-1\}$, a representation $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is P_k -Anosov if:

- (1) There exists a ρ -equivariant, continuous, transverse map

$$\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$$

which induces a splitting $\widehat{E}_\rho = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$.

- (2) For some family of norms $\|\cdot\|_v$ on the fibers of \widehat{E}_ρ there exists $C > 1$ and $c > 0$ so that

$$\frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq C e^{-ct} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $t > 0$, $v \in \widehat{U}(\Gamma)$, $Y \in \widehat{\Theta}^k|_v$ and non-zero $Z \in \widehat{\Xi}^{d-k}|_v$.

We refer to ξ as a P_k -Anosov *limit map* of ρ .

Remark 3.2. Notice that a continuous family of norms $\|\cdot\|_{v \in \widehat{U}(\Gamma)}$ on the fibers of \widehat{E}_ρ lifts to a continuous family of norms $\|\cdot\|_{v \in U(\Gamma)}$ on \mathbb{K}^d which is ρ -equivariant in the following sense: if $v \in U(\Gamma)$, $Y \in \mathbb{K}^d$ and $\gamma \in \Gamma$, then

$$(2) \quad \|\rho(\gamma)(Y)\|_{\gamma(v)} = \|Y\|_v.$$

Conversely, any continuous family of norms satisfying Equation (2) descends to a continuous family of norms on the fibers of \widehat{E}_ρ .

If one prefers a bundle-theoretic definition, one can consider the vector bundle $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ over $\widehat{U}(\Gamma)$. Notice that since the splitting is flow-invariant, ϕ_t induces a flow on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ given by

$$f \mapsto \phi_t \circ f \circ \phi_{-t},$$

with some abuse of notation we use ϕ_t to denote this flow. Moreover, any norm on the fibers of \widehat{E}_ρ induces an operator norm on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$. We say that the flow ϕ_t on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ is *uniformly contracting* if there exists $C, c > 0$ so that

$$\|\phi_t(f)\|_{\phi_t(v)} \leq C e^{-ct} \|f\|_v$$

for all $t > 0$, $v \in \widehat{U}(\Gamma)$ and $f \in \text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)_v$. One may easily check that ρ is P_k -Anosov if and only if there exists a ρ -equivariant, continuous transverse map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$ so that the flow is uniformly contracting on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ with respect to an operator norm associated to a continuous family of norms on \widehat{E}_ρ . (The details of this equivalence are worked out carefully in the proof of [7, Prop. 2.3].) Moreover, by duality, ϕ_t is uniformly contracting on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ if and only if the flow, also called ϕ_t , on $\text{Hom}(\widehat{\Theta}^k, \widehat{\Xi}^{d-k})$ is uniformly expanding, i.e. the inverse flow ϕ_{-t} is uniformly contracting. We record these observations for future use.

Proposition 3.3. *Suppose that $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group, $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation and $k \in \{1, \dots, d-1\}$. Then the following are equivalent:*

- (1) ρ is P_k -Anosov,
(2) There exists a ρ -equivariant, continuous transverse map

$$\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \mathrm{Gr}_k(\mathbb{R}^d) \times \mathrm{Gr}_{d-k}(\mathbb{R}^d)$$

so that the flow is uniformly contracting on $\mathrm{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ with respect to an operator norm induced by a norm on \widehat{E}_ρ .

- (3) There exists a ρ -equivariant, continuous transverse map

$$\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \mathrm{Gr}_k(\mathbb{R}^d) \times \mathrm{Gr}_{d-k}(\mathbb{R}^d)$$

so that the flow is uniformly expanding on $\mathrm{Hom}(\widehat{\Theta}^k, \widehat{\Xi}^{d-k})$ with respect to an operator norm induced by a norm on \widehat{E}_ρ .

As an immediate corollary we obtain:

Corollary 3.4. *Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group, $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a representation and $k \in \{1, \dots, d-1\}$. Then ρ is P_k -Anosov if and only if ρ is P_{d-k} -Anosov.*

Remark 3.5. If we wish to allow Γ to be a geometrically finite subgroup of $\mathrm{PGL}(2, \mathbb{R})$, then we simply consider $\Gamma_0 = \Gamma \cap \mathrm{PSL}(2, \mathbb{R})$ and say that $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov if and only if $\rho|_{\Gamma_0}$ is P_k -Anosov. With this definition, all of our results for geometrically finite Fuchsian groups remain true for geometrically finite subgroups of $\mathrm{PGL}(2, \mathbb{R})$.

3.1. Canonical norms and cusp representations. It will be useful to construct certain ‘‘canonical’’ norms on the fibers of the vector bundles \widehat{E}_ρ , and hence also on $\mathrm{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$, that are well behaved on the cusps. Later we will show that when ρ is P_k -Anosov, the flow ϕ_t on $\mathrm{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ is uniformly contracting for any canonical norm (see Corollary 6.2).

The crucial property of our canonical norms is that they have a standard form over the thin part of the geodesic flow. In order to describe this standard form we will use the following result about representations of $\mathrm{SL}(2, \mathbb{R})$. Recall that $g = g_{ss}g_u$ denotes the multiplicative Jordan-Chevalley decomposition of $g \in \mathrm{SL}(d, \mathbb{K})$.

Proposition 3.6. *(see Appendix A) If $g \in \mathrm{SL}(d, \mathbb{K})$ is weakly unipotent, then there exists a representation $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ where $\Psi \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right) = g_u$ and g_{ss} commutes with the elements of $\Psi(\mathrm{SL}(2, \mathbb{R}))$.*

Proposition 3.6 follows easily from the Jordan normal form of a weakly unipotent matrix and we delay the proof until [Appendix A](#).

A representation $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ of a geometrically finite Fuchsian group is *type-preserving* if ρ sends every parabolic element in Γ to a weakly unipotent element in $\mathrm{SL}(d, \mathbb{K})$. If ρ is a type preserving representation and $\alpha \in \Gamma$ is parabolic, then we say that a representation $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a *cusp representation* for α and $\rho(\alpha)$ if

- (1) $\Psi(\tilde{\alpha}) = \rho(\alpha)_u$, where $\tilde{\alpha}$ is the (unique) unipotent lift of α to $\mathrm{SL}(2, \mathbb{R})$ and
- (2) $\rho(\alpha)_{ss}$ commutes with the elements of $\Psi(\mathrm{SL}(2, \mathbb{R}))$.

Proposition 3.6 implies that cusp representations always exist.

Suppose that Γ is a geometrically finite Fuchsian group, $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is type-preserving and $C = \langle \alpha \rangle \backslash H$ is an embedded cusp neighborhood where $\alpha \in \Gamma$ is parabolic. Further suppose that Ψ is a cusp representation for α and $\rho(\alpha)$, and $\|\cdot\|_v$ is a family of norms of the fibers $T^1 \mathbb{H}^2 \times \mathbb{K}^d \rightarrow T^1 \mathbb{H}^2$ such that

- (1) each $\|\cdot\|_v$ is $\rho(\alpha)_{ss}$ -invariant
- (2) $\|\Psi(g)Z\|_{g(v)} = \|Z\|_v$ for all $g \in \mathrm{SL}(2, \mathbb{R})$, $Z \in \mathbb{K}^d$ and $v \in T^1 \mathbb{H}^2$.

Such families are easy to construct: the group $K := \overline{\langle \rho(\alpha)_{ss}, \Psi(-\mathrm{id}_2) \rangle}$ is abelian and compact, so there exists a norm $\|\cdot\|_0$ which is K -invariant, then if we fix some $v_0 \in T^1 \mathbb{H}^2$, the family of norms defined by $\|Z\|_{g(v_0)} := \|\Psi(g)^{-1}Z\|_0$ has the desired properties.

Also, for such a family of norms

$$\|\rho(\alpha)Z\|_{\alpha(v)} = \|\Psi(\tilde{\alpha})\rho(\alpha)_{ss}Z\|_{\alpha(v)} = \|Z\|_v$$

for all $Z \in \mathbb{K}^d$ and $v \in T^1\mathbb{H}^2$. So this family descends (and restricts) to a family of norms on the fibers of \widehat{E}_ρ over $\widehat{U}(\Gamma)_C$ which we call a *canonical family of norms on the cusp neighborhood* C . Observe that if C' is an embedded cusp neighborhood properly contained in C , then $\widehat{U}(\Gamma)_C - \widehat{U}(\Gamma)_{C'}$ is precompact, and a canonical family of norms on the fibers of \widehat{E}_ρ over $\widehat{U}(\Gamma)_C$ is determined completely by its restriction to the fibers over $\widehat{U}(\Gamma)_C - \widehat{U}(\Gamma)_{C'}$.

Definition 3.7. Let $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ be a type preserving representation of a geometrically finite Fuchsian group. A continuous family of norms $\|\cdot\|_v$ on the fibers of \widehat{E}_ρ is *canonical* if there exists a full collection \mathcal{C} of embedded cusp neighborhoods for Γ , such that for all $C \in \mathcal{C}$, the restriction of the family of norms to the fibers over $\widehat{U}(\Gamma)_C$ is a canonical family of norms on C .

It is straightforward to construct a canonical family of norms on the fibers of \widehat{E}_ρ . One first chooses a full collection of embedded cusp neighborhoods \mathcal{C} . For each $C \in \mathcal{C}$, one chooses a canonical norm on the cusp neighborhood C . One then chooses any continuous norm on a compact neighborhood of the fibers over the thick part $\widehat{U}(\Gamma) - \widehat{U}(\Gamma)_C$. One may then use a cut off function to interpolate between the norms on their interface and obtain a family of norms on all of \widehat{E}_ρ which is canonical with respect to a full collection of cusp neighborhoods contained in \mathcal{C} .

We observe that there are uniform upper and lower bounds on the growth rate of a canonical norm with respect to the (lift of the) geodesic flow ϕ_t .

Lemma 3.8. *Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group, $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is type-preserving and $\|\cdot\|_v$ is a canonical family of norms on the bundle \widehat{E}_ρ . There exist $C_0 > 1, c_0 > 0$ so that if $v \in \widehat{U}(\Gamma)$, $t \in \mathbb{R}$ and $Z \in \widehat{E}_\rho|_v$, then*

$$\frac{1}{C_0} e^{-c_0|t|} \|Z\|_v \leq \|\phi_t(Z)\|_{\phi_t(v)} \leq C_0 e^{c_0|t|} \|Z\|_v.$$

Proof. For $v \in \widehat{U}(\Gamma)$, let

$$f(v) = \max \left\{ \|\phi_t(Z)\|_{\phi_t(v)} : |t| \leq 1, Z \in \widehat{E}_\rho|_v, \|Z\|_v = 1 \right\}.$$

Let \mathcal{C} be a full collection of embedded cusp neighborhoods such that for all $C \in \mathcal{C}$, $\|\cdot\|_v$ restricted to the fibers over each component of $\widehat{U}(\Gamma)_C$ is a canonical family of norms on C . For each $C \in \mathcal{C}$, let C' be an embedded cusp neighborhood such that

$$\bigcup_{|t| \leq 1} \phi_t(C') \subset C.$$

Then f is constant on each $\widehat{U}(\Gamma)_{C'}$. Further,

$$K = \widehat{U}(\Gamma) - \bigcup_{C \in \mathcal{C}} \widehat{U}(\Gamma)_{C'}$$

is compact. Hence

$$C_0 := \sup_{v \in \widehat{U}(\Gamma)} f(v) = \max_{v \in K} f(v)$$

is finite. Set $c_0 = \log C_0$. For any $t \in \mathbb{R}$, let n be the largest integer such that $n \leq |t|$. If $t > 0$, then

$$\|\phi_t(Z)\|_{\phi_t(v)} \leq C_0^n \|\phi_{t-n}(Z)\|_{\phi_{t-n}(v)} \leq e^{c_0 n} C_0 \|Z\|_v \leq C_0 e^{c_0|t|} \|Z\|_v.$$

On the other hand, if $t < 0$, then

$$\|\phi_t(Z)\|_{\phi_t(v)} \leq C_0^n \|\phi_{t+n}(Z)\|_{\phi_{t+n}(v)} \leq e^{c_0 n} C_0 \|Z\|_v \leq C_0 e^{c_0|t|} \|Z\|_v.$$

This proves the required upper bound. The lower bound is similar. \square

Remark 3.9. We will not need this for our work, but with a bit more effort, one can show that any two canonical families of norms on \widehat{E}_ρ are bilipschitz.

4. BASIC PROPERTIES OF ANOSOV REPRESENTATIONS

In this section we prove Theorem 1.1 which we restate here.

Theorem 4.1. *If $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov, then*

(1) *For any $z_0 \in \mathbb{H}^2$, there exists $A, a > 1$ so that if $\gamma \in \Gamma$, then*

$$\frac{1}{A} \exp\left(\frac{1}{a} d_{\mathbb{H}^2}(z_0, \gamma(z_0))\right) \leq \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq A \exp(ad_{\mathbb{H}^2}(z_0, \gamma(z_0))).$$

(2) *There exists $B, b > 1$ so that if $\gamma \in \Gamma$, then*

$$\frac{1}{B} \exp\left(\frac{1}{b} \ell(\gamma)\right) \leq \frac{\lambda_k(\rho(\gamma))}{\lambda_{k+1}(\rho(\gamma))} \leq B \exp(b\ell(\gamma))$$

where $\ell(\gamma)$ is the translation length of γ on \mathbb{H}^2 .

(3) *The P_k -Anosov limit map ξ_ρ is strongly dynamics-preserving and unique. In particular, if $\alpha \in \Gamma$ is parabolic, then $\rho(\alpha)$ is weakly unipotent, while if $\gamma \in \Gamma$ is hyperbolic, then $\rho(\gamma)$ is P_k -proximal.*

(4) *If $z_0 \in \mathbb{H}^2$ and x_0 is a point in the symmetric space $X_d(\mathbb{K})$ for $\mathrm{SL}(d, \mathbb{K})$, then the orbit map $\tau_\rho : \Gamma(z_0) \rightarrow X_d(\mathbb{K})$ given by $\tau_\rho(\gamma(z_0)) = \rho(\gamma)(x_0)$ is a quasi-isometric embedding.*

Before proving the theorem we note the following consequences which will be useful in [5].

Corollary 4.2. *Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov with P_k -Anosov limit map ξ_ρ .*

(1) *If $\{\gamma_n\}$ is a sequence in Γ with $\gamma_n \rightarrow x \in \Lambda(\Gamma)$, then $\lim U_k(\rho(\gamma_n)) = \xi_\rho(x)$.*

(2) *If $\alpha \in \Gamma$ is parabolic and $j \in \{1, \dots, d\}$, then there exists an integer $c(j, \alpha)$ and $C_j > 1$ so that*

$$\frac{1}{C_j} \leq \frac{\sigma_j(\rho(\alpha^n))}{n^{c(j, \alpha)}} \leq C_j \quad \text{for all } n \in \mathbb{N}.$$

Moreover,

$$c(k, \alpha) - c(k+1, \alpha) > 0.$$

Proof. Property (1) in Corollary 4.2 is known as the P_k -Cartan property and is an immediate consequence of the fact that ξ_ρ is strongly dynamics-preserving and Lemma 2.2.

If $\alpha \in \Gamma$ is parabolic, then part (3) of Theorem 1.1 implies that $\rho(\alpha)$ is weakly unipotent. So the group $\overline{\langle \rho(\alpha)_{ss} \rangle}$ is compact. Further, there exists $\{d_1, \dots, d_m\}$, so that $\rho(\alpha)_u$ is conjugate to $\oplus_{i=1}^m \tau_{d_i} \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right)$.

Thus there exists a constant $C_0 > 1$ such that

$$\frac{1}{C_0} \sigma_j \left(\oplus_{i=1}^m \tau_{d_i} \left(\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \right) \right) \leq \sigma_j(\rho(\alpha^n)) \leq C_0 \sigma_j \left(\oplus_{i=1}^m \tau_{d_i} \left(\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \right) \right)$$

for all $n \in \mathbb{N}$ and $j \in \{1, \dots, d\}$.

Given $j \in \Delta$ and $n \in \mathbb{N}$, there exists i and $k \in \{1, \dots, d_i\}$, so that

$$\sigma_j \left(\oplus_{i=1}^m \tau_{d_i} \left(\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \right) \right) = \sigma_k \left(\tau_{d_i} \left(\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \right) \right).$$

Then Lemma 2.3 implies the first claim in part (2) of Corollary 4.2, while the second claim follows from part (1) of Theorem 4.1. □

Proof of Theorem 4.1. Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov. By definition there exist $C > 1$, $c > 0$, and a ρ -equivariant family of norms $\|\cdot\|_v$ on the fibers of $E_\rho \rightarrow \mathrm{U}(\Gamma)$ so that

$$(3) \quad \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq C e^{-ct} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $t > 0$, $v \in \mathrm{U}(\Gamma)$, $Y \in \xi^k(v^+)$ and non-zero $Z \in \xi^{d-k}(v^-)$.

Fix a distance d_∞ on $\partial \mathbb{H}^2 \cong \mathbb{S}^1$ which is induced by a Riemannian metric. The following very special case of a result of Abels-Margulis-Soifer [1, Theorem 4.1] plays a key role in the proof.

Lemma 4.3. *There exist $\delta > 0$ and a finite subset \mathcal{B} of Γ such that if $\gamma \in \Gamma$, then there exists $\beta \in \mathcal{B}$ so that $\gamma\beta$ is hyperbolic and $d_\infty((\gamma\beta)^+, (\gamma\beta)^-) \geq \delta$.*

Lemma 4.3 allows us to reduce much of the proof to considering hyperbolic elements $\gamma \in \Gamma$ with $d_\infty(\gamma^+, \gamma^-) \geq \delta$. Since ξ is transverse and continuous these elements have the following decomposition: there exists a compact set $\mathcal{A} \subset \mathrm{SL}(d, \mathbb{K})$ so that if $\gamma \in \Gamma$ is hyperbolic and $d_\infty(\gamma^+, \gamma^-) \geq \delta$, then there exists $g_\gamma \in \mathcal{A}$, $A_\gamma \in \mathrm{GL}(k, \mathbb{K})$ and $B_\gamma \in \mathrm{GL}(d-k, \mathbb{K})$ with

$$(4) \quad \rho(\gamma) = g_\gamma \begin{pmatrix} A_\gamma & 0 \\ 0 & B_\gamma \end{pmatrix} g_\gamma^{-1}, \quad \xi^k(\gamma^+) = g_\gamma(\langle e_1, \dots, e_k \rangle) \quad \text{and} \quad \xi^{d-k}(\gamma^-) = g_\gamma(\langle e_{k+1}, \dots, e_d \rangle).$$

There also exists a compact set K of \mathbb{H}^2 so that any bi-infinite geodesic whose endpoints are a distance at least δ apart intersects K . Let

$$R = \max\{d_{\mathbb{H}^2}(i, z) : z \in K\}.$$

The next two lemmas establish (1) for any hyperbolic element γ with $d_\infty(\gamma^+, \gamma^-) \geq \delta$. The proof of the first lemma makes crucial use of the contraction properties of the flow, while the second lemma does not depend on the contraction properties.

Lemma 4.4. *There exist $C_1, C_2 > 0$ such that if $\gamma \in \Gamma$ is hyperbolic and $d_\infty(\gamma^+, \gamma^-) \geq \delta$, then*

$$\frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \geq C_1 e^{cd_{\mathbb{H}^2}(i, \gamma(i))}$$

and

$$\sigma_k(A_\gamma) \geq C_2 e^{cd_{\mathbb{H}^2}(i, \gamma(i))} \sigma_1(B_\gamma).$$

Proof. Recall that $r_v : \mathbb{R} \rightarrow \mathbb{H}^2$ denotes the geodesic with $r'_v(0) = v \in T^1 \mathbb{H}^2$. Since the geodesic joining γ^+ to γ^- intersects K , there exists $v_0 \in \mathrm{U}(\Gamma)$ with $r_{v_0}(0) \in K$ and $v_0^\pm = \gamma^\pm$. Also, notice that

$$\ell(\gamma) = d_{\mathbb{H}^2}(r_{v_0}(0), \gamma(r_{v_0}(0))) \geq d_{\mathbb{H}^2}(i, \gamma(i)) - 2R.$$

Since $\phi_{\ell(\gamma)}(v_0) = \gamma(v_0)$, the ρ -equivariance of the norms and Equation (3) imply that

$$\frac{\|Y\|_{v_0}}{\|Z\|_{v_0}} = \frac{\|\rho(\gamma)(Y)\|_{\gamma(v_0)}}{\|\rho(\gamma)(Z)\|_{\gamma(v_0)}} \leq C e^{-c\ell(\gamma)} \frac{\|\rho(\gamma)(Y)\|_{v_0}}{\|\rho(\gamma)(Z)\|_{v_0}}$$

for all $Y \in \xi^k(\gamma^+)$ and non-zero $Z \in \xi^{d-k}(\gamma^-)$.

Since K is compact, there exists L so that if $v \in \mathrm{U}(\Gamma)$ and $r_v(0) \in K$, then $\|\cdot\|_v$ is L -bilipschitz to the standard Euclidean norm $\|\cdot\|_2$ on \mathbb{K}^d . Therefore,

$$\frac{\|\rho(\gamma)(Y)\|_2}{\|\rho(\gamma)(Z)\|_2} \geq \frac{1}{CL^4} e^{c\ell(\gamma)} \frac{\|Y\|_2}{\|Z\|_2}$$

for all $Y \in \xi^k(\gamma^+)$ and non-zero $Z \in \xi^{d-k}(\gamma^-)$. So by the max-min/min-max characterization of singular values

$$\sigma_k(\rho(\gamma)) \geq \min_{\substack{Y \in \xi^k(\gamma^+) \\ Y \neq 0}} \frac{\|\rho(\gamma)(Y)\|_2}{\|Y\|_2} \geq \frac{1}{CL^4} e^{c\ell(\gamma)} \max_{\substack{Z \in \xi^{d-k}(\gamma^-) \\ Z \neq 0}} \frac{\|\rho(\gamma)(Z)\|_2}{\|Z\|_2} \geq \frac{1}{CL^4} e^{c\ell(\gamma)} \sigma_{k+1}(\rho(\gamma)).$$

Hence $C_1 := \frac{1}{CL^4} e^{-2Rc}$ suffices.

Since \mathcal{A} is compact,

$$S = \max \left\{ \frac{\sigma_1(g)}{\sigma_d(g)} : g \in \mathcal{A} \right\}$$

is finite. So, if $Y = (Y', 0) \in \mathbb{K}^k \times \{0\}$ and $Z = (0, Z') \in \{0\} \times \mathbb{K}^{d-k}$, then

$$\frac{\|A_\gamma(Y')\|_2}{\|B_\gamma(Z')\|_2} = \frac{\|g_\gamma^{-1} \rho(\gamma) g_\gamma(Y)\|_2}{\|g_\gamma^{-1} \rho(\gamma) g_\gamma(Z)\|_2} \geq \frac{1}{S} \frac{\|\rho(\gamma) g_\gamma(Y)\|_2}{\|\rho(\gamma) g_\gamma(Z)\|_2} \geq \frac{1}{CL^4 S} e^{c\ell(\gamma)} \frac{\|g_\gamma(Y)\|_2}{\|g_\gamma(Z)\|_2} \geq \frac{1}{CL^4 S^2} e^{c\ell(\gamma)} \frac{\|Y'\|_2}{\|Z'\|_2}$$

So, again by the max-min/min-max characterization of singular values,

$$\sigma_k(A_\gamma) \geq \min_{Y' \neq 0} \frac{\|A_\gamma(Y')\|_2}{\|Y'\|_2} \geq \frac{1}{CL^4 S^2} e^{c\ell(\gamma)} \max_{Z' \neq 0} \frac{\|B_\gamma(Z')\|_2}{\|Z'\|_2} \geq \frac{1}{CL^4 S^2} e^{c\ell(\gamma)} \sigma_1(B_\gamma).$$

Hence $C_2 := \frac{1}{CL^4 S^2} e^{-2Rc}$ suffices. \square

Lemma 4.5. *There exist $C_3, c_0 > 0$ so that if $\gamma \in \Gamma$ is hyperbolic and $d_\infty(\gamma^+, \gamma^-) \geq \delta$, then*

$$\sigma_1(\rho(\gamma)) \leq C_3 e^{c_0 d_{\mathbb{H}^2}(i, \gamma(i))}.$$

In particular,

$$\frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq C_3^2 e^{2c_0 d_{\mathbb{H}^2}(i, \gamma(i))}.$$

Proof. Fix a ρ -equivariant family of norms $\|\cdot\|_{v \in \mathbf{U}(\Gamma)}^*$ which descends to a canonical family of norms on \widehat{E}_ρ . Then by Lemma 3.8 there exist $C_0, c_0 > 0$ so that

$$(5) \quad \frac{1}{C_0} e^{-c_0 |t|} \|Y\|_v^* \leq \|Y\|_{\phi_t(v)}^* \leq C_0 e^{c_0 |t|} \|Y\|_v^*$$

for all $v \in \mathbf{U}(\Gamma)$, $Y \in \mathbb{K}^d$ and $t \in \mathbb{R}$.

Since the geodesic joining γ^+ to γ^- intersects K , there exists $v_0 \in \mathbf{U}(\Gamma)$ with $r_{v_0}(0) \in K$ and $v_0^\pm = \gamma^\pm$. Also, recall that

$$\ell(\gamma) = d_{\mathbb{H}^2}(r_{v_0}(0), \gamma(r_{v_0}(0))) \leq d_{\mathbb{H}^2}(i, \gamma(i)) + 2R.$$

Since $\phi_{\ell(\gamma)}(v_0) = \gamma(v_0)$, the ρ -equivariance of the norms and Equation (5) imply that

$$\|Y\|_{v_0}^* = \|\rho(\gamma)(Y)\|_{\gamma(v_0)}^* \geq \frac{1}{C_0} e^{-c_0 \ell(\gamma)} \|\rho(\gamma)(Y)\|_{v_0}^*$$

if $Y \in \mathbb{K}^d$.

Since K is compact, there exists L so that if $v \in \mathbf{U}(\Gamma)$ and $r_v(0) \in K$, then $\|\cdot\|_v^*$ is L -bilipschitz to the standard Euclidean norm $\|\cdot\|_2$ on \mathbb{K}^d . Therefore, if $Y \in \mathbb{K}^d$, then

$$\|\rho(\gamma)(Y)\|_2 \leq C_3 e^{c_0 d_{\mathbb{H}^2}(i, \gamma(i))} \|Y\|_2$$

where $C_3 := C_0 L^2 e^{2c_0 R}$. So

$$\sigma_1(\rho(\gamma)) \leq C_3 e^{c_0 d_{\mathbb{H}^2}(i, \gamma(i))}.$$

Finally, notice that

$$\frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq \frac{\sigma_1(\rho(\gamma))}{\sigma_d(\rho(\gamma))} = \sigma_1(\rho(\gamma)) \sigma_1(\rho(\gamma^{-1})) \leq C_3^2 e^{2c_0 d_{\mathbb{H}^2}(i, \gamma(i))}. \quad \square$$

We can now prove part (1). Since \mathcal{B} is finite, both

$$S_{\mathcal{B}} = \max \left\{ \frac{\sigma_1(\rho(\beta))}{\sigma_d(\rho(\beta))} \mid \beta \in \mathcal{B} \right\} \quad \text{and} \quad b = \max \{ d_{\mathbb{H}^2}(i, \beta(i)) : \beta \in \mathcal{B} \}$$

are finite.

Given any $\gamma \in \Gamma$, Lemma 4.3 implies that there exists $\beta \in \mathcal{B}$ so that $d_\infty((\gamma\beta)^+, (\gamma\beta)^-) \geq \delta$. Then

$$\frac{1}{S_{\mathcal{B}}} \frac{\sigma_k(\rho(\gamma\beta))}{\sigma_{k+1}(\rho(\gamma\beta))} \leq \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq S_{\mathcal{B}} \frac{\sigma_k(\rho(\gamma\beta))}{\sigma_{k+1}(\rho(\gamma\beta))}$$

and

$$|d_{\mathbb{H}^2}(i, \gamma(i)) - d_{\mathbb{H}^2}(i, \gamma\beta(i))| \leq d_{\mathbb{H}^2}(i, \beta(i)) \leq b.$$

Therefore,

$$\frac{1}{A} \exp\left(\frac{1}{a} d_{\mathbb{H}^2}(i, \gamma(i))\right) \leq \frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \leq A \exp(a d_{\mathbb{H}^2}(i, \gamma(i)))$$

where $A = \max\{\frac{1}{C_1} S_{\mathcal{B}} e^{cb}, C_3^2 S_{\mathcal{B}} e^{2c_0 b}\}$ and $a = \max\{1/c, 2c_0\}$. This proves (1).

Recall that if $T \in \mathrm{SL}(d, \mathbb{K})$, then

$$\lambda_j(T^n) = \lim_{n \rightarrow \infty} (\sigma_j(T^n))^{1/n}$$

and $\ell(\gamma) = \lim_{n \rightarrow \infty} \frac{d_{\mathbb{H}^2}(i, \gamma^n(i))}{n}$ for all $\gamma \in \Gamma$. Therefore, part (2) follows immediately from part (1). (One may give a direct proof of (2) in the spirit of Lemma 4.4 by noting that there is a compact subset \hat{K} of \mathbb{H}^2 such that every hyperbolic element of Γ is conjugate to an element of Γ whose axis passes through \hat{K} .)

Part (4) is a simple consequence of part (1) and Lemma 4.5. Recall that $X_d(\mathbb{R}) = \mathrm{SL}(d, \mathbb{R})/\mathrm{SO}(d)$ and $X_d(\mathbb{C}) = \mathrm{SL}(d, \mathbb{C})/\mathrm{SU}(d)$. We may choose x_0 to be either $[\mathrm{SO}(d)]$ or $[\mathrm{SU}(d)]$. Then, after possibly scaling, we have the following formula for the distance on $X_d(\mathbb{K})$, see for instance [8, Cor. 10.42],

$$d_{X_d}(x_0, g(x_0)) = \sqrt{\sum_{j=1}^d |\log \sigma_j(g)|^2}$$

Therefore, applying part (1), we see that

$$d_{X_d}(x_0, \rho(\gamma)(x_0)) \geq \frac{1}{2} \log \left(\frac{\sigma_k(\rho(\gamma))}{\sigma_{k+1}(\rho(\gamma))} \right) \geq \frac{a}{2} d_{\mathbb{H}^2}(i, \gamma(i)) - \frac{\log A}{2}$$

for all $\gamma \in \Gamma$. On the other hand, by Lemma 4.5

$$d_{X_d}(x_0, \rho(\gamma)(x_0)) \leq 2 \log \left(\frac{\sigma_1(\rho(\gamma))}{\sigma_d(\rho(\gamma))} \right) \leq 4 \log C_3 + 4c_0 d_{\mathbb{H}^2}(i, \gamma(i)),$$

so the orbit map is a quasi-isometry, which completes the proof of part (4).

We now show that ξ is strongly dynamics preserving. Notice that this immediately implies that ξ is unique. Lemma 2.2 will then imply that if $\alpha \in \Gamma$ is parabolic, then $\rho(\alpha)$ is weakly unipotent, while if $\gamma \in \Gamma$ is hyperbolic, then $\rho(\gamma)$ is P_k -proximal.

Fix a sequence $\{\gamma_n\}$ with $\gamma_n \rightarrow x \in \Lambda(\Gamma)$ and $\gamma_n^{-1} \rightarrow y \in \Lambda(\Gamma)$. For each n there exists $\beta_n \in \mathcal{B}$ such that $\gamma_n \beta_n$ is hyperbolic and

$$d_{\infty}((\gamma_n \beta_n)^+, (\gamma_n \beta_n)^-) \geq \delta.$$

Since the set \mathcal{B} is finite, we can divide $\{\gamma_n\}$ into finitely many subsequences and only consider the case when $\beta_n = \beta$ for all n . Then $(\gamma_n \beta)^+ \rightarrow x$ and $(\gamma_n \beta)^- \rightarrow \beta^{-1}(y)$.

Let

$$\rho(\gamma_n \beta) = g_n \begin{pmatrix} A_n & 0 \\ 0 & B_n \end{pmatrix} g_n^{-1}$$

be the block diagonal decomposition from Equation (4). Then by Lemma 4.4

$$\lim_{n \rightarrow \infty} \frac{\sigma_1(B_n)}{\sigma_k(A_n)} = 0.$$

so

$$g_n^{-1} \rho(\gamma_n \beta) g_n(W) \rightarrow \langle e_1, \dots, e_k \rangle$$

for all $W \in \mathrm{Gr}_k(\mathbb{K}^d)$ transverse to $\langle e_{k+1}, \dots, e_d \rangle$. Further, by construction

$$\xi^k((\gamma_n \beta)^+) = g_n(\langle e_1, \dots, e_k \rangle) \quad \text{and} \quad \xi^{d-k}((\gamma_n \beta)^-) = g_n(\langle e_{k+1}, \dots, e_d \rangle).$$

So by the continuity of ξ ,

$$\rho(\gamma_n \beta)(V) \rightarrow \xi^k(x)$$

for all V transverse to $\xi^{d-k}(\beta^{-1}(y))$. This implies that

$$\rho(\gamma_n)(V) \rightarrow \xi^k(x)$$

for all V transverse to $\xi^{d-k}(y) = \rho(\beta)(\xi^{d-k}(\beta^{-1}(y)))$ which completes the proof of (3). \square

5. BASIC PROPERTIES OF CUSP REPRESENTATIONS

In this section we establish some useful properties of the cusp representations associated to type preserving representations introduced in Section 3.1.

We say that a representation $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -proximal if $\Psi(\gamma)$ is P_k -proximal for some (any) hyperbolic element $\gamma \in \mathrm{SL}(2, \mathbb{R})$. We first observe that a cusp representation associated to a P_k -Anosov representation is P_k -proximal.

Proposition 5.1. *Suppose that $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{K})$ is P_k -Anosov. If $\alpha \in \Gamma$ is parabolic and $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a cusp representation associated to α and $\rho(\alpha)$, then Ψ is P_k -proximal.*

Proof. Let $\tilde{\alpha} \in \mathrm{SL}(2, \mathbb{R})$ be the unique unipotent lift of α . Since $\rho(\alpha)$ is weakly unipotent, the group $\overline{\langle \rho(\alpha)_{ss} \rangle}$ is compact. Then, since $\Psi(\tilde{\alpha}^n) = \rho(\alpha)_{ss}^{-n} \rho(\alpha^n)$, there exists $L > 1$ such that

$$\frac{1}{L} \sigma_j(\rho(\alpha^n)) \leq \sigma_j(\Psi(\tilde{\alpha}^n)) \leq L \sigma_j(\rho(\alpha^n))$$

for all $j \in \{1, \dots, d\}$ and $n \in \mathbb{Z}$. Then, since ρ is strongly dynamics-preserving, Lemma 2.2 implies that

$$\lim_{n \rightarrow \infty} \frac{\sigma_k(\Psi(\tilde{\alpha}^n))}{\sigma_{k+1}(\Psi(\tilde{\alpha}^n))} \geq \frac{1}{L^2} \lim_{n \rightarrow \infty} \frac{\sigma_k(\rho(\alpha^n))}{\sigma_{k+1}(\rho(\alpha^n))} = \infty.$$

Now, write $\tilde{\alpha}^n = k_n a_{t_n} m_n$, where $k_n, m_n \in \mathrm{SO}(2)$ and $a_t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$. Since $\{\Psi(a_t) : t \in \mathbb{R}\}$ is simultaneously diagonalizable, by increasing $L > 1$ we may assume that

$$\frac{1}{L} \sigma_k(\Psi(a_t)) \leq \lambda_k(\Psi(a_t)) \leq L \sigma_k(\Psi(a_t))$$

for all $t \in \mathbb{R}$. Since $\Psi(\mathrm{SO}(2))$ is compact, we may increase $L > 1$ further and assume that

$$\frac{1}{L} \leq \sigma_j(\Psi(g)) \leq L$$

for all $g \in \mathrm{SO}(2)$ and $j \in \{1, \dots, d\}$.

Then

$$\lim_{n \rightarrow \infty} \frac{\lambda_k(\Psi(a_{t_n}))}{\lambda_{k+1}(\Psi(a_{t_n}))} \geq \frac{1}{L^2} \lim_{n \rightarrow \infty} \frac{\sigma_k(\Psi(a_{t_n}))}{\sigma_{k+1}(\Psi(a_{t_n}))} \geq \frac{1}{L^6} \lim_{n \rightarrow \infty} \frac{\sigma_k(\Psi(\tilde{\alpha}^n))}{\sigma_{k+1}(\Psi(\tilde{\alpha}^n))} = \infty,$$

which implies that $\Psi(a_t)$ is P_k -proximal for all $t \in \mathbb{R} - \{0\}$. \square

Next we observe that a P_k -proximal representation is itself P_k -Anosov and admits a P_k -limit map, in the following sense.

Proposition 5.2. *If $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a P_k -proximal representation, then there exists a continuous, Ψ -equivariant transverse map*

$$\eta = (\eta^k, \eta^{d-k}) : \partial \mathbb{H}^2 \rightarrow \mathrm{Gr}_k(\mathbb{K}^d) \times \mathrm{Gr}_{d-k}(\mathbb{K}^d)$$

with the following properties:

- (1) *If $\|\cdot\|$ is a Ψ -invariant family of norms on the fibers of $T^1 \mathbb{H}^2 \times \mathbb{K}^d$, then there exists $B, b > 0$ so that if $t > 0$, $v \in T^1 \mathbb{H}^2$, $Y \in \eta^k(v^+)$ and $Z \in \eta^{d-k}(v^-)$ is non-zero, then*

$$(6) \quad \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq B e^{-bt} \frac{\|Y\|_v}{\|Z\|_v}.$$

(2) If $\{\gamma_n\}$ is a sequence in $\mathrm{SL}(2, \mathbb{R})$ with $\gamma_n \rightarrow x \in \partial \mathbb{H}^2$ and $\gamma_n^{-1} \rightarrow y \in \partial \mathbb{H}^2$, then

$$\Psi(\gamma_n)(V) \rightarrow \eta^k(x)$$

locally uniformly for all $V \in \mathrm{Gr}_k(\mathbb{K}^d)$ transverse to $\eta^{d-k}(y)$.

(3) If $g \in \mathrm{SL}(d, \mathbb{K})$ commutes with the elements of $\Psi(\mathrm{SL}(2, \mathbb{R}))$, then $g \circ \eta = \eta$.

Proof. By conjugating Ψ we can assume that

$$(7) \quad \Psi = \bigoplus_{i=1}^m \tau_{d_i}.$$

Let $\xi_{d_i} : \partial \mathbb{H}^2 \rightarrow \mathcal{F}(\mathbb{K}^{d_i})$ denote the τ_{d_i} -equivariant boundary map described in Section 2.3.

By definition, if $\gamma \in \mathrm{SL}(2, \mathbb{R})$ is a hyperbolic element, then $\tau_{d_i}(\gamma)$ is diagonalizable with eigenvalues having pairwise distinct absolute values. Furthermore, for all $k = 1, \dots, d_i - 1$, $\xi_{d_i}^k(\gamma^+)$ is the direct sum of the eigenspaces of the k largest eigenvalues of $\tau_{d_i}(\gamma)$.

First, we construct the map η . Observe that since Ψ is P_k -proximal, $\Psi(\gamma)$ is P_k -proximal and P_{d-k} -proximal for every hyperbolic element $\gamma \in \mathrm{SL}(2, \mathbb{R})$. Thus, for all $i = 1, \dots, m$, there are integers $k_i \in \{0, \dots, d_i\}$ such that

- $\sum_{i=1}^m k_i = k$, and for all hyperbolic $\gamma \in \mathrm{SL}(2, \mathbb{R})$, $\bigoplus_{i=1}^m \xi_{d_i}^{k_i}(\gamma^+) \in \mathrm{Gr}_k(\mathbb{K}^d)$ is the attracting fixed point for the action of $\Psi(\gamma)$ on $\mathrm{Gr}_k(\mathbb{K}^d)$, and
- $\sum_{i=1}^m d - k_i = d - k$, and for all hyperbolic $\gamma \in \mathrm{SL}(2, \mathbb{R})$, $\bigoplus_{i=1}^m \xi_{d_i}^{d_i - k_i}(\gamma^-) \in \mathrm{Gr}_{d-k}(\mathbb{K}^d)$ is the attracting fixed point for the action of $\Psi(\gamma^{-1})$ on $\mathrm{Gr}_{d-k}(\mathbb{K}^d)$.

Then set

$$\eta^k := \bigoplus_{i=1}^m \xi_{d_i}^{k_i} : \partial \mathbb{H}^2 \rightarrow \mathrm{Gr}_k(\mathbb{K}^d)$$

and

$$\eta^{d-k} := \bigoplus_{i=1}^m \xi_{d_i}^{d_i - k_i} : \partial \mathbb{H}^2 \rightarrow \mathrm{Gr}_{d-k}(\mathbb{K}^d).$$

Note that this pair of maps are continuous, Ψ -equivariant, and transverse.

For every $v \in T^1 \mathbb{H}^2$, there exists a one-parameter subgroup $\{a_t\}_{t \in \mathbb{R}}$ of hyperbolic elements in $\mathrm{SL}(2, \mathbb{R})$ so that $a_t(v) = \phi_t(v)$ for all $t \in \mathbb{R}$. Since $\Psi(a_1)$ is P_k -proximal,

$$b := \log \left(\frac{\lambda_k(\Psi(a_1))}{\lambda_{k+1}(\Psi(a_1))} \right) > 0,$$

so

$$\frac{\lambda_k(\Psi(a_t))}{\lambda_{k+1}(\Psi(a_t))} = e^{bt}$$

for all $t > 0$.

If $X \in \mathbb{K}^d$, then

$$\|X\|_{\phi_t(v)} = \|\Psi(a_t)^{-1}(X)\|_v$$

so, if $Y \in \eta^k(v^+)$ and $Z \in \eta^{d-k}(v^-)$ are non-zero, then

$$\frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq \frac{\lambda_{k+1}(\Psi(a_t))}{\lambda_k(\Psi(a_t))} \frac{\|Y\|_v}{\|Z\|_v} = e^{-bt} \frac{\|Y\|_v}{\|Z\|_v}$$

which completes the proof of (1).

If $\{\gamma_n\}$ is a sequence in $\mathrm{SL}(2, \mathbb{R})$ with $\gamma_n \rightarrow x$ and $\gamma_n^{-1} \rightarrow y$, one can write $\gamma_n = k_n a_{t_n} m_n$ where $k_n, m_n \in \mathrm{SO}(2)$, $t_n \geq 0$, and $a_t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$. By assumption, $t_n \rightarrow \infty$, $m_n \rightarrow m$ and $k_n \rightarrow k$ with $k(\infty) = x$ and $m^{-1}(0) = y$.

Notice that if $W \in \mathrm{Gr}_k(\mathbb{R}^d)$ is transverse to $\eta^{d-k}(0)$, then $\{\Psi(a_{t_n})(W)\}$ converges to $\eta^k(\infty)$ (since, by definition, $\eta^k(\infty)$ is the attracting k -plane of $\Psi(a_1)$ and $\eta^{d-k}(0)$ is the repelling $(d-k)$ -plane). So, by equivariance, if V is transverse to $\eta^{d-k}(y)$, then $\Psi(m_n)(V)$ is transverse to $\eta^{d-k}(0) = \lim \Psi(m_n)(\eta^{d-k}(y))$ for all large enough n . Thus, $\{\Psi(a_{t_n} m_n)(V)\}$ converges to $\eta^k(\infty)$ locally uniformly in V . Therefore, $\{\Psi(\gamma_n)(V)\} = \{\Psi(k_n a_{t_n} m_n)(V)\}$ converges to $\Psi(k)(\eta^k(\infty)) = \eta^k(x)$ locally uniformly in V . This proves (2).

To prove (3), fix $x \in \partial\mathbb{H}^2$ and a hyperbolic element $\gamma \in \mathrm{SL}(2, \mathbb{R})$ so that $\gamma^+ = x$. Then $\Psi(\gamma)$ is P_k -proximal and $\eta^k(x) \in \mathrm{Gr}_k(\mathbb{K}^d)$ is the attracting fixed point of $\Psi(\gamma)$, so $g\eta^k(x) = \eta^k(x)$ since g commutes with $\Psi(\gamma)$. Similar reasoning shows that $g\eta^{d-k}(x) = \eta^{d-k}(x)$. \square

The following technical result says that the image of the limit map of an Anosov representation is asymptotically homogeneous at a parabolic fixed point. In one of the arguments that follow, we need uniform control over continuous families of Anosov representations, so we introduce a parameter u .

Proposition 5.3. *Suppose that $\alpha \in \mathrm{SL}(2, \mathbb{R})$ is a parabolic element, X is a closed α -invariant subset of $\partial\mathbb{H}^2$ containing the fixed point $\alpha^+ = \alpha^-$ of α , $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ is a P_k -proximal representation with P_k -limit map η , and $\ell \in \mathrm{SL}(d, \mathbb{K})$ is elliptic and commutes with the elements of $\Psi(\mathrm{SL}(2, \mathbb{R}))$.*

Let U be a compact metric space and $\{g_u\}_{u \in U}$ be a continuous family of elements in $\mathrm{SL}(d, \mathbb{R})$. Suppose

$$\xi = (\xi^k, \xi^{d-k}) : U \times X \rightarrow \mathrm{Gr}_k(\mathbb{K}^d) \times \mathrm{Gr}_{d-k}(\mathbb{K}^d)$$

is continuous and for each $u \in U$, the map $\xi_u := \xi(u, \cdot)$ is transverse, $\xi_u(\alpha^+) = g_u\eta(\alpha^+)$ and

$$(8) \quad \xi_u \circ \alpha = g_u \ell \Psi(\alpha) g_u^{-1} \circ \xi_u.$$

If $\gamma \in \mathrm{SL}(2, \mathbb{R})$ is a hyperbolic element with attracting fixed point α^+ , then

$$\lim_{n \rightarrow \infty} (g_u \Psi(\gamma)^{-n} g_u^{-1} \circ \xi_u \circ \gamma^n)(x_n) = g_u \eta(x)$$

if $\{x_n\} \subset X$ and $\lim x_n = x \in \partial\mathbb{H}^2$. Moreover, the convergence is uniform in $u \in U$.

Proof. Fix a distance d_G on $\mathrm{Gr}_k(\mathbb{K}^d) \times \mathrm{Gr}_{d-k}(\mathbb{K}^d)$ induced by a Riemannian metric.

Suppose the proposition fails for a sequence $\{x_n\} \subset X$ with $\lim x_n = x \in X$. Then there exist $\epsilon > 0$, $\{n_j\}$ converging to infinity and a sequence $\{u_j\}$ in U such that

$$d_G \left((g_{u_j} \Psi(\gamma)^{-n_j} g_{u_j}^{-1} \circ \xi_{u_j} \circ \gamma^{n_j})(x_{n_j}), g_{u_j} \eta(x) \right) > \epsilon$$

for all j . Passing to subsequences we may suppose that $u_j \rightarrow u_\infty \in U$.

For notational convenience, let

$$\eta_j = g_{u_j} \Psi(\gamma)^{-n_j} g_{u_j}^{-1} \circ \xi_{u_j} \circ \gamma^{n_j}$$

for all $j \in \mathbb{N}$. Since η is Ψ -equivariant and $\xi_u(\alpha^+) = g_u \eta(\alpha^+)$ for all $u \in U$,

$$\begin{aligned} g_{u_j} \eta(\alpha^+) &= g_{u_j} \Psi(\gamma)^{-n_j} \eta(\alpha^+) = g_{u_j} \Psi(\gamma)^{-n_j} g_{u_j}^{-1} \circ \xi_{u_j}(\alpha^+) \\ &= g_{u_j} \Psi(\gamma)^{-n_j} g_{u_j}^{-1} \circ \xi_{u_j} \circ \gamma^{n_j}(\alpha^+) = \eta_j(\alpha^+). \end{aligned}$$

So, by passing to a tail of our sequences we may assume that $x_{n_j} \neq \alpha^+$ for every j .

First suppose that $\{y_j = \gamma^{n_j}(x_{n_j})\}$ lies in a compact subset of $\partial\mathbb{H}^2 - \{\alpha^+\}$. Then $x = \gamma^-$. Since $\xi_{u_j} \rightarrow \xi_{u_\infty}$, there exists $N > 0$ sufficiently large so that $\{g_{u_j}^{-1} \xi_{u_j}(y_j)\}_{n \geq N}$ lies in a compact subset of flags transverse to $g_{u_\infty}^{-1} \xi_{u_\infty}(\alpha^+) = \eta(\gamma^+)$. Hence Proposition 5.2 part (2) implies that

$$g_{u_j}^{-1} \eta_j(x_{n_j}) = \Psi(\gamma)^{-n_j} (g_{u_j}^{-1} \xi_{u_j}(y_j))$$

converges to $\eta(\gamma^-)$. Thus

$$\epsilon \leq \liminf_{j \rightarrow \infty} d_G(\eta_j(x_{n_j}), g_{u_j} \eta(x)) = d_G(g_{u_\infty} \eta(\gamma^-), g_{u_\infty} \eta(\gamma^-)) = 0$$

and we have a contradiction.

Now suppose that $\{y_j\}$ does not lie in a compact subset of $\partial\mathbb{H}^2 - \{\alpha^+\}$. We may assume without loss of generality that $\gamma^- = 0$, $\gamma^+ = \infty = \alpha^+$, and $\alpha = u_1$ where

$$u_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}(2, \mathbb{R}).$$

Then $y_j \in \mathbb{R} \subset \partial\mathbb{H}^2$ for all j . Let $z_j = \lfloor y_j \rfloor \in \mathbb{Z}$ and $w_j = y_j - z_j \in [0, 1]$ and set $\delta_j = \gamma^{-n_j} u_{z_j} = \gamma^{-n_j} \alpha^{z_j}$. Notice that $\alpha^{z_j}(w_j) = y_j$, $\delta_j(w_j) = x_{n_j}$, $\delta_j^- = \infty$ and $\delta_j^+ \rightarrow x$. Passing to a subsequence, we can suppose

that $w_j \rightarrow w \in [0, 1]$. Proposition 5.2 part (2) then implies that $\Psi(\delta_j)(V)$ converges to $\eta^k(x)$ locally uniformly for all $V \in \text{Gr}_k(\mathbb{K}^d)$ which are transverse to $\eta^{d-k}(\infty)$. Also,

$$\begin{aligned} \eta_j^k(x_{n_j}) &= g_{u_j} \Psi(\gamma^{-n_j}) g_{u_j}^{-1} (\xi_{u_j}^k(\alpha^{z_j}(w_j))) = g_{u_j} \Psi(\gamma^{-n_j}) g_{u_j}^{-1} (g_{u_j} \ell^{z_j} \Psi(\alpha^{z_j}) g_{u_j}^{-1} \xi_{u_j}^k(w_j)) \\ &= g_{u_j} \ell^{z_j} \Psi(\gamma^{-n_j} \alpha^{z_j}) g_{u_j}^{-1} (\xi_{u_j}^k(w_j)) = g_{u_j} \ell^{z_j} \Psi(\delta_j) (g_{u_j}^{-1} \xi_{u_j}^k(w_j)) \end{aligned}$$

(where in the first line we apply assumption (8)). We may pass to a subsequence so that $\ell^{z_j} \rightarrow \ell_\infty \in \text{SL}(d, \mathbb{K})$. Then ℓ_∞ is also elliptic and commutes with the elements of $\Psi(\text{SL}(2, \mathbb{R}))$ and hence, by Proposition 5.2 part (3), fixes each element in the image of η . Then

$$\lim_{j \rightarrow \infty} \eta_j^k(x_{n_j}) = g_{u_\infty} \ell_\infty \eta^k(x) = g_{u_\infty} \eta^k(x)$$

since $g_{u_j}^{-1} \xi_{u_j}^k(w_j) \rightarrow g_{u_\infty}^{-1} \xi_{u_\infty}^k(w)$, $g_{u_\infty}^{-1} \xi_{u_\infty}^k(w)$ is transverse to $\eta^{d-k}(\infty) = g_{u_\infty}^{-1} \xi_{u_\infty}^k(\infty)$, and $\ell_\infty \circ \eta = \eta$.

Reversing the roles of k and $d - k$, we may similarly show that $\eta_j^{d-k}(x_{n_j}) \rightarrow g_{u_\infty} \eta^{d-k}(x)$. Hence we again have a contradiction. \square

6. A DYNAMICAL CHARACTERIZATION OF LINEAR ANOSOV REPRESENTATIONS

In this section, we prove Theorem 1.2. The forward implication has already been established as part (3) of Theorem 1.1. The reverse implication follows from the following more general statement.

First, recall from Section 3 that a transverse, ρ -equivariant, continuous map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$ induces a continuous decomposition of \widehat{E}_ρ into a pair of sub-bundles $\widehat{E}_\rho = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$ of rank k and $d - k$ respectively. Further, the flow ϕ_t induces a flow on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$, which we denote by ϕ_t , and any canonical norm on \widehat{E}_ρ induces a canonical norm on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ which is simply given by the associated operator norm.

Theorem 6.1. *Suppose $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation. If there exists a ρ -equivariant, transverse, continuous, strongly dynamics preserving map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$, then ρ is type-preserving and the flow ϕ_t on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ is uniformly contracting with respect to any canonical norm on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$. In particular, ρ is P_k -Anosov and ξ is its P_k -Anosov limit map.*

The following is an immediate corollary of Theorem 6.1 and Theorem 4.1.

Corollary 6.2. *Suppose $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a P_k -Anosov representation. Then the flow ϕ_t on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ is uniformly contracting with respect to any canonical norm on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$.*

As another corollary, we see that if ρ is Zariski dense, then ρ is P_k -Anosov if it admits a transverse limit map, which generalizes a result of Guichard and Wienhard from the uncusped Anosov setting [16, Theorem 4.11].

Corollary 6.3. *Suppose $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation.*

- (1) *If ρ is irreducible, and there exists a ρ -equivariant, transverse, continuous, map $\xi = (\xi^1, \xi^{d-1}) : \Lambda(\Gamma) \rightarrow \text{Gr}_1(\mathbb{K}^d) \times \text{Gr}_{d-1}(\mathbb{K}^d)$, then ρ is P_1 -Anosov and ξ is its P_1 -Anosov limit map.*
- (2) *If $\wedge^k \rho : \Gamma \rightarrow \text{SL}(\wedge^k \mathbb{K}^d)$ is irreducible (e.g. if $\rho(\Gamma)$ is Zariski dense in $\text{SL}(d, \mathbb{K})$) and there exists a ρ -equivariant, transverse, continuous, map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$, then ρ is P_k -Anosov and ξ is its P_k -Anosov limit map.*

Proof. (1): It is enough to show that ξ is strongly dynamics preserving. Fix an escaping sequence $\{\gamma_n\}$ in Γ with $\gamma_n \rightarrow x$ and $\gamma_n^{-1} \rightarrow y$. Let $[\rho(\gamma_n)]$ denote the image of $\rho(\gamma_n)$ in $\mathbb{P}(\text{End}(\mathbb{K}^d))$. Then it is enough to show that $[\rho(\gamma_n)]$ converges to the element $T \in \mathbb{P}(\text{End}(\mathbb{K}^d))$ with $\ker(T) = \xi^{d-1}(y)$ and $\text{Image}(T) = \xi^1(x)$. Since $\mathbb{P}(\text{End}(\mathbb{K}^d))$ is compact it is enough to show that every convergent subsequence of $\rho(\gamma_n)$ converges to T . So suppose that $[\rho(\gamma_n)] \rightarrow S$ in $\mathbb{P}(\text{End}(\mathbb{K}^d))$. Then

$$S(v) = \lim_{n \rightarrow \infty} \rho(\gamma_n)(v)$$

for all $v \in \mathbb{P}(\mathbb{K}^d) \setminus \mathbb{P}(\ker S)$.

We first claim that $\text{Image}(S) = \xi^1(x)$. Since $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is irreducible, there exists $x_1, \dots, x_d \in \Lambda(\Gamma)$ so that $\xi(x_1), \dots, \xi(x_d)$ spans \mathbb{K}^d . Since $\partial\Gamma$ is perfect, we can perturb each x_j and assume that

$$y \notin \{x_1, \dots, x_d\}.$$

Then $\rho(\gamma_n)(\xi^1(x_j)) \rightarrow \xi^1(x)$. Since $\{\xi^1(x_1), \dots, \xi^1(x_d)\}$ spans \mathbb{K}^d , we can relabel and suppose that

$$\ker S \oplus \xi^1(x_1) \oplus \dots \oplus \xi^1(x_m) = \mathbb{K}^d$$

where $m = d - \dim \ker S$. Then

$$S(\xi^1(x_j)) = \lim_{n \rightarrow \infty} \rho(\gamma_n)(\xi^1(x_j)) = \xi^1(x)$$

for all $1 \leq j \leq m$. Hence $\text{Image}(S) = \xi^1(x)$.

To compute the kernel, we notice that $\text{Gr}_{d-1}(\mathbb{K}^d)$ may be identified with $\mathbb{P}(\mathbb{K}^{d*})$ by identifying a hyperplane Q in \mathbb{K}^d with the projective class of linear functionals with kernel Q . Notice that $[{}^t\rho(\gamma_n)]$ converges to tS in $\mathbb{P}(\text{End}(\mathbb{K}^{d*}))$. Repeating the argument above shows that $\text{Image}({}^tS) = \xi^{d-1}(y)$, so the kernel of S is $\xi^{d-1}(y)$.

(2): One can argue similarly using the Plücker embeddings. \square

Proof of Theorem 6.1. Suppose that $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group, $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is a representation and that

$$\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$$

is a continuous, transverse, ρ -equivariant, strongly dynamics-preserving map. Lemma 2.2 implies that ρ is type-preserving.

Let $\|\cdot\|$ be a canonical family of norms on \widehat{E}_ρ and let \mathcal{C} be a full collection of embedded cusp neighborhoods so that the restriction to the fibers over $\widehat{U}(\Gamma)_C$ is canonical for all $C \in \mathcal{C}$. We will also use $\|\cdot\|$ to denote the lift of $\|\cdot\|$ to a continuous family of norms on the fibers of $\text{U}(\Gamma) \times \mathbb{K}^d$.

The proof divides into two parts. We first use properties of the canonical family of norms to control the flow over the thin part $\widehat{U}(\Gamma)_C$. We then use a compactness argument to control the flow on the complement.

Proposition 6.4. *If $C \in \mathcal{C}$, then there exist constants b_C and B_C and an embedded cusp sub-neighborhood $C' \subset C$ such that if $v \in \widehat{U}(\Gamma)_{C'}$, $t \geq 0$ and $\phi_s(v) \in \widehat{U}(\Gamma)_{C'}$ for all $s \in [0, t]$, then*

$$(9) \quad \frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq B_C e^{-b_C t} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $Y \in \widehat{\Theta}^k|_v$ and non-zero $Z \in \widehat{\Xi}^{d-k}|_v$.

Proof. Suppose that $C = \langle \alpha \rangle \backslash H$. Then it suffices to find a horodisc $H' \subset H$ and constants b_C, B_C such that: if $v \in \text{U}(\Gamma)_{H'}$, $t \geq 0$ and $\phi_s(v) \in \text{U}(\Gamma)_{H'}$ for all $s \in [0, t]$, then

$$(10) \quad \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq B_C e^{-b_C t} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $Y \in \xi^k(v^+)$ and non-zero $Z \in \xi^{d-k}(v^-)$.

After possibly replacing C with a subcusp, there exists a cusp representation $\Psi : \text{SL}(2, \mathbb{R}) \rightarrow \text{SL}(d, \mathbb{K})$ for α and $\rho(\alpha)$ such that $\|\cdot\|$ coincides with a $\rho(\alpha)_{ss}$ -invariant, Ψ -equivariant family of norms on $\text{U}(\Gamma)_H$.

Let η be the P_k -limit map of Ψ . Proposition 5.2 implies that there exists $B, b > 0$ such that

$$(11) \quad \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq B e^{-bt} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $t \geq 0$, $v \in \text{U}(\Gamma)_H$ with $\phi_t(v) \in \text{U}(\Gamma)_H$, $Y \in \eta^k(v^+)$ and non-zero $Z \in \eta^{d-k}(v^-)$. Choose $b_C = \frac{b}{2}$ and $T > 0$ so that

$$(12) \quad B e^{-bt} < e^{-b_C t} \text{ for all } t > T.$$

We claim that there is a horodisk $H' \subset H$ so that if $v \in \mathbf{U}(\Gamma)_{H'}$, $t \in [T, 2T]$ and $\phi_s(v) \in \mathbf{U}(\Gamma)_{H'}$ for all $s \in [0, t]$, then

$$(13) \quad \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq e^{-b_C t} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $Y \in \xi^k(v^+)$ and non-zero $Z \in \xi^{d-k}(v^-)$. If this is not the case, then there exists

- a sequence $\{t_n\}$ in $[T, 2T]$,
- a nested sequence $\{H_n\}$ of horodisks centered at the fixed point of α whose intersection is empty,
- a sequence $\{v_n\}$ such that $\phi_s(v_n) \in \mathbf{U}(\Gamma)_{H_n}$ for all $s \in [0, t_n]$,
- a sequence vectors $\{Y_n\}$ such that $Y_n \in \xi^k(v_n^+)$,
- a sequence of non-zero vectors $\{Z_n\}$ such that $Z_n \in \xi^{d-k}(v_n^-)$,

such that

$$(14) \quad \frac{\|Y_n\|_{\phi_{t_n}(v_n)}}{\|Z_n\|_{\phi_{t_n}(v_n)}} > e^{-b_C t_n} \frac{\|Y_n\|_{v_n}}{\|Z_n\|_{v_n}}.$$

Let

$$u_t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad a_s = \begin{pmatrix} e^s & 0 \\ 0 & e^{-s} \end{pmatrix}.$$

As usual, we assume that $\Psi = \oplus_{i=1}^m \tau_{d_i}$, and $\alpha = [u_1]$. Choose a compact set $K \subset \mathbf{U}(\Gamma)_{\bar{H}}$ (where \bar{H} is the closure of H), so that

$$\bigcup_{s \in [0, \infty)} \bigcup_{m \in \mathbb{Z}} u_m a_s(K) = \mathbf{U}(\Gamma)_{\bar{H}}.$$

There is a sequence $\{s_n\} \rightarrow \infty$ of positive real numbers and a sequence $\{m_n\}$ of integers such that $w_n := a_{-s_n} u_{m_n}(v_n) \in K$ for all $n > 0$. By passing to a subsequence, we may assume that w_n converges to some $w_\infty \in \mathbf{U}(\Gamma)_{\bar{H}}$, and that

$$(V_n, W_n) := \left(\frac{\Psi(a_{-s_n} u_{m_n})(Y_n)}{\|\Psi(a_{-s_n} u_{m_n})(Y_n)\|_{w_n}}, \frac{\Psi(a_{-s_n} u_{m_n})(Z_n)}{\|\Psi(a_{-s_n} u_{m_n})(Z_n)\|_{w_n}} \right)$$

converges to some $(V_\infty, W_\infty) \in \mathbb{K}^d \times \mathbb{K}^d$. By definition,

$$(V_n, W_n) \in \Psi(a_{-s_n} u_{m_n})(\xi^k(v_n^+) \times \xi^{d-k}(v_n^-)) = \xi_n^k(w_n^+) \times \xi_n^{d-k}(w_n^-)$$

where $\xi_n = \Psi(a_{-s_n}) \circ \xi \circ a_{s_n}$. Proposition 5.3 (applied when U is a singleton) implies that $\lim_{n \rightarrow \infty} \xi_n = \eta$, so

$$(V_\infty, W_\infty) \in \eta^k(w_\infty^+) \times \eta^{d-k}(w_\infty^-).$$

Since $\|\cdot\|$ coincides with a Ψ -equivariant family of norms on $\mathbf{U}(\Gamma)_H$, and hence on $\mathbf{U}(\Gamma)_{\bar{H}}$, Equation (14) implies that

$$(15) \quad \frac{\|V_n\|_{\phi_{t_n}(w_n)}}{\|W_n\|_{\phi_{t_n}(w_n)}} = C_n \frac{\|Y_n\|_{\phi_{t_n}(v_n)}}{\|Z_n\|_{\phi_{t_n}(v_n)}} > C_n e^{-b_C t_n} \frac{\|Y_n\|_{v_n}}{\|Z_n\|_{v_n}} = e^{-b_C t_n} \frac{\|V_n\|_{w_n}}{\|W_n\|_{w_n}}.$$

where $C_n = \frac{\|\Psi(a_{-s_n} u_{m_n})(Z_n)\|_{w_n}}{\|\Psi(a_{-s_n} u_{m_n})(Y_n)\|_{w_n}}$. By passing to a final subsequence, we can suppose that $t_n \rightarrow t \in [T, 2T]$. Then taking $n \rightarrow \infty$ in Equation (15) we obtain

$$\frac{\|V_\infty\|_{\phi_t(w_\infty)}}{\|W_\infty\|_{\phi_t(w_\infty)}} \geq e^{-b_C t} \frac{\|V_\infty\|_{w_\infty}}{\|W_\infty\|_{w_\infty}} > B e^{-bt} \frac{\|V_\infty\|_{w_\infty}}{\|W_\infty\|_{w_\infty}}.$$

Since $V_\infty \in \eta^k(w_\infty^+)$ and $W_\infty \in \eta^{d-k}(w_\infty^-)$, this contradicts (11), thus proving the claim.

Repeated applications of the claim implies that if $v \in \mathbf{U}(\Gamma)_{H'}$ and $t \geq T$ is a number such that $\phi_s(v) \in \mathbf{U}(\Gamma)_{H'}$ for all $s \in [0, t]$, then

$$\frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} \leq e^{-b_C t} \frac{\|Y\|_v}{\|Z\|_v}$$

for all non-zero $Y \in \xi^k(v^+)$ and $Z \in \xi^{d-k}(v^-)$. Hence, if we define

$$B_C = \max \left\{ e^{-b_C t} \frac{\|Z\|_v \|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)} \|Y\|_v} : 0 \leq t \leq T, v \in K, Y, Z \in \mathbb{K}^d - \{0\} \right\},$$

(notice that B_C is finite by Lemma 3.8) then the proposition follows. \square

It remains to control the behavior of the flow on the thick part. The proof of the following proposition is inspired by arguments of Tsouvalas [33, Theorem 1.1].

Proposition 6.5. *Suppose that \widehat{K} is a compact subset of $\widehat{\mathbf{U}}(\Gamma)$, $\{u_n\}$ is a sequence in \widehat{K} , $Y_n \in \widehat{\Theta}^k|_{u_n}$, $Z_n \in \widehat{\Xi}^{d-k}|_{u_n}$ and $\|Y_n\|_{u_n} = \|Z_n\|_{u_n} = 1$ for all n . If $\{t_n\} \subset \mathbb{R}$, $\lim t_n = +\infty$ and $\phi_{t_n}(u_n) \in \widehat{K}$ for all n , then*

$$\lim_{n \rightarrow \infty} \frac{\|\phi_{t_n}(Y_n)\|_{\phi_{t_n}(u_n)}}{\|\phi_{t_n}(Z_n)\|_{\phi_{t_n}(u_n)}} = 0.$$

Proof. We first choose a compact set $K \subset \mathbf{U}(\Gamma)$ so that $\widehat{K} \subset \pi(K)$ (where $\pi : \mathbf{U}(\Gamma) \rightarrow \widehat{\mathbf{U}}(\Gamma)$ is the quotient map). For each n , choose $v_n \in K$ so that $\pi(v_n) = u_n$ and $\gamma_n \in \Gamma$ so that $w_n = \gamma_n^{-1}(\phi_{t_n}(v_n)) \in K$. We may assume that $v_n \rightarrow v_\infty$ and $w_n \rightarrow w_\infty$ for some $v_\infty, w_\infty \in K$. Notice that $\gamma_n \rightarrow v_\infty^+$ and $\gamma_n^{-1} \rightarrow w_\infty^-$.

Let $V_n \in \xi^k(v_n^+)$ and $W_n \in \xi^{d-k}(v_n^-)$ denote lifts of Y_n and Z_n respectively. Then

$$\frac{\|\phi_{t_n}(Y_n)\|_{\phi_{t_n}(u_n)}}{\|\phi_{t_n}(Z_n)\|_{\phi_{t_n}(u_n)}} = \frac{\|V_n\|_{\phi_{t_n}(v_n)}}{\|W_n\|_{\phi_{t_n}(v_n)}} = \frac{\|\rho(\gamma_n)^{-1}(V_n)\|_{w_n}}{\|\rho(\gamma_n)^{-1}(W_n)\|_{w_n}},$$

Also, since K is compact, there exists L so that if $v \in K$, then $\|\cdot\|_v$ is L -bilipschitz to the standard norm $\|\cdot\|_2$ on \mathbb{K}^d . So it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{\|\rho(\gamma_n)^{-1}(V_n)\|_2}{\|\rho(\gamma_n)^{-1}(W_n)\|_2} = 0.$$

Since ξ is strongly dynamics preserving, Lemma 2.2 implies that

$$(16) \quad \lim_{n \rightarrow \infty} \frac{\sigma_{d-k}(\rho(\gamma_n)^{-1})}{\sigma_{d-k+1}(\rho(\gamma_n)^{-1})} = \infty,$$

$U_k(\rho(\gamma_n)) \rightarrow \xi^k(v_\infty^+)$, and $U_{d-k}(\rho(\gamma_n)^{-1}) \rightarrow \xi^{d-k}(w_\infty^-)$. By the ρ -equivariance of ξ , $\rho(\gamma_n)^{-1}(W_n) \in \xi^{d-k}(w_n^-)$, which implies that

$$\lim_{n \rightarrow \infty} \angle(U_{d-k}(\rho(\gamma_n)^{-1}), \rho(\gamma_n)^{-1}(W_n)) = 0.$$

Therefore,

$$\liminf_{n \rightarrow \infty} \frac{\|\rho(\gamma_n)^{-1}(W_n)\|_2}{\sigma_{d-k}(\rho(\gamma_n)^{-1})} \geq \liminf_{n \rightarrow \infty} \|W_n\|_2 \geq \frac{1}{L}.$$

We now write $V_n = V'_n + V''_n$, where $V'_n \in U_k(\rho(\gamma_n))$ and $V''_n \in \rho(\gamma_n)(U_{d-k}(\rho(\gamma_n)^{-1}))$ are orthogonal. Notice that

$$\limsup_{n \rightarrow \infty} \frac{\|\rho(\gamma_n)^{-1}(V'_n)\|_2}{\sigma_{d-k+1}(\rho(\gamma_n)^{-1})} \leq L$$

so

$$\lim_{n \rightarrow \infty} \frac{\|\rho(\gamma_n)^{-1}(V'_n)\|_2}{\|\rho(\gamma_n)^{-1}(W_n)\|_2} = 0.$$

As such, if $\frac{\|\rho(\gamma_n)^{-1}(V_n)\|_2}{\|\rho(\gamma_n)^{-1}(W_n)\|_2}$ does not converge to 0 it must be the case that $\frac{\|\rho(\gamma_n)^{-1}(V''_n)\|_2}{\|\rho(\gamma_n)^{-1}(W_n)\|_2}$ does not converge to 0, and hence that

$$\limsup_{n \rightarrow \infty} \frac{\|\rho(\gamma_n)^{-1}(V''_n)\|_2}{\|\rho(\gamma_n)^{-1}(W_n)\|_2} = \infty.$$

We may then pass to a subsequence so that the limits

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)^{-1}(\text{Span}_{\mathbb{K}}(V''_n)) = \lim_{n \rightarrow \infty} \rho(\gamma_n)^{-1}(\text{Span}_{\mathbb{K}}(V_n))$$

are equal and exist. At the same time,

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)^{-1} \left(\text{Span}_{\mathbb{K}}(V_n'') \right) \subset \lim_{n \rightarrow \infty} U_{d-k}(\rho(\gamma_n)^{-1}) = \xi^{d-k}(w_\infty^-).$$

and

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)^{-1} \left(\text{Span}_{\mathbb{K}}(V_n) \right) \subset \lim_{n \rightarrow \infty} \rho(\gamma_n)^{-1} (\xi^k(v_n^+)) = \lim_{n \rightarrow \infty} \xi^k(w_n^+) = \xi^k(w_\infty^+).$$

This contradicts the transversality of ξ , and completes the proof. \square

We now combine Proposition 6.4 and Proposition 6.5 to finish the proof of Theorem 1.2.

For each $C \in \mathcal{C}$, let $C' \subset C$ be the embedded cusp subneighborhood given by Proposition 6.4. Let $\mathcal{C}' = \{C' : C \in \mathcal{C}\}$ and let $\widehat{K} = \widehat{U}(\Gamma) - \widehat{U}(\Gamma)_{\mathcal{C}'}$, which is compact. Let $b = \min\{b_C : C \in \mathcal{C}\}$ and $B = \max\{B_C : C \in \mathcal{C}\}$.

Proposition 6.5 implies that there exists $T_1 > 0$ so that: if $t \geq T_1$, $v \in \widehat{K}$ with $\phi_t(v) \in \widehat{K}$, $Y \in \widehat{\Theta}^k|_v$, and $Z \in \widehat{\Xi}^{d-k}|_v$ is non-zero, then

$$(17) \quad \frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq \frac{1}{2B^2} \frac{\|Y\|_v}{\|Z\|_v}.$$

Moreover, since \widehat{K} is compact, there exists $R > 0$ so that: if $v \in \widehat{K}$, $0 \leq t \leq T_1$, $Y \in \widehat{\Theta}^k|_v$ and $Z \in \widehat{\Xi}^{d-k}|_v$ is non-zero, then

$$(18) \quad \frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq R e^t \frac{\|Y\|_v}{\|Z\|_v}.$$

Now choose $T > 0$ so that

$$B^2 R e^{T_1(1+b)-bT} \leq \frac{1}{2} \quad \text{and} \quad B e^{-bT} \leq \frac{1}{2}.$$

We claim that if $t \geq T$, $v \in \widehat{U}(\Gamma)$, $Y \in \widehat{\Theta}^k|_v$ and $Z \in \widehat{\Xi}^{d-k}|_v$ is non-zero, then

$$(19) \quad \frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq \frac{1}{2} \frac{\|Y\|_v}{\|Z\|_v}.$$

Once we have proven our claim, we can choose

$$a := \frac{\log(2)}{T} \quad \text{and} \quad A := \max \left\{ \frac{\|\phi_t(Y)\|_{\phi_t(v)} \|Z\|_v}{\|\phi_t(Z)\|_{\phi_t(v)} \|Y\|_v} : 0 \leq t \leq T, Y \in \widehat{\Theta}^k|_v - 0, Z \in \widehat{\Xi}^{d-k}|_v - 0 \right\},$$

(notice that A is finite by Lemma 3.8) and observe that

$$\frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq A e^{-at} \frac{\|Y\|_v}{\|Z\|_v}$$

for all $t > 0$, $v \in \widehat{U}(\Gamma)$, $Y \in \widehat{\Theta}^k|_v$ and non-zero $Z \in \widehat{\Xi}^{d-k}|_v$. Hence, we will have shown that ρ is P_k -Anosov.

We now establish (19). Fix $v \in \widehat{U}(\Gamma)$, $t \geq T$, $Y \in \widehat{\Theta}^k|_v$ and non-zero $Z \in \widehat{\Xi}^{d-k}|_v$. If $\phi_s(v) \notin \widehat{K}$ for all $s \in [0, t]$, then Proposition 6.4 implies that

$$\frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} \leq B e^{-bt} \frac{\|Y\|_v}{\|Z\|_v} \leq \frac{1}{2} \frac{\|Y\|_v}{\|Z\|_v}.$$

Otherwise let

$$s_1 = s_1(t, v) := \min \left\{ s \in [0, t] : \phi_s(v) \in \widehat{K} \right\}$$

and

$$s_2 = s_2(t, v) := \max \left\{ s \in [0, t] : \phi_s(v) \in \widehat{K} \right\}.$$

If $s_2 - s_1 \geq T_1$, then Proposition 6.4 and (17) give

$$\begin{aligned} \frac{\|\phi_t(Y)\|_{\phi_t(v)}}{\|\phi_t(Z)\|_{\phi_t(v)}} &\leq (Be^{-bs_1}) \left(\frac{1}{2B^2}\right) (Be^{-b(t-s_2)}) \frac{\|Y\|_v}{\|Z\|_v} \\ &= \frac{1}{2} e^{b(s_2-s_1-t)} \frac{\|Y\|_v}{\|Z\|_v} \leq \frac{1}{2} \frac{\|Y\|_v}{\|Z\|_v}. \end{aligned}$$

On the other hand, if $s_2 - s_1 < T_1$, then by Proposition 6.4 and (18), we have

$$\begin{aligned} \frac{\|Y\|_{\phi_t(v)}}{\|Z\|_{\phi_t(v)}} &\leq (Be^{-bs_1}) (Re^{s_2-s_1}) (Be^{-b(t-s_2)}) \frac{\|Y\|_v}{\|Z\|_v} \\ &\leq (B^2 Re^{T_1(1+b)-bT}) \frac{\|Y\|_v}{\|Z\|_v} \leq \frac{1}{2} \frac{\|Y\|_v}{\|Z\|_v}. \end{aligned} \quad \square$$

7. HITCHIN REPRESENTATIONS ARE BOREL ANOSOV

In this section, we show that Hitchin representations are irreducible and Borel Anosov, i.e. P_k -Anosov for all k . Theorem 1.2 reduces the proof that Hitchin representations are Borel Anosov to the claim that their limit maps are strongly dynamics preserving.

Theorem 7.1. *Suppose $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho : \Gamma \rightarrow \mathrm{SL}(d, \mathbb{R})$ is Hitchin with continuous positive ρ -equivariant map $\xi : \Lambda(\Gamma) \rightarrow \mathcal{F}_d$. If $\{\gamma_n\}$ is a sequence in Γ with $\gamma_n \rightarrow x \in \Lambda(\Gamma)$ and $\gamma_n^{-1} \rightarrow y \in \Lambda(\Gamma)$, then*

$$\rho(\gamma_n)(V) \rightarrow \xi^k(x)$$

for all V transverse to $\xi^{d-k}(y)$.

Proof. Our proof relies on an observation about convergence of sequences of flags.

Definition 7.2. For any positive triple of flags (F_1, F_2, F_3) in $\mathcal{F}(\mathbb{R}^d)$, define the open set

$$(20) \quad \mathcal{O}(F_1, F_2, F_3) := \left\{ F \in \mathcal{F}(\mathbb{R}^d) : (F_1, F, F_2, F_3) \text{ is positive} \right\}.$$

The following result is a corrected version of Lemma 3.16 in [6] (whose statement omits an additional assumption given here).

Lemma 7.3. *Let (F_+, F_-) be a transverse pair of flags in $\mathcal{F}(\mathbb{R}^d)$. Suppose that $\{F_{1,n}\}$, $\{F_{2,n}\}$ and $\{F_{3,n}\}$ are sequences in $\mathcal{F}(\mathbb{R}^d)$ such that*

- (1) $F_{1,n} \rightarrow F^+$ and $F_{2,n} \rightarrow F^+$,
- (2) $F_{3,n} \rightarrow F^-$, and
- (3) $(F_{1,n}, F_{2,n}, F_{3,n})$ is positive for all n .

If $\{F_n\}$ is a sequence in $\mathcal{F}(\mathbb{R}^d)$ such that $F_n \in \mathcal{O}(F_{1,n}, F_{2,n}, F_{3,n})$ for all n , then $F_n \rightarrow F^+$.

We first suppose that $x \neq y$. Then by passing to the tail of the sequence $\{\gamma_n\}$, we may assume that each γ_n is hyperbolic with attractor and repeller γ_n^+ and γ_n^- in $\Lambda(\Gamma)$. Then $\gamma_n^+ \rightarrow x$, $\gamma_n^- \rightarrow y$, and $\gamma_n(z) \rightarrow x$ for all $z \in \Lambda(\Gamma) - \{y\}$.

Since $\Lambda(\Gamma)$ is infinite, there are points $a, b \in \Lambda(\Gamma) - \{x, y\}$ such that

- either $x < a < b < y$ or $y < b < a < x$, and
- up to taking subsequences, the sequences $\{\gamma_n(a)\}$ and $\{\gamma_n(b)\}$ both converge monotonically to x , and from the same direction.

Observe that

$$\lim_{n \rightarrow \infty} \xi(\gamma_n(a)) = \xi(x) = \lim_{n \rightarrow \infty} \xi(\gamma_n(b)).$$

Now, consider the open sets

$$\mathcal{O}_n := \mathcal{O}(\xi(a), \xi(b), \xi(\gamma_n^-))$$

for all n . Since $\gamma_n^- \rightarrow y$, and either $a < b < y$ or $b < a < y$, it follows that there is some $N > 0$ such that either $a < b < \gamma_n^-$ for all $n \geq N$, or $b < a < \gamma_n^-$ for all $n \geq N$. Lemma 3.15 in [6] then implies that $\mathcal{O}_n = \mathcal{O}_m$ for all $n, m \geq N$. Hence, if we set $\mathcal{O} := \mathcal{O}_N$, then for all $n \geq N$, we have

$$\rho(\gamma_n)(\mathcal{O}) = \rho(\gamma_n)(\mathcal{O}_n) = \mathcal{O}(\xi(\gamma_n(a)), \xi(\gamma_n(b)), \xi(\gamma_n^-)).$$

Since $\xi(\gamma_n(a)), \xi(\gamma_n(b)) \rightarrow \xi(x)$ and $\xi(\gamma_n^-) \rightarrow \xi(y)$, we may apply Lemma 7.3 to deduce that

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)(F) = \xi(x)$$

for all $F \in \mathcal{O}$. Repeating the same argument with γ_n^{-1} , we see that there exists an open set $\mathcal{O}' \subset \mathcal{F}(\mathbb{R}^d)$ where

$$\lim_{n \rightarrow \infty} \rho(\gamma_n^{-1})(F) = \xi(y)$$

for all $F \in \mathcal{O}'$. Hence, we may apply Lemma 2.2 to deduce the proposition when $x \neq y$.

Now suppose that $x = y$. Pick $\gamma \in \Gamma$ such that $z := \gamma^{-1}(x) \neq x$. Then $\gamma_n \gamma \rightarrow x$, $(\gamma_n \gamma)^{-1} \rightarrow z \neq x$. By the first part, $\rho(\gamma_n \gamma)(F) \rightarrow \xi(x)$ for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi(z)$. Equivalently, $\rho(\gamma_n)(F) \rightarrow \xi(x)$ for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi(x)$. \square

We recall that positive tuples of flags are in general position in the following sense.

Proposition 7.4. (Fock-Goncharov [14, Prop. 9.4], Sun-Wienhard-Zhang [32, Prop 2.21]) *If (F_1, \dots, F_k) is a positive tuple of flags, $(n_i)_{i=1}^k \in \mathbb{N}^k$ and $n = n_1 + \dots + n_k \leq d$, then $\oplus_{i=1}^k F_i^{n_i}$ has dimension n .*

We also use the following equivalent formulation of the positivity of a quadruple of flags.

Lemma 7.5. *A quadruple of flags (F_1, F_2, F_3, F_4) is positive if and only if there is a basis (b_1, \dots, b_d) of \mathbb{R}^d such that $b_i \in F_1^i \cap F_3^{d-i+1}$ for all $i \in \{1, \dots, d\}$, and some $u, v \in U_{>0}(b_1, \dots, b_d)$ such that $u(F_3) = F_2$ and $v^{-1}(F_3) = F_4$.*

Proof. Suppose first that $(F_1, F_2, F_3, F_4) = (F_1, u(F_3), F_3, v^{-1}(F_3))$ for some basis (b_1, \dots, b_d) of \mathbb{R}^d such that $b_i \in F_1^i \cap F_3^{d-i+1}$ for all $i \in \{1, \dots, d\}$, and some $u, v \in U_{>0}(b_1, \dots, b_d)$. Then $v(F_1, F_2, F_3, F_4) = (F_1, vu(F_3), v(F_3), F_3)$, which implies that $v(F_1, F_2, F_3, F_4)$ is positive. Thus, (F_1, F_2, F_3, F_4) is positive.

Conversely, suppose that (F_1, F_2, F_3, F_4) is positive. By Proposition 7.4, F_4 and F_3 are both transverse to F_1 , there is a unique unipotent $w \in \mathrm{SL}(d, \mathbb{R})$ that fixes F_1 and sends F_4 to F_3 . Then $w(F_1, F_2, F_3, F_4)$ is positive, which implies that there is a basis (b_1, \dots, b_d) of \mathbb{R}^d such that $b_i \in F_1^i \cap F_3^{d-i+1}$ for all $i \in \{1, \dots, d\}$, and some $u, v \in U_{>0}(b_1, \dots, b_d)$ such that

$$(F_1, w(F_2), w(F_3), F_3) = w(F_1, F_2, F_3, F_4) = (F_1, vu(F_3), v(F_3), F_3).$$

Since v is unipotent, fixes F_1 , and sends F_3 to $w(F_3)$, it follows that $v = w$. Therefore,

$$(F_1, F_2, F_3, F_4) = v^{-1}(F_1, vu(F_3), v(F_3), F_3) = (F_1, u(F_3), F_3, v^{-1}(F_4)). \quad \square$$

Proof of Theorem 1.4. The fact that ρ is P_k -Anosov, and that $x \mapsto \xi^k(x)$ is the Anosov limit map for $k = 1, \dots, d-1$, follows from Theorems 1.2 and 7.1. Further, Theorem 7.1 and Lemma 2.2 imply that:

- (1) If $\alpha \in \Gamma$ is parabolic, then $\rho(\alpha)$ is weakly unipotent.
- (2) If $\gamma \in \Gamma$ is hyperbolic, then $\rho(\gamma)$ is loxodromic.

If α is parabolic, let $x \in \Lambda(\Gamma) - \{\alpha^+\}$, and note that $(\alpha^+, \alpha^{-1}(x), x, \alpha(x))$ is a cyclically ordered set of distinct points in $\Lambda(\Gamma)$. Then we may apply the following lemma to $(\xi(\alpha^+), \xi(\alpha^{-1}(x)), \xi(x), \xi(\alpha(x)))$ to further conclude that $\rho(\alpha) = \pm u$ for some unipotent $u \in \mathrm{SL}(d, \mathbb{R})$ with a single Jordan block.

Lemma 7.6. *Let $g \in \mathrm{SL}(d, \mathbb{R})$ be weakly unipotent, and suppose that there are flags $F_1, F_2 \in \mathcal{F}(\mathbb{R}^d)$ such that F_1 is fixed by g and $(F_1, g^{-1}(F_2), F_2, g(F_2))$ is positive. Then F_1 is the unique fixed flag of g , or equivalently, $g = \pm u$ for some unipotent $u \in \mathrm{SL}(d, \mathbb{R})$ with a single Jordan block.*

Proof. By Lemma 7.5, there is a basis (b_1, \dots, b_d) of \mathbb{R}^d such that $b_i \in F_1^i \cap F_2^{d-i+1}$ for all $i \in \{1, \dots, d\}$, and some $u, v \in U_{>0}(b_1, \dots, b_d)$ such that $u(F_2) = g(F_2)$ and $v^{-1}(F_2) = g^{-1}(F_2)$. Then $a = u^{-1}g$ and $b = gv^{-1}$ both fix F_1 and F_2 , so they are diagonal in the basis (b_1, \dots, b_d) . Furthermore, since g is weakly unipotent, the diagonal entries of a and b are either 1 or -1 .

Assume for contradiction that there is some $i, j \in \{1, \dots, d\}$ such that the i -th diagonal entry of b is 1, while the j -th diagonal entry of b is -1 . Since all the upper triangular entries of v are positive, this implies that the upper triangular entries of bv along the i -th row are positive, while the upper triangular entries of bv along the j -th row are negative. But this is impossible since $ua = g = bv$, and for every column of ua , the upper triangular entries in that column must have the same sign. As such, $b = \pm \text{id}$. This implies that $g_{ss} = \pm \text{id}$ and $g_u = v = u$.

It now suffices to show that u has a unique fixed flag in $\mathcal{F}(\mathbb{R}^d)$. Observe the following linear algebra facts:

- (1) If w is a unipotent element that is represented in a basis (e_1, \dots, e_d) by upper triangular matrix where all the upper triangular entries are positive, then the line spanned by e_1 is the unique fixed line of w .
- (2) If $w \in U_{>0}(b_1, \dots, b_d)$, then for all $k \in \{1, \dots, d-1\}$, the linear action of w on $\bigwedge^k \mathbb{R}^d$ is represented in the basis $(b_{i_1} \wedge \dots \wedge b_{i_k})_{1 \leq i_1 < \dots < i_k \leq d}$ by an upper triangular matrix where all the upper triangular entries are positive.

These two observations imply that the unique fixed flag of u is the flag F given by $F^k = \text{Span}_{\mathbb{R}}(b_1, \dots, b_k)$ for all $k \in \{1, \dots, d-1\}$. \square

It only remains to show that ρ is irreducible. Suppose that ρ is not irreducible. Then there is a proper subspace $W \subset \mathbb{R}^d$ which is invariant under $\rho(\Gamma)$. By Theorem 7.1, $\rho(\gamma)$ is loxodromic for any hyperbolic $\gamma \in \Gamma$, so either W contains the attracting fixed point (in $\mathbb{P}(\mathbb{R}^d)$) of $\rho(\gamma)$, or W lies in the repelling hyperplane (in $\mathbb{P}(\mathbb{R}^d)$) of $\rho(\gamma)$. Since ξ is transverse, this implies that either W lies in the repelling hyperplanes of $\rho(\gamma)$ for all hyperbolic $\gamma \in \Gamma$, or W contains the attracting fixed point of $\rho(\gamma)$ for all hyperbolic $\gamma \in \Gamma$. However, this contradicts Proposition 7.4 in either case. Therefore, ρ is irreducible. \square

8. STABILITY OF ANOSOV REPRESENTATIONS

In this section, we prove Theorem 1.3, which we restate here.

Theorem 8.1. *If $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is a geometrically finite group and $\rho_0 : \Gamma \rightarrow \text{SL}(d, \mathbb{K})$ is P_k -Anosov, then there exists an open neighborhood \mathcal{O} of ρ_0 in $\text{Hom}_{\text{tp}}(\rho_0)$, so that*

- (1) *If $\rho \in \mathcal{O}$, then ρ is P_k -Anosov.*
- (2) *There exists $\alpha > 0$ so that if $\rho \in \mathcal{O}$, then ξ_ρ is α -Hölder (with respect to any distance on $\Lambda(\Gamma)$ induced by a Riemannian metric on $\partial \mathbb{H}^2$ and any distance on $\text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$ induced by a Riemannian metric).*
- (3) *If $\{\rho_u\}_{u \in M}$ is a \mathbb{K} -analytic family of representations in \mathcal{O} and $z \in \Lambda(\Gamma)$ then the map from M to $\text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$ given by $u \rightarrow \xi_{\rho_u}(z)$ is \mathbb{K} -analytic.*

The proof of (1) is based on the proof of stability for Anosov diffeomorphisms on compact manifolds given in Shub's book [31, Cor. 5.19]. The two key features which allow us overcome the non-compactness of the base space is the smooth conjugacy of the flows on the cusps, see Equation (21), and uniform estimates for families of canonical norms, see Lemma 3.8.

Proof of Theorem 8.1: We define

$$E(\mathcal{O}) = \mathcal{O} \times \mathbf{U}(\Gamma) \times \mathbb{K}^d \quad \text{and} \quad \widehat{E}(\mathcal{O}) = \Gamma \backslash (\mathcal{O} \times \mathbf{U}(\Gamma) \times \mathbb{K}^d).$$

The geodesic flow on $\mathbf{U}(\Gamma)$ extends to a flow on $E(\mathcal{O})$ whose action is trivial on the first and third factor. This in turn descends to a flow on $\widehat{E}(\mathcal{O})$. As usual, we use ϕ_t to denote these flows. Also, notice that $\widehat{E}(\mathcal{O})|_\rho$ naturally identifies with \widehat{E}_ρ .

Let $\|\cdot\|^0$ be a canonical family of norms for \widehat{E}_{ρ_0} and let \mathcal{C} be a full collection of embedded cusp neighborhoods for Γ so that $\|\cdot\|^0$ is canonical with respect to \mathcal{C} . Suppose that $\widehat{E}_{\rho_0} = \widehat{\Theta}_{\rho_0}^k \oplus \widehat{\Xi}_{\rho_0}^{d-k}$ is the P_k -Anosov splitting of \widehat{E}_{ρ_0} .

A ϕ_t -invariant splitting of $\widehat{E}(\mathcal{O})$. First, we prove that (after possibly shrinking \mathcal{O}) there exists a continuous ϕ_t -invariant splitting

$$\widehat{E}(\mathcal{O}) = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$$

that restricts to the splitting $\widehat{E}_{\rho_0} = \widehat{\Theta}_{\rho_0}^k \oplus \widehat{\Xi}_{\rho_0}^{d-k}$ over ρ_0 .

If $C \in \mathcal{C}$ and $C = \langle \alpha \rangle \backslash H$, then by shrinking \mathcal{O} if necessary, we may assume that there is a continuous map $g_C : \mathcal{O} \rightarrow \mathrm{SL}(d, \mathbb{K})$ such that

$$g_C(\rho)\rho_0(\alpha)g_C(\rho)^{-1} = \rho(\alpha)$$

for all $\rho \in \mathcal{O}$. Moreover, if $\rho \in \mathcal{O}$, the bundle isomorphism

$$\Phi_\rho^H : E_{\rho_0}|_{\mathrm{U}(\Gamma)_H} \rightarrow E_\rho|_{\mathrm{U}(\Gamma)_H} \quad \text{given by } (\rho_0, v, Z) \rightarrow (\rho, v, g_C(\rho)(Z))$$

descends to a bundle isomorphism

$$\widehat{\Phi}_\rho^C : \widehat{E}_{\rho_0}|_{\widehat{\mathrm{U}}(\Gamma)_C} \rightarrow \widehat{E}_\rho|_{\widehat{\mathrm{U}}(\Gamma)_C}$$

so that if $\phi_s(Z) \in \widehat{E}_{\rho_0}|_{\widehat{\mathrm{U}}(\Gamma)_C}$ for all $s \in [0, t]$, then

$$(21) \quad \widehat{\Phi}_\rho^C(\phi_t(Z)) = \phi_t(\widehat{\Phi}_\rho^C(Z)).$$

With this, we may extend the splitting $\widehat{E}_{\rho_0} = \widehat{\Theta}_{\rho_0}^k \oplus \widehat{\Xi}_{\rho_0}^{d-k}$ to a global splitting

$$\widehat{E}(\mathcal{O}) = \widehat{F}^k \oplus \widehat{G}^{d-k}$$

by first setting

$$\widehat{F}^k|_{(\rho, v)} = \widehat{\Phi}_\rho^C(\widehat{\Theta}_{\rho_0}^k|_v) \quad \text{and} \quad \widehat{G}^{d-k}|_{(\rho, v)} = \widehat{\Phi}_\rho^C(\widehat{\Xi}_{\rho_0}^{d-k}|_v)$$

for all $\rho \in \mathcal{O}$, $C \in \mathcal{C}$ and $v \in \mathrm{U}(\Gamma)_C$, and then extending this globally after perhaps shrinking \mathcal{O} and each C .

The flow ϕ_t does not necessarily preserve the splitting $\widehat{E}(\mathcal{O}) = \widehat{F}^k \oplus \widehat{G}^{d-k}$. To find the required ϕ_t -invariant splitting, we will use the contraction mapping theorem. For that purpose, we extend $\|\cdot\|^0$ to a canonical family of norms $\|\cdot\|$ on the fibers of $\widehat{E}(\mathcal{O})$ over $\mathcal{O} \times \widehat{\mathrm{U}}(\Gamma)$ as follows. If $C \in \mathcal{C}$, we define

$$\left\| \widehat{\Phi}_\rho^C(Z) \right\|_{(\rho, v)} = \|Z\|_{(\rho_0, v)}^0 \quad \text{for all } \rho \in \mathcal{O}, \quad v \in \widehat{\mathrm{U}}(\Gamma)_C \text{ and } Z \in \widehat{E}_{\rho_0}|_v.$$

This gives us canonical norms over all $C \in \mathcal{C}$. Then, perhaps after once more shrinking \mathcal{O} and each C , we may extend this to a continuous family of norms $\|\cdot\|$ for the fibers of $\widehat{E}(\mathcal{O})$ over $\mathcal{O} \times \widehat{\mathrm{U}}(\Gamma)$ such that the restriction to \widehat{E}_ρ is canonical for all $\rho \in \mathcal{O}$.

Suppose that V is a subspace of $\widehat{E}(\mathcal{O})|_{(\rho, v)}$, W is a subspace of $\widehat{E}(\mathcal{O})|_{(\rho, w)}$ and $T \in \mathrm{Hom}(V, W)$. We define the operator norm

$$\|T\|_{(\rho, v)} := \max \left\{ \|T(Z)\|_{(\rho, w)} : Z \in V, \|Z\|_{(\rho, v)} = 1 \right\}.$$

Then, if $V, W \subset \widehat{E}(\mathcal{O})$ are subbundles, $Q : \widehat{\mathrm{U}}(\Gamma) \rightarrow \widehat{\mathrm{U}}(\Gamma)$ is an isomorphism, and $T : V \rightarrow W$ is a map that restricts to a linear map $T|_{(\rho, v)} : V|_{(\rho, v)} \rightarrow W|_{(\rho, Q(v))}$ for all $\rho \in \mathcal{O}$ and $v \in \widehat{\mathrm{U}}(\Gamma)$, we define

$$\|T\|_X := \sup_{\rho \in \mathcal{O}, v \in X} \|T|_{(\rho, v)}\|_{(\rho, v)}$$

for any subset $X \subset \widehat{\mathrm{U}}(\Gamma)$. In the case when $Q = \mathrm{id}$, we may view T as an element of $S(\mathrm{Hom}(V, W))$, the vector space of sections of the bundle $\mathrm{Hom}(V, W)$. Note then that $\|\cdot\|_{\widehat{\mathrm{U}}(\Gamma)}$ defines a norm on $S(\mathrm{Hom}(V, W))$ whose corresponding distance is Cauchy complete.

We may decompose the flow ϕ_t as

$$\phi_t = \begin{pmatrix} A_t & B_t \\ C_t & D_t \end{pmatrix}$$

relative to the splitting $\widehat{E}(\mathcal{O}) = \widehat{F}^k \oplus \widehat{G}^{d-k}$. Here, $A_t : \widehat{F}^k \rightarrow \widehat{F}^k$ is a map such that for all $\rho \in \mathcal{O}$ and $v \in \widehat{U}(\Gamma)$, A_t restricts to a linear map $A_{(\rho,v,t)} : \widehat{F}^k|_{(\rho,v)} \rightarrow \widehat{F}^k|_{(\rho,\phi_t(v))}$, etc. In the case when this splitting happens to be ϕ_t -invariant, then $B_t \equiv 0$ and $C_t \equiv 0$.

Lemma 8.2. *Up to taking a subneighborhood of \mathcal{O} , there exists $T > 0$ so that if $t \in [T, 2T]$, then A_t and D_t are invertible and*

$$\max \{ \|A_t^{-1}B_t\|, \|D_t^{-1}C_t\|, L_t \} < \frac{1}{3},$$

where $L_t := \sup_{\rho \in \mathcal{O}, v \in \widehat{U}(\Gamma)} \left(\|A_{(\rho,v,t)}\|_{(\rho,v)} \|D_{(\rho,v,t)}^{-1}\|_{(\rho,\phi_t(v))} \right)$.

Proof. For a subset $X \subset \widehat{U}(\Gamma)$, define

$$L_t(X) := \sup_{\rho \in \mathcal{O}, v \in X} \left(\|A_{(\rho,v,t)}\|_{(\rho,v)} \|D_{(\rho,v,t)}^{-1}\|_{(\rho,\phi_t(v))} \right).$$

First notice that for all $v \in \widehat{U}(\Gamma)$, $B_{(\rho_0,v,t)} = 0$, $C_{(\rho_0,v,t)} = 0$ and both $A_{(\rho_0,v,t)}$, $D_{(\rho_0,v,t)}$ are invertible since $F^k \oplus G^{d-k}|_{\rho_0} = \widehat{\Theta}_{\rho_0}^k \oplus \widehat{\Xi}_{\rho_0}^{d-k}$ is a flow invariant splitting. Since ρ_0 is P_k -Anosov, there exist $c, C > 0$ such that

$$\|A_{(\rho_0,v,t)}\|_{(\rho_0,v)} \|D_{(\rho_0,v,t)}^{-1}\|_{(\rho_0,\phi_t(v))} \leq Ce^{-ct}.$$

for all $v \in \widehat{U}(\Gamma)$ and $t > 0$. Choose T so that $Ce^{-cT} < \frac{1}{3}$.

Consider the compact set

$$X = \left\{ v \in \widehat{U}(\Gamma) : \phi_t(v) \notin \widehat{U}(\Gamma)_C \text{ for some } t \in [0, 2T] \right\}.$$

By shrinking \mathcal{O} if necessary, we can ensure that on $\widehat{E}(\mathcal{O})|_{\mathcal{O} \times X}$, if $t \in [T, 2T]$, then A_t and D_t are invertible, and

$$\max \{ \|A_t^{-1}B_t\|_X, \|D_t^{-1}C_t\|_X, L_t(X) \} < \frac{1}{3}.$$

On the other hand, if $v \in \widehat{U}(\Gamma) - X$, then there is some $C \in \mathcal{C}$ such that $\phi_t(v) \in \widehat{U}(\Gamma)_C$ for all $t \in [0, 2T]$. Then by construction,

$$\phi_{(\rho,v,t)} = \widehat{\Phi}_{\rho}^C \circ \phi_{(\rho_0,v,t)} \circ (\widehat{\Phi}_{\rho}^C)^{-1}$$

for all $\rho \in \mathcal{O}$ and $t \in [0, 2T]$. Since $\widehat{\Phi}_{\rho}^C$ is an isometry that preserves the splitting, it follows that on $\widehat{E}(\mathcal{O})|_{\mathcal{O} \times (\widehat{U}(\Gamma) - X)}$, if $t \in [T, 2T]$, then A_t and D_t are invertible, $B_t = 0$, $C_t = 0$, and

$$L_t(\widehat{U}(\Gamma) - X) = \sup_{v \in \widehat{U}(\Gamma) - X} \|A_{(\rho_0,v,t)}\|_{(\rho_0,v)} \|D_{(\rho_0,v,t)}^{-1}\|_{(\rho_0,\phi_t(v))} < \frac{1}{3}.$$

□

Consider the bundle $\text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k) \rightarrow \mathcal{O} \times \widehat{U}(\Gamma)$ with its induced operator norm $\|\cdot\|$. Let $\mathcal{R}_r \subset \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)$ denote the ball bundle of radius r about the zero section.

Proposition 8.3. *If $t \in [T, 2T]$, then there is a well-defined map $\psi_t : \mathcal{R}_1 \rightarrow \mathcal{R}_{\frac{2}{3}}$ given by*

$$\psi_t(f) = (B_t + A_t f)(D_t + C_t f)^{-1}$$

for all $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f \in \mathcal{R}_1|_{(\rho,v)}$. Furthermore:

- (1) $\|\psi_t(f_1) - \psi_t(f_2)\| \leq \frac{5}{6} \|f_1 - f_2\|$ for all $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f_1, f_2 \in \mathcal{R}_1|_{(\rho,v)}$.
- (2) If $B_t \equiv 0$ and $C_t \equiv 0$, then ψ_t extends to a flow on $\text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)$,

$$\psi_t(f) = \phi_t \circ f \circ \phi_{-t}$$

and ψ_t is uniformly contracting.

Remark 8.4. One can verify that the map ψ_t has the defining property

$$\text{Graph}(\psi_t(f)) = \phi_t(\text{Graph}(f))$$

for all $t \in [T, 2T]$, $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f \in \mathcal{R}_1|_{(\rho, v)}$. Similarly, if $B_t \equiv 0$ and $C_t \equiv 0$, then

$$\psi_t(f) = A_t f D_t^{-1} = \phi_t \circ f \circ \phi_{-t}$$

is a well defined flow on $\text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)$.

Proof of Proposition 8.3. If $t \in [T, 2T]$, then

$$D_t + C_t f = D_t (\text{id} + D_t^{-1} C_t f).$$

for all $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f \in \mathcal{R}_1|_{(\rho, v)}$. By Lemma 8.2, if $\|f\|_{(\rho, v)} < 1$, then

$$\|D_t^{-1} C_t f\|_{(\rho, v)} \leq \|D_t^{-1} C_t\|_{\widehat{U}(\Gamma)} \|f\|_{(\rho, v)} < \frac{1}{3}$$

which implies that $\text{id} + D_t^{-1} C_t f$ has trivial kernel for all $\rho \in \mathcal{O}$ and $v \in \widehat{U}(\Gamma)$. Hence, $\text{id} + D_t^{-1} C_t f$ is invertible. Since Lemma 8.2 also gives that D_t is invertible, it follows that

$$(B_t + A_t f) (D_t + C_t f)^{-1} \in \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)|_{(\rho, \phi_t(v))}$$

is a well-defined for all $f \in \mathcal{R}_1|_{(\rho, v)}$.

It remains to show that $(B_t + A_t f) (D_t + C_t f)^{-1} \in \mathcal{R}_{\frac{2}{3}}|_{(\rho, \phi_t(v))}$. By Lemma 8.2,

$$\|A_t^{-1} B_t + f\|_{(\rho, v)} \leq \frac{1}{3} + 1 = \frac{4}{3}$$

and

$$\|(\text{id} + D_t^{-1} C_t f)^{-1}\|_{(\rho, v)} \leq \frac{1}{1 - \frac{1}{3}} = \frac{3}{2}.$$

Thus,

$$\left\| (B_t + A_t f) (D_t + C_t f)^{-1} \right\|_{(\rho, \phi_t(v))} \leq L_t \|A_t^{-1} B_t + f\|_{(\rho, v)} \|(\text{id} + D_t^{-1} C_t f)^{-1}\|_{(\rho, v)} \leq \frac{1}{3} \cdot \frac{4}{3} \cdot \frac{3}{2} = \frac{2}{3}.$$

Next, we prove (1). For any $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$, $f \in \mathcal{R}_1|_{(\rho, v)}$ and any $\eta \in \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)|_{(\rho, v)}$,

$$\begin{aligned} \left. \frac{d}{ds} \right|_{s=0} (\psi_t)(f + s\eta) &= A_t \eta (D_t + C_t f)^{-1} - (B_t + A_t f) (D_t + C_t f)^{-1} C_t \eta (D_t + C_t f)^{-1} \\ &= A_t \left(\eta (\text{id} + D_t^{-1} C_t f)^{-1} - (A_t^{-1} B_t + f) (\text{id} + D_t^{-1} C_t f)^{-1} D_t^{-1} C_t \eta (\text{id} + D_t^{-1} C_t f)^{-1} \right) D_t^{-1} \\ &\in \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)|_{(\rho, \phi_t(v))}. \end{aligned}$$

Thus,

$$\left\| \left. \frac{d}{ds} \right|_{s=0} (\psi_t)(f + s\eta) \right\|_{(\rho, \phi_t(v))} \leq \frac{1}{3} \left(\frac{3}{2} + \frac{4}{3} \cdot \frac{3}{2} \cdot \frac{1}{3} \cdot \frac{3}{2} \right) \|\eta\|_{(\rho, v)} = \frac{5}{6} \|\eta\|_{(\rho, v)},$$

which implies (1).

Finally, we prove (2). Note that if $B_t \equiv 0$ and $C_t \equiv 0$, then A_t and D_t are invertible for all $t \in \mathbb{R}$. Thus, ψ_t extends to a well-defined flow given by

$$\psi_t(f) = A_t f D_t^{-1} = \phi_t \circ f \circ \phi_{-t}$$

for all $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f \in \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)|_{(\rho, v)}$. By Lemma 8.2, if $t \in [T, 2T]$, then

$$\|\psi_t(f)\|_{(\rho, \phi_t(v))} \leq L_t \|f\|_{(\rho, v)} \leq \frac{1}{3} \|f\|_{(\rho, v)}$$

for all $\rho \in \mathcal{O}$, $v \in \widehat{U}(\Gamma)$ and $f \in \text{Hom}(\widehat{G}^{d-k}, \widehat{F}^k)|_{(\rho, v)}$. Hence, for all $t \geq 0$,

$$\|f\|_{(\rho, \phi_t(v))} \leq C_0 e^{-c_0 t} \|f\|_{(\rho, v)},$$

where $c_0 := \frac{\log 3}{T}$ and

$$C_0 := \sup \left\{ e^{c_0 t} \frac{\|f\|_{(\rho, \phi_t(v))}}{\|f\|_{(\rho, v)}} : \rho \in \mathcal{O}, v \in \mathbf{U}(\Gamma), f \in \text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)|_{(\rho, v)} - 0, t \in [0, T] \right\}$$

(notice that C_0 is finite by Lemma 3.8). \square

Let $S(\mathcal{R}_r)$ be the space of continuous sections of $\mathcal{R}_r \rightarrow \mathcal{O} \times \mathbf{U}(\Gamma)$. Notice that ψ_t induces a map $\psi_t^S : S(\mathcal{R}_1) \rightarrow S(\mathcal{R}_{2/3})$ given by

$$\psi_t^S(\sigma)(\rho, v) = \psi_t(\sigma(\rho, \phi_{-t}(v))).$$

By part (1) of Proposition 8.3, the map ψ_t^S is a contraction mapping on $S(\mathcal{R}_1)$ for each $t \in [T, 2T]$. We may now apply the contraction mapping theorem to conclude that for each $t \in [T, 2T]$ there exists a unique ψ_t^S -invariant section $\sigma^{(t)}$ of the bundle $\mathcal{R}_{2/3}$.

We claim that $\sigma^{(t)}$ does not depend on t . If $t_1, t_2 \in \mathbb{Q} \cap [T, 2T]$, then there exist sequences $\{n_j\}, \{m_j\}$ with $n_j, m_j \rightarrow \infty$ and $n_j t_1 = m_j t_2$ for all $j \geq 1$. Then by the proof of the contraction mapping theorem

$$\sigma^{(t_1)} = \lim_{j \rightarrow \infty} \psi_{n_j t_1}^S(\sigma^{(t_2)}) = \lim_{j \rightarrow \infty} \psi_{m_j t_2}^S(\sigma^{(t_2)}) = \sigma^{(t_2)}.$$

So $\sigma^{(t)}$ does not depend on t when $t \in \mathbb{Q} \cap [T, 2T]$. Then by uniqueness of invariant sections and the continuity of ψ_t^S , we see that $\sigma^{d-k} := \sigma^{(t)}$ does not depend on t . Then σ^{d-k} determines a ϕ_t -invariant $(d-k)$ -dimensional subbundle $\widehat{\Xi}^{d-k}$ of $\widehat{E}(\mathcal{O})$ defined by

$$\widehat{\Xi}^{d-k}|_{(\rho, v)} = \text{Graph } \sigma^{d-k}(\rho, v),$$

see Remark 8.4.

Applying a similar argument to the bundle $\text{Hom}(\widehat{F}^k, \widehat{G}^{d-k})$ we obtain, by further shrinking \mathcal{O} if necessary, a ϕ_t -invariant k -dimensional subbundle $\widehat{\Theta}^k$ of $\widehat{E}(\mathcal{O})$.

To show that $\widehat{E}(\mathcal{O}) = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$, it now suffices to show that the fibers $\widehat{\Xi}^{d-k}|_{(\rho, v)}$ and $\widehat{\Theta}^k|_{(\rho, v)}$ are transverse for every $(\rho, v) \in \mathcal{O} \times \widehat{\mathbf{U}}(\Gamma)$. Suppose for contradiction that there is some non-zero $Z \in \widehat{\Xi}^{d-k}|_{(\rho, v)} \cap \widehat{\Theta}^k|_{(\rho, v)}$ for some $\rho \in \mathcal{O}$ and $v \in \widehat{\mathbf{U}}(\Gamma)$. We may write Z uniquely as $Z_k + Z_{d-k}$ where $Z_k \in \widehat{F}^k|_{(\rho, v)}$ and $Z_{d-k} \in \widehat{G}^{d-k}|_{(\rho, v)}$. Since $\widehat{\Xi}^{d-k}$ corresponds to the section $\sigma^{d-k} \in S(\mathcal{D}_{\frac{2}{3}})$, the fact that $Z \in \widehat{\Xi}^{d-k}|_{(\rho, v)}$ is non-zero implies that $Z_{d-k} \neq 0$ and

$$\frac{\|Z_k\|_{(\rho, v)}}{\|Z_{d-k}\|_{(\rho, v)}} = \frac{\|\sigma^{d-k}(\rho, v)(Z_{d-k})\|_{(\rho, v)}}{\|Z_{d-k}\|_{(\rho, v)}} \leq \|\sigma^{d-k}(\rho, v)\|_{(\rho, v)} < \frac{2}{3}.$$

For the same reasons, $Z_k \neq 0$ and

$$\frac{\|Z_{d-k}\|_{(\rho, v)}}{\|Z_k\|_{(\rho, v)}} < \frac{2}{3},$$

which is a contradiction. Thus, $\widehat{\Theta}^k$ and $\widehat{\Xi}^{d-k}$ indeed give a ϕ_t -invariant splitting of $\widehat{E}(\mathcal{O})$.

Relative to the splitting $\widehat{E}(\mathcal{O}) = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$ the decomposition of the flow ϕ_t on $\widehat{E}(\mathcal{O})$ simplifies to

$$\phi_t = \begin{pmatrix} A_t & 0 \\ 0 & D_t \end{pmatrix}.$$

Then by part (2) of Proposition 8.3 and Remark 8.4, we have the following corollary.

Corollary 8.5. *The flow $f \mapsto \phi_t \circ f \circ \phi_{-t}$ on $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ is uniformly contracting.*

Existence of limit maps. Next, we use the ϕ_t -invariant splitting $\widehat{E}(\mathcal{O}) = \widehat{\Theta}^k \oplus \widehat{\Xi}^{d-k}$ to define limit maps

$$\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d).$$

Lift this splitting of $\widehat{E}(\mathcal{O})$ to a splitting

$$E(\mathcal{O}) = \Theta^k \oplus \Xi^{d-k}.$$

The flow ϕ_t on $\widehat{E}(\mathcal{O})$ lifts to a flow, also denoted ϕ_t , on $E(\mathcal{O})$, under which this splitting is invariant. Then the bundle $\text{Hom}(\widehat{\Theta}^k, \widehat{\Xi}^{d-k})$ lifts to the bundle $\text{Hom}(\Theta^k, \Xi^{d-k})$. Finally, we use $\|\cdot\|$ to denote the lifted norms on $E(\mathcal{O})$ and $\text{Hom}(\Theta^k, \Xi^{d-k})$.

By Corollary 8.5, there exists $C_0, c_0 > 0$ such that

$$(22) \quad \|f\|_{(\rho, \phi_t(v))} \leq C_0 e^{-c_0 t} \|f\|_{(\rho, v)}$$

for all $f \in \text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v)} = \text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, \phi_t(v))}$ and $t \geq 0$.

Let

$$\sigma = (\sigma^k, \sigma^{d-k}) : \mathcal{O} \times \text{U}(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}) \times \text{Gr}_{d-k}(\mathbb{K}^d)$$

be the map so that $\sigma^k(\rho, v) = \Theta^k|_{(\rho, v)}$ and $\sigma^{d-k}(\rho, v) = \Xi^{d-k}|_{(\rho, v)}$. Since σ is ϕ_t -invariant, $\sigma^k(\rho, v)$ and $\sigma^{d-k}(\rho, v)$ depend only on ρ , v^+ and v^- . We now check that σ^{d-k} depends only on ρ and v^- . Let $\gamma \in \Gamma$ be a hyperbolic element, let $v_\gamma \in \text{U}(\Gamma)$ be a vector so that $v_\gamma^+ = \gamma^+$ and $v_\gamma^- = \gamma^-$. Let $\ell(\gamma)$ be the translation distance of γ on \mathbb{H}^2 . Then $\phi_{n\ell(\gamma)}(v_\gamma) = \gamma^n(v_\gamma)$ for all n . Since σ^{d-k} is equivariant and ϕ_t -invariant, it follows that

$$\sigma^{d-k}(\rho, v_\gamma) = \sigma^{d-k}(\rho, \phi_{-\ell(\gamma)}(\gamma(v_\gamma))) = \rho(\gamma)(\sigma^{d-k}(\rho, v_\gamma))$$

for all ρ . Furthermore, if $W \subset \mathbb{K}^d$ is a $(d-k)$ -dimensional subspace that is transverse to $\sigma^k(\rho, v_\gamma)$, we may view W as the graph of an element $f_W \in \text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v_\gamma)}$.

Equation (22) implies that

$$\|\rho(\gamma)^{-n}(f_W)\|_{(\rho, v_\gamma)} = \|f_W\|_{(\rho, \gamma^n(v_\gamma))} = \|f_W\|_{(\rho, \phi_{n\ell(\gamma)}(v_\gamma))} \rightarrow 0,$$

which implies that $\rho(\gamma)^{-n}(W) \rightarrow \sigma^{d-k}(\rho, v_\gamma)$. Thus, $\sigma^{d-k}(\rho, v_\gamma)$ is the repelling fixed point of $\rho(\gamma)$ in $\text{Gr}_{d-k}(\mathbb{K}^d)$. Now if $x \in \Lambda(\Gamma) \setminus \{\gamma^-\}$, then there exists $v \in \text{U}(\Gamma)$ so that $v^+ = x$, $v^- = \gamma^-$ and

$$\lim_{t \rightarrow \infty} d_{\text{U}(\Gamma)}(\phi_{-t}(v_\gamma), \phi_{-t}(v)) = 0,$$

so $\gamma^n(\phi_{-n\ell(\gamma)}(v)) \rightarrow v_\gamma$. Thus,

$$\sigma^{d-k}(\rho, v_\gamma) = \lim_{n \rightarrow \infty} \sigma^{d-k}(\rho, \gamma^n(\phi_{-n\ell(\gamma)}(v))) = \lim_{n \rightarrow \infty} \rho(\gamma)^n(\sigma^{d-k}(\rho, v)).$$

Since $\sigma^{d-k}(\rho, v_\gamma)$ is the repelling fixed point of $\rho(\gamma)$ in $\text{Gr}_{d-k}(\mathbb{K}^d)$, this implies that $\sigma^{d-k}(\rho, v) = \sigma^{d-k}(\rho, v_\gamma)$. Therefore, since σ^{d-k} is ϕ_t -invariant, if $v^- = \gamma^-$, then $\sigma^{d-k}(\rho, v)$ is the repelling fixed point of $\rho(\gamma)$. Since every point in $\Lambda(\Gamma)$ is a limit of repelling fixed points of hyperbolic elements of Γ , this implies that σ^{d-k} depends only on ρ and v^- .

One may similarly show that σ^k depends only on ρ and v^+ , so there exists

$$\xi = (\xi^k, \xi^{d-k}) : \mathcal{O} \times \Lambda(\Gamma) \rightarrow \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$$

so that $\sigma(\rho, v) = (\xi^k(\rho, v^+), \xi^{d-k}(\rho, v^-))$. As such, if $\rho \in \mathcal{O}$, then ρ is P_k -Anosov. This proves (1).

The limit maps are Hölder. We now prove that, perhaps after shrinking our neighborhood \mathcal{O} again, that the limit maps are uniformly Hölder. It is possible to establish this using Shub's C^r -Section theorem [31, Thm. 5.18], however setting up bundles with the correct regularity (see [31, Cor. 5.19]) and verifying the admissibility condition is somewhat involved when $\widehat{\text{U}}(\Gamma)$ is non-compact. Instead we provide a direct argument based on the proof of Lemma 4.4 in [36].

We will continue to work with $E(\mathcal{O})$ and $\text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v)}$. For $\rho \in \mathcal{O}$, $v \in \text{U}(\Gamma)$ and $x \in \Lambda(\Gamma) \setminus \{v^+\}$, let $f_{\rho, v, x} \in \text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v)}$ denote the unique element with

$$\text{Graph}(f_{\rho, v, x}) = \xi_\rho^{d-k}(x).$$

Notice that $f_{\rho, \phi_t(v), x} = f_{\rho, v, x}$ for all $t \in \mathbb{R}$. If $v \in \mathbf{U}(\Gamma)$, we let $v^\perp \subset \partial\mathbb{H}^2$ denote the endpoints of the geodesic through the basepoint of v which is orthogonal to v . (One can use the orientation to canonically identify them as $v^{\perp,+}$ and $v^{\perp,-}$ but this will not be needed for our purposes.)

Lemma 8.6. *Up to taking a subneighborhood of \mathcal{O} , there exists $C_1 > 1$ so that if $\rho \in \mathcal{O}$, $v \in \mathbf{U}(\Gamma)$ and $x \in v^\perp \cap \Lambda(\Gamma)$, then*

$$\frac{1}{C_1} \leq \|f_{\rho, v, x}\|_{(\rho, v)} \leq C_1.$$

In the case when Γ is convex co-compact, the lemma is a simple consequence of equivariance and compactness, but in the general geometrically finite case the proof is somewhat involved. Delaying the proof of the lemma, we first complete the proof of part (2) of Theorem 8.1.

Let $\mathcal{O}' \subset \mathcal{O}$ be a subneighborhood of ρ_0 such that the closure of $\overline{\mathcal{O}'}$ of \mathcal{O}' is a compact subset of \mathcal{O} . Fix a compact set $K \subset \mathbf{U}(\Gamma)$ such that

$$\Lambda(\Gamma) = \{v^+ : v \in K\}.$$

Let d_∞ denote the distance induced by a Riemannian metric on $\partial\mathbb{H}$, and let d_G denote the distance induced by a Riemannian metric on $\text{Gr}_{d-k}(\mathbb{K}^d)$. Fix $\delta > 0$ such that: if $v \in K$ and $x \in \Lambda(\Gamma) - \{v^-\}$ satisfies $d_\infty(x, v^-) \leq \delta$, then there exists $t \geq 0$ such that $x \in \phi_{-t}(v)^\perp$ (in particular $x \neq v^+$).

Fix $\rho \in \mathcal{O}'$ and $v \in K$. Then d_G is bilipschitz to the norm $\|\cdot\|_{(\rho, v)}$ on any compact subset of the affine chart

$$\text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v)} \simeq \{V \in \text{Gr}_{d-k}(\mathbb{K}^d) : V \text{ is transverse to } \xi_\rho^k(v^+)\},$$

where the isomorphism identifies each $f \in \text{Hom}(\Xi^{d-k}, \Theta^k)|_{(\rho, v)}$ with its graph in $\text{Gr}_{d-k}(\mathbb{K}^d)$. By the compactness of $\overline{\mathcal{O}'} \times K$, there exists $C_2 > 0$ such that: if $\rho \in \mathcal{O}'$, $v \in K$ and $x \in \Lambda(\Gamma)$ with $d_\infty(x, v^-) \leq \delta$, then

$$d_G(\xi_\rho^{d-k}(x), \xi_\rho^{d-k}(v^-)) \leq C_2 \|f_{\rho, v, x}\|_{(\rho, v)}$$

There also exists $C_3 > 0$ such that: if $v \in K$, $x \in \Lambda(\Gamma)$, $d_\infty(x, v^-) \leq \delta$ and $x \in \phi_{-t}(v)^\perp$, then

$$\frac{1}{C_3} e^{-t} \leq d_\infty(x, v^-) \leq C_3 e^{-t}.$$

Finally, let

$$C_4 = \text{diam}\left(\text{Gr}_{d-k}(\mathbb{K}^d), d_G\right)$$

and $C = \max\{\delta^{-c_0} C_4, C_0 C_1 C_2 C_3^{c_0}\}$.

Now suppose $x, y \in \Lambda(\Gamma)$ and $\rho \in \mathcal{O}'$. If $d_\infty(x, y) > \delta$, then

$$d_G(\xi_\rho^{d-k}(x), \xi_\rho^{d-k}(y)) \leq C_4 \leq C_4 \frac{d_\infty(x, y)^{c_0}}{\delta^{c_0}} \leq C d_\infty(x, y)^{c_0}.$$

If $d_\infty(x, y) \leq \delta$, then there exists $t \geq 0$ and $v \in K$ such that $y = v^-$ and $x \in \phi_{-t}(v)^\perp$. Then

$$\begin{aligned} d_G(\xi_\rho^{d-k}(x), \xi_\rho^{d-k}(v^-)) &\leq C_2 \|f_{\rho, v, x}\|_{(\rho, v)} \leq C_0 C_2 e^{-c_0 t} \|f_{\rho, \phi_{-t}(v), x}\|_{(\rho, \phi_{-t}(v))} \\ &\leq C_0 C_1 C_2 C_3^{c_0} d_\infty(x, v^-)^{c_0} \leq C d_\infty(x, y)^{c_0}. \end{aligned}$$

Therefore, ξ_ρ^{d-k} is c_0 -Hölder.

One may similarly prove that, perhaps after passing to another sub-neighborhood, that if $\rho \in \mathcal{O}'$, then ξ_ρ^k is c_1 -Hölder for some $c_1 > 0$. So, (2) holds with $\alpha = \min\{c_0, c_1\}$.

Proof of Lemma 8.6. Let $\mathcal{O}' \subset \mathcal{O}$ be a subneighborhood of ρ_0 such that the closure of $\overline{\mathcal{O}'}$ of \mathcal{O}' is a compact subset of \mathcal{O} . If the lemma fails for \mathcal{O}' , then there exist sequences $\{\rho_m\}$ in \mathcal{O}' , $\{v_m\}$ in $\mathbf{U}(\Gamma)$ and $\{x_m\}$ in $\Lambda(\Gamma)$ so that $x_m \in v_m^\perp$ for all m and

$$(23) \quad \lim_{m \rightarrow \infty} \left| \log \|f_{\rho_m, v_m, x_m}\|_{(\rho_m, v_m)} \right| = +\infty.$$

By passing to a subsequence we can suppose that $\rho_m \rightarrow \rho \in \mathcal{O}$. After passing to a further subsequence and translating by elements in Γ , either

- (1) $v_m \rightarrow v \in \mathbf{U}(\Gamma)$ and $x_m \rightarrow x \in \Lambda(\Gamma)$, or
- (2) there exists an embedded cusp neighborhood $C = \langle \alpha \rangle \backslash H$ such that $\{v_m\} \subset \mathbf{U}(\Gamma)_H$ and $\{v_m\}$ projects to an escaping sequence in $\widehat{\mathbf{U}}(\Gamma)$.

In the first case, $x \in v^\perp$ and $\lim_{m \rightarrow \infty} \|f_{\rho_m, v_m, x_m}\|_{(\rho_m, v_m)} = \|f_{\rho, v, x}\|_{(\rho, v)} \neq 0$. Thus we must be in the second case.

By conjugating, we may assume that $\rho_m(\alpha) = \rho_0(\alpha)$ for all $m \in \mathbb{Z}^+ \cup \{\infty\}$, and that the restriction of the canonical norm on $E(\mathcal{O}')$ to each E_{ρ_m} is with respect to the same cusp representation Ψ for α and $\rho_0(\alpha)$. We may also assume, by translating by a power of α , that each v_m lies in a standard fundamental domain F for the action of α on $\mathbf{U}(\Gamma)_H$. Let $y \in \Lambda(\Gamma)$ be the center of H and fix a hyperbolic element $\gamma \in \mathbf{SL}(2, \mathbb{R})$ with attracting fixed point y , and whose axis intersects the interior of F . After passing to a subsequence, we can find $n_m \rightarrow \infty$ such that $\gamma^{-n_m}(v_m) \rightarrow v \in F$ and $\gamma^{-n_m}(x_m) \rightarrow x \in \Lambda(\Gamma)$.

Let $\eta^j : \partial \mathbb{H}^2 \rightarrow \text{Gr}_j(\mathbb{K}^d)$ denote the boundary maps associated to Ψ for $j = k, d-k$, see Proposition 5.2. Let $\hat{f} \in \text{Hom}(\eta^{d-k}(v^-), \eta^k(v^+))$ denote the unique element with

$$\text{Graph}(\hat{f}) = \eta^{d-k}(x).$$

Notice that $\hat{f} \neq 0$, since η^k and η^{d-k} are transverse and $x \in v^\perp$. By Proposition 5.3,

$$\Psi(\gamma^{-n_m})\left(\xi_{\rho_m}^j(v_m^\pm)\right) = \left(\Psi(\gamma^{-n_m}) \circ \xi_{\rho_m}^j \circ \gamma^{n_m}\right)(\gamma^{-n_m}(v_m^\pm)) \rightarrow \eta^j(v^\pm).$$

for $j = k, d-k$. Similarly, $\Psi(\gamma^{-n_m})(\xi_{\rho_m}^j(x)) \rightarrow \eta^j(x)$, for $j = k, d-k$, so

$$\Psi(\gamma^{-n_m}) \circ f_{\rho_m, v_m, x_m} \circ \Psi(\gamma^{n_m}) \rightarrow \hat{f}.$$

Notice, that if $X \in \Xi^{d-k}|_{(\rho_m, v_m)}$, then since $\|\cdot\|$ is a canonical family of norms,

$$\|f_{\rho_m, v_m, x_m}(X)\|_{(\rho_m, v_m)} = \left\| \left(\Psi_m(\gamma^{-n_m}) \circ f_{\rho_m, v_m, x_m} \circ \Psi_m(\gamma^{n_m}) \right) \left(\Psi_m(\gamma^{-n_m})(X) \right) \right\|_{(\rho_m, \gamma^{-n_m}(v_m))}$$

and $\|\Psi(\gamma^{-n_m})(X)\|_{(\rho_m, \gamma^{-n_m}(v_m))} = \|X\|_{(\rho_m, v_m)}$. Thus

$$\lim_{m \rightarrow \infty} \|f_{\rho_m, v_m, x_m}\|_{(\rho_m, v_m)} = \lim_{m \rightarrow \infty} \left\| \Psi_m(\gamma^{-n_m}) \circ f_{\rho_m, v_m, x_m} \circ \Psi_m(\gamma^{n_m}) \right\|_{(\rho_m, \gamma^{-n_m}(v_m))} = \left\| \hat{f} \right\|_{(\rho, v)} \neq 0$$

and we have a contradiction. This completes the proof of Lemma 8.6 and hence the proof of (2).

The limits maps vary analytically. It remains to prove the analytic variation of the limit maps. First suppose that $\mathbb{K} = \mathbb{R}$. The general strategy is to complexify and then exploit the fact that locally uniform limits of complex analytic functions are complex analytic.

Suppose that $h : M \rightarrow \text{Hom}_{\text{tp}}(\rho_0)$ is a real analytic map and every representation in $h(M)$ is P_k -Anosov.

If $\rho : \Gamma \rightarrow \mathbf{SL}(d, \mathbb{R})$ is P_k -Anosov, we may compose with the inclusion map $\iota_2 : \mathbf{SL}(d, \mathbb{R}) \rightarrow \mathbf{SL}(d, \mathbb{C})$ to obtain a P_k -Anosov representation $\rho^{\mathbb{C}} = \iota_2 \circ \rho : \Gamma \rightarrow \mathbf{SL}(d, \mathbb{C})$. Fix generators g_1, \dots, g_N of Γ and view $\text{Hom}(\Gamma, \mathbf{SL}(d, \mathbb{C}))$ as a subset of $\mathbf{SL}(d, \mathbb{C})^N$. We can then view h as a map $h : M \rightarrow \mathbf{SL}(d, \mathbb{C})^N$. We can also realize M as a totally real submanifold of a complex manifold $M^{\mathbb{C}}$ and then extend h to a complex analytic map $h : M^{\mathbb{C}} \rightarrow \mathbf{SL}(d, \mathbb{C})^N$. Notice that $h(M)$ and $h(M^{\mathbb{C}})$ have the same Zariski closure in $\mathbf{SL}(d, \mathbb{C})^N$.

We claim, after possibly shrinking $M^{\mathbb{C}}$, that $h(M^{\mathbb{C}}) \subset \text{Hom}_{\text{tp}}(\rho_0^{\mathbb{C}})$. For any $\alpha \in \mathbf{SL}(d, \mathbb{C})$, the set

$$\{g \in \mathbf{SL}(d, \mathbb{C}) : g \text{ is conjugate to } \alpha\}$$

is locally closed (i.e. open in its closure) in the Zariski topology, see for instance [10, Theorem 3.6]. This implies that $\text{Hom}_{\text{tp}}(\rho_0^{\mathbb{C}})$ is itself a locally closed set in the Zariski topology on $\text{SL}(d, \mathbb{C})^N$. Then since $h(M) \subset \text{Hom}_{\text{tp}}(\rho_0^{\mathbb{C}})$, by shrinking $M^{\mathbb{C}}$ we may assume that $h(M^{\mathbb{C}}) \subset \text{Hom}_{\text{tp}}(\rho_0^{\mathbb{C}})$.

Since every representation in $h(M)$ is P_k -Anosov, by shrinking $M^{\mathbb{C}}$ again if necessary, we may assume that every representation in $h(M^{\mathbb{C}})$ is also P_k -Anosov. Thus, if we can prove that for any $x \in \Lambda(\Gamma)$, the map $u \mapsto \xi_{h(u)}^k(x)$ from $M^{\mathbb{C}}$ to $\text{Gr}_k(\mathbb{C}^d)$ is complex analytic, then its restriction to M is real analytic.

If γ is a hyperbolic element and $\rho \in h(M^{\mathbb{C}})$, then $\rho(\gamma)$ is P_k -proximal and $\xi_{\rho}^k(\gamma^+)$ is the attracting k -plane of $\rho(\gamma)$. Then it follows from standard results in the perturbation theory of linear operators, see, for example, [20, Chapter 6], that the function from $M^{\mathbb{C}}$ to $\text{Gr}_k(\mathbb{C}^d)$ given by $u \mapsto \xi_{h(u)}^k(\gamma^+)$ is complex analytic. If $x \in \Lambda(\Gamma)$, then there exists a sequence $\{\gamma_n\}$ of hyperbolic elements of Γ , so that $\gamma_n^+ \rightarrow x$. Then, since the map

$$(u, y) \in M^{\mathbb{C}} \times \Lambda(\Gamma) \mapsto \xi_{h(u)}^k(y) \in \text{Gr}_k(\mathbb{C}^d)$$

is continuous, the function $u \mapsto \xi_{h(u)}^k(x)$ is a locally uniform limit of complex analytic functions, hence complex analytic. This completes the proof of (3) in the case when $\mathbb{K} = \mathbb{R}$.

The case when $\mathbb{K} = \mathbb{C}$ case follows by simply repeating the argument in the previous paragraph. \square

9. POSITIVE REPRESENTATIONS IN THE SENSE OF FOCK-GONCHAROV

In Fock and Goncharov's work [14], they define positive representations in the following way. Suppose that $\Gamma_0 \subset \text{PSL}(2, \mathbb{R})$ is a discrete group and $\Gamma_0 \backslash \mathbb{H}^2$ is a non-compact, finite area hyperbolic surface. Recall that $\gamma \in \Gamma_0$ is *peripheral* if it is represented by a curve which may be freely homotoped off of every compact subset of $S = \Gamma \backslash \mathbb{H}^2$. Let $\Lambda_p(\Gamma_0)$ be the set of fixed points of the peripheral elements in Γ_0 (in this case, all of which are parabolic). Notice that $\Lambda_p(\Gamma_0)$ inherits two natural cyclic orders as a subset of $\partial\mathbb{H}^2$. A representation $\rho : \Gamma_0 \rightarrow \text{SL}(d, \mathbb{R})$ is *positive* if there is a positive, ρ -equivariant map $\zeta : \Lambda_p(\Gamma_0) \rightarrow \mathcal{F}(\mathbb{R}^d)$. Notice that, with Fock and Goncharov's definition, every Hitchin representation of a convex cocompact, but not cocompact, Fuchsian group Γ_0 is also a positive representation of a lattice Γ . (If $\Gamma \backslash \mathbb{H}^2$ is homeomorphic to the interior of a compact surface S , then Γ is a finite area uniformization of the interior of S .) In this case, every peripheral element is mapped to a loxodromic element and there are many different positive ρ -equivariant maps from $\Lambda_p(\Gamma)$ to $\mathcal{F}(\mathbb{R}^d)$, corresponding to a choice of fixed point for the (unique) fixed point of each peripheral element of Γ .

Motivated by their definition, we define the notion of a positive type preserving representation in the following way. Let $\Gamma \subset \text{PSL}(2, \mathbb{R})$ be a geometrically finite group. Then $\gamma \in \Gamma$ is peripheral if it is either unipotent or it is a hyperbolic element whose fixed points both lie in the boundary of $\Lambda(\Gamma)$.

Definition 9.1. A representation $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{R})$ of a geometrically finite Fuchsian group is *positive type preserving* if

- $\rho(\gamma)$ is weakly unipotent for every parabolic $\gamma \in \Gamma$, and
- there is a positive, ρ -equivariant map $\zeta : \Lambda_p(\Gamma) \rightarrow \mathcal{F}(\mathbb{R}^d)$.

Observe that if $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{R})$ is a positive type preserving representation, then $\rho \circ f_* : \Gamma_0 \rightarrow \text{SL}(d, \mathbb{R})$ is a positive representation for all homeomorphisms $f : \Gamma_0 \backslash \mathbb{H}^2 \rightarrow \Gamma \backslash \mathbb{H}^2$.

It is clear that every Hitchin representation from Γ to $\text{SL}(d, \mathbb{R})$ is a positive type preserving representation. We show that the converse is also true.

Theorem 9.2. *If $\Gamma \subset \text{PSL}(2, \mathbb{R})$ is geometrically finite group, then every positive type preserving representation $\rho : \Gamma \rightarrow \text{SL}(d, \mathbb{R})$ is Hitchin.*

Let ρ be a positive, type preserving representation, and let $\zeta : \Lambda_p(\Gamma) \rightarrow \mathcal{F}(\mathbb{R}^d)$ denote a positive, ρ -equivariant map. We make use of the following well-known fact which is implicit in Fock-Goncharov [14], see Kim-Tan-Zhang [21, Observation 3.18] for details. It may be viewed as a generalization of the fact that every bounded monotone sequence in \mathbb{R} is convergent.

Proposition 9.3. *Let $\{F_n\}$ be a sequence of flags in $\mathcal{F}(\mathbb{R}^d)$ and $H_1, H_2 \in \mathcal{F}(\mathbb{R}^d)$ such that $(F_1, \dots, F_n, H_1, H_2)$ is positive for all n . Then the sequence $\{F_n\}$ converges to a flag $F_{\infty} \in \mathcal{F}(\mathbb{R}^d)$, and $(F_1, \dots, F_n, F_{\infty}, H_2)$ is positive for all n .*

We fix for the remainder of the section, one of the two natural cyclic orders on $\partial\mathbb{H}^2$. With this convention, it is natural to define one-sided convergence of sequences.

Definition 9.4. A sequence $\{x_n\}$ in $\partial\mathbb{H}^2$ converges to $x \in \partial\mathbb{H}^2$ *in the positive direction* (respectively, *in the negative direction*) if $\{x_n\}$ converges to x , and there exists $N > 0$ such that

$$x_N < x_{N+1} < \cdots < x \quad (\text{respectively, } x_N > x_{N+1} > \cdots > x)$$

For short, we write $x_n \nearrow x$ (respectively, $x_n \searrow x$) if $\{x_n\}$ converges to x in the positive direction (respectively, in the negative direction). If $\{x_n\}$ converges to x in either the positive direction or the negative direction, we say that $\{x_n\}$ converges *monotonically* to x .

Definition 9.5. Let $x \in \Lambda(\Gamma)$.

- If there are sequences in $\Lambda_p(\Gamma)$ that converge to x from the positive direction (respectively, from the negative direction), set

$$\xi_+(x) := \lim_{y \in \Lambda_p(\Gamma), y \nearrow x} \zeta(y) \quad \left(\text{respectively, } \xi_-(x) := \lim_{y \in \Lambda_p(\Gamma), y \searrow x} \zeta(y) \right).$$

- If there are no sequences in $\Lambda_p(\Gamma)$ that converge to x in the positive direction (respectively, in the negative direction), then there necessarily are sequences in $\Lambda_p(\Gamma)$ that converge to x in the negative direction (respectively, in the positive direction). Thus, we may set

$$\xi_+(x) := \xi_-(x) \quad (\text{respectively, } \xi_-(x) := \xi_+(x)).$$

Since $\Lambda_p(\Gamma)$ is dense in $\Lambda(\Gamma)$, Proposition 9.3 implies that these limit maps are well-defined. We refer to the maps $\xi_+ : \Lambda(\Gamma) \rightarrow \mathcal{F}(\mathbb{R}^d)$ (respectively, $\xi_- : \Lambda(\Gamma) \rightarrow \mathcal{F}(\mathbb{R}^d)$) as the *plus limit map* (respectively, *minus limit map*).

Since ζ is ρ -equivariant, both ξ_+ and ξ_- are ρ -equivariant. We next check that they satisfy the following positivity property, which implies in particular, that they are both positive.

Proposition 9.6. *Let $x_1 < x_2 < \cdots < x_k$ be points in $\Lambda(\Gamma)$, and let $s_1, \dots, s_k \in \{+, -\}$. Then*

$$(\xi_{s_1}(x_1), \xi_{s_2}(x_2), \dots, \xi_{s_k}(x_k))$$

is a positive tuple of flags.

Proof. For each $i = 1, \dots, k$, let $q_i, q'_i \in \Lambda_p(\Gamma)$ be points satisfying the following conditions:

- $q_i = x_i$ if there are no sequences in $\Lambda_p(\Gamma)$ that converge to x_i in the positive direction,
- $q'_i = x_i$ if there are no sequences in $\Lambda_p(\Gamma)$ that converge to x_i in the negative direction,
- $x_i < q'_i < q_{i+1}$ if there is a sequence in $\Lambda_p(\Gamma)$ that converges to x_i in the negative direction, and
- $q'_{i-1} < q_i < x_i$ if there is a sequence in $\Lambda_p(\Gamma)$ that converges to x_i in the positive direction.

Here, arithmetic in the subscripts is done modulo k .

By the definition of ξ_{\pm} , for each $i = 1, \dots, k$, there is a sequence $(y_{i,n})_{n \geq 1}$ in $\Lambda_p(\Gamma)$ that converges monotonically to x_i , and satisfies $\lim_{n \rightarrow \infty} \zeta(y_{i,n}) = \xi_{s_i}(x_i)$. By passing to the tail of the sequences $(y_{i,n})_{n \geq 1}$, we may assume that for all n ,

$$q_1 < y_{1,n} < q'_1 < q_2 < y_{2,n} < q'_2 < \cdots < q_k < y_{k,n} < q'_k,$$

which implies that

$$(\zeta(q_1), \zeta(y_{1,n}), \zeta(q'_1), \zeta(q_2), \zeta(y_{2,n}), \zeta(q'_2), \dots, \zeta(q_k), \zeta(y_{k,n}), \zeta(q'_k))$$

is a positive tuple of flags in $\mathcal{F}(\mathbb{R}^d)$. Our result then follows from Proposition 9.3. \square

We next prove that ξ_{\pm} satisfy an analogue of the strongly dynamics-preserving property of Anosov limit maps. In particular, if $\xi_+ = \xi_-$, then it implies that ξ_{\pm} is strongly dynamics-preserving. In fact, the proof mimics the proof of Theorem 7.1.

Proposition 9.7. *Suppose $\{\gamma_n\}$ is a sequence in Γ with $\gamma_n \rightarrow x \in \Lambda(\Gamma)$ and $\gamma_n^{-1} \rightarrow y \in \Lambda(\Gamma)$. Then after passing to a subsequence, there exists $s_1, s_2 \in \{+, -\}$ such that*

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)(F) = \xi_{s_1}(x)$$

for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi_{s_2}(y)$.

Proof. We first suppose that $x \neq y$. By passing to the tail of the sequence, we may assume that each γ_n is hyperbolic. Then $\gamma_n^+ \rightarrow x$, $\gamma_n^- \rightarrow y$, and $\gamma_n(z) \rightarrow x$ for all $z \in \Lambda(\Gamma) - \{y\}$.

Since $\Lambda_p(\Gamma)$ is infinite, there are points $a, b \in \Lambda_p(\Gamma) - \{x, y\}$ such that

- either $x < a < b < y$ or $y < b < a < x$, and
- up to taking subsequences, the sequences $\{\gamma_n(a)\}$ and $\{\gamma_n(b)\}$ both converge monotonically to x , and from the same direction.

Therefore,

$$\lim_{n \rightarrow \infty} \zeta(\gamma_n(a)) = \xi_{s_1}(x) = \lim_{n \rightarrow \infty} \zeta(\gamma_n(b))$$

for some $s_1 \in \{+, -\}$. For each $n \geq 1$, choose a point $c_n \in \Lambda_p(\Gamma) \setminus \{\gamma_n^-\}$ such that $c_n \rightarrow y$ and $\gamma_n(c_n) \rightarrow y$. Passing to a further subsequence, we can assume that $\{\gamma_n(c_n)\}$ converges monotonically to y . Thus,

$$\lim_{n \rightarrow \infty} \zeta(\gamma_n(c_n)) = \xi_{s_3}(y)$$

for some $s_3 \in \{+, -\}$.

Now, consider the open sets (defined in Definition 7.2)

$$\mathcal{O}_n := \mathcal{O}(\zeta(a), \zeta(b), \zeta(c_n))$$

for all n . Since $c_n \rightarrow y$, and either $a < b < y$ or $b < a < y$, there exists $N > 0$ such that either $a < b < c_n$ for all $n \geq N$, or $b < a < c_n$ for all $n \geq N$.

We may then argue, exactly as in the proof of Theorem 7.1, that

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)(F) = \xi_{s_1}(x)$$

for all $F \in W$. Repeating the same argument with γ_n^{-1} , we see that there exists $s_2 \in \{+, -\}$ and an open set $W' \subset \mathcal{F}(\mathbb{R}^d)$ where

$$\lim_{n \rightarrow \infty} \rho(\gamma_n^{-1})(F) = \xi_{s_2}(y)$$

for all $F \in W'$. Hence, we may apply Lemma 2.2 to deduce the proposition when $x \neq y$.

If $x = y$, pick $\eta \in \Gamma$ such that $z := \eta^{-1}(x) \neq x$. Then $\gamma_n \eta \rightarrow x$, $(\gamma_n \eta)^{-1} \rightarrow z \neq x$. By the first part, there exists $s_1, s_2 \in \{+, -\}$ such that $\rho(\gamma_n \eta)(F) \rightarrow \xi_{s_1}(x)$ for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi_{s_2}(z)$. Equivalently, $\rho(\gamma_n)(F) \rightarrow \xi_{s_1}(x)$ for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi_{s_2}(x)$. \square

Given $x \in \Lambda(\Gamma) - \Lambda_p(\Gamma)$, it will be useful to construct a sequence in Γ and a pair of points in $\Lambda(\Gamma)$, so that the orbits of the points under the sequence of elements of Γ approach x from opposite sides.

Lemma 9.8. *If $x \in \Lambda(\Gamma) - \Lambda_p(\Gamma)$, then there exists a sequence $\{\gamma_n\}$ in Γ , a point $y \in \Lambda(\Gamma)$, and points $a, b \in \Lambda(\Gamma) - \{y\}$ so that:*

- (1) $\gamma_n \rightarrow x$,
- (2) $\gamma_n^{-1} \rightarrow y$,
- (3) $\gamma_n(a) \nearrow x$ and $\gamma_n(b) \searrow x$.

Proof. Let $r : \mathbb{R} \rightarrow \mathbb{H}^2$ be a geodesic joining a point $w \in \Lambda(\Gamma) - \{x\}$ to x . Since x is a conical limit point, there exists sequences $\{t_n\} \subset \mathbb{R}$ and $\{\gamma_n\} \subset \Gamma$ so that $t_n \rightarrow \infty$ and $\{d_{\mathbb{H}^2}(r(t_n), \gamma_n(r(0)))\}$ is bounded. Therefore, after passing to a subsequence, $v_n = (\gamma_n^{-1} \circ r)'(t_n)$ converges to a vector $v \in \mathbf{U}(\Gamma)$. Notice that $\gamma_n(v_n^+) = x$ and $\gamma_n(v_n^-) = w$ for all n .

If there is a point $a \in \Lambda(\Gamma)$ such that $v^- < a < v^+$, then $v_n^- < a < v_n^+$ for sufficiently large n . It follows that $w < \gamma_n(a) < x$ for sufficiently large n . Since $\gamma_n(a) \rightarrow x$, it follows that $\gamma_n(a) \nearrow x$. On the other hand, if there are no points $a \in \Lambda(\Gamma)$ such that $v^- < a < v^+$, then v^+ and v^- are the fixed points of

a hyperbolic, peripheral element. In particular, $\gamma_n^{-1}(x) \neq v^+$ for any n . Since $\gamma_n^{-1}(x) \rightarrow v^+$, this implies that $\gamma_n^{-1}(x) \searrow v^+$, so $\gamma_n(v^+) \nearrow x$. In either case, there is some $a \in \Lambda(\Gamma)$ such that $\gamma_n(a) \nearrow x$.

Similarly, we may find a point $b \in \Lambda(\Gamma)$ so that $\gamma_n(b) \searrow x$. \square

Lemma 9.8 allows us to complete the proof of Theorem 9.2.

Proof of Theorem 9.2. We first notice that it suffices to prove that $\xi_+ = \xi_-$ and $\xi_+|_{\Lambda_p(\Gamma)} = \zeta$. Indeed, if we can do so, then we may set $\xi : \Lambda(\Gamma) \rightarrow \mathcal{F}(\mathbb{R}^d)$ to be the map given by $\xi(x) := \xi_{\pm}(x)$. Proposition 9.6 implies that ξ is positive, while Proposition 9.3 implies that ξ is continuous.

The proof proceeds in three cases. In the first case, we assume that x is not the fixed point of a peripheral element. If x is the fixed point of a peripheral element, then x is the fixed point of either a parabolic element or a peripheral hyperbolic element.

Case 1: $x \in \Lambda(\Gamma) - \Lambda_p(\Gamma)$. Let $a, b, y \in \Lambda(\Gamma)$ and $\{\gamma_n\}$ be as in Lemma 9.8. By Proposition 9.7, we can pass to a subsequence so that there exists $s_1, s_2 \in \{+, -\}$ so that

$$\rho(\gamma_n)(F) \rightarrow \xi_{s_1}(x)$$

for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi_{s_2}(y)$. By Proposition 9.6, the flags $\xi_+(a)$ and $\xi_+(b)$ are both transverse to $\xi_{s_2}(y)$. So

$$(24) \quad \xi_{s_1}(x) = \lim_{n \rightarrow \infty} \rho(\gamma_n)(\xi_+(a)) = \lim_{n \rightarrow \infty} \xi_+(\gamma_n(a)) = \xi_+(x)$$

and

$$(25) \quad \xi_{s_1}(x) = \lim_{n \rightarrow \infty} \rho(\gamma_n)(\xi_+(b)) = \lim_{n \rightarrow \infty} \xi_+(\gamma_n(b)) = \xi_-(x).$$

The second equality in (24) and (25) holds because $\gamma_n \in \Gamma$ for all n , and the last equality in (24) and (25) is a consequence of Proposition 9.3. Thus $\xi_+(x) = \xi_-(x)$.

Case 2: $x \in \Lambda_p(\Gamma)$ is the fixed point of a parabolic element $\alpha \in \Gamma$. As in the proof of Theorem 1.4, Lemma 7.6 implies that $\rho(\alpha)$ has a unique fixed flag in $\mathcal{F}(\mathbb{R}^d)$. It follows that $\xi_+(x) = \xi_-(x) = \zeta(x)$ is this unique fixed flag.

Case 3: $x \in \Lambda_p(\Gamma)$ is the fixed point of a hyperbolic peripheral element $\gamma \in \Gamma$. By replacing γ with γ^{-1} we can assume that $x = \gamma^+$. Then either $\gamma^+ < a < \gamma^-$ for all $a \in \Lambda_p(\Gamma) - \{\gamma^+, \gamma^-\}$ or $\gamma^- < a < \gamma^+$ for all $a \in \Lambda_p(\Gamma) - \{\gamma^+, \gamma^-\}$. Also, by definition, $\xi_+(x) = \xi_-(x)$.

We now show $\xi_+(x) = \zeta(x)$. Since $\gamma^n \rightarrow \gamma^+$ and $\gamma^{-n} \rightarrow \gamma^-$, Proposition 9.7 implies that there is an increasing sequence $\{m_n\}$ of integers so that

$$\rho(\gamma^{m_n})(F) \rightarrow \xi_+(x)$$

for all $F \in \mathcal{F}(\mathbb{R}^d)$ transverse to $\xi_+(\gamma^-)$. It follows that $\rho(\gamma)$ is loxodromic, and that $\xi_+(\gamma^+)$ and $\xi_+(\gamma^-)$ are respectively the attracting and repelling fixed flag of $\rho(\gamma)$.

To finish the proof, it is sufficient to show that $\zeta(\gamma^+)$ is also the attracting fixed flag of $\rho(\gamma)$. Let $\{x_n\}$ be a sequence in $\Lambda_p(\Gamma)$ that converges monotonically to γ^- . Since ζ is positive, the tuple

$$(\zeta(\gamma^+), \zeta(x_1), \dots, \zeta(x_n), \zeta(\gamma^-))$$

is positive for all n . Since $\xi_+(\gamma^-) = \lim_{n \rightarrow \infty} \zeta(x_n)$, Proposition 9.3 implies that

$$(\zeta(\gamma^+), \zeta(x_1), \dots, \zeta(x_n), \xi_+(\gamma^-))$$

is positive for all n . In particular, $\zeta(\gamma^+)$ and $\xi_+(\gamma^-)$ are transverse. Since $\zeta(\gamma^+)$ is fixed by $\rho(\gamma)$ and $\xi_+(\gamma^-)$ is the repelling fixed point of $\rho(\gamma)$, it follows that $\zeta(\gamma^+)$ is the attracting fixed flag of $\rho(\gamma)$. \square

APPENDIX A. CONSTRUCTING CUSP REPRESENTATIONS

In this appendix, we prove Proposition 3.6. Suppose $g \in \mathrm{SL}(d, \mathbb{K})$ is weakly unipotent and let

$$u = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

We will construct a representation $\Psi : \mathrm{SL}(2, \mathbb{R}) \rightarrow \mathrm{SL}(d, \mathbb{K})$ where $\Psi(u) = g_u$ and g_{ss} commutes with the elements of $\Psi(\mathrm{SL}(2, \mathbb{R}))$.

First suppose that $\mathbb{K} = \mathbb{C}$. Then using the Jordan normal form there exist $p \in \mathrm{SL}(d, \mathbb{C})$, integers $d_1 \geq d_2 \geq \dots \geq d_m > 0$, and complex numbers $\lambda_1, \dots, \lambda_m \in \mathbb{S}^1$ so that $d_1 + \dots + d_m = d$,

$$pg_u p^{-1} = \bigoplus_{j=1}^m \tau_{d_j}(u) \quad \text{and} \quad pg_{ss} p^{-1} = \bigoplus_{j=1}^m \lambda_j \mathrm{id}_{d_j}.$$

It follows that Ψ defined by

$$\Psi(\beta) = p^{-1} \left(\bigoplus_{i=1}^m \tau_{d_i}(\beta) \right) p$$

for all $\beta \in \mathrm{SL}(2, \mathbb{R})$ has the desired properties.

Next suppose that $\mathbb{K} = \mathbb{R}$. Given two matrices A, B we will let $A \otimes B$ denote the *Kronecker product*, that is if $A = [a_{ij}]$ is an m -by- n matrix, then

$$A \otimes B = \begin{pmatrix} a_{11}B & \cdots & a_{1n}B \\ \vdots & & \vdots \\ a_{m1}B & \cdots & a_{mn}B \end{pmatrix}.$$

Also given $\theta \in \mathbb{R}$ let

$$M(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}.$$

Then using the real Jordan normal form there exist $p \in \mathrm{SL}(d, \mathbb{R})$, integers $d_1 \geq d_2 \geq \dots \geq d_{m+n} > 0$ and numbers $\lambda_1, \dots, \lambda_m \in \{-1, 1\}$, $\theta_{m+1}, \dots, \theta_{m+n} \in \mathbb{R}$ so that $d_1 + \dots + d_m + 2d_{m+1} + \dots + 2d_{m+n} = d$,

$$pg_u p^{-1} = \left[\bigoplus_{j=1}^m \tau_{d_j}(u) \right] \oplus \left[\bigoplus_{j=m+1}^{m+n} \tau_{d_j}(u) \otimes \mathrm{id}_2 \right]$$

and

$$pg_{ss} p^{-1} = \left[\bigoplus_{j=1}^m \lambda_j \mathrm{id}_{d_j} \right] \oplus \left[\bigoplus_{j=m+1}^{m+n} \mathrm{id}_{d_j} \otimes M(\theta_j) \right].$$

By the multiplicative property of the Kronecker product

$$\left[\tau_{d_j}(\beta) \otimes \mathrm{id}_2 \right] \left[\mathrm{id}_{d_j} \otimes M(\theta_j) \right] = \left[\mathrm{id}_{d_j} \otimes M(\theta_j) \right] \left[\tau_{d_j}(\beta) \otimes \mathrm{id}_2 \right] = \tau_{d_j}(\beta) \otimes M(\theta_j)$$

for all $\beta \in \mathrm{SL}(2, \mathbb{R})$ and so it follows that Ψ defined by

$$\Psi(\beta) = p^{-1} \left(\left[\bigoplus_{j=1}^m \tau_{d_j}(\beta) \right] \oplus \left[\bigoplus_{j=m+1}^{m+n} \tau_{d_j}(\beta) \otimes \mathrm{id}_2 \right] \right) p$$

has the desired properties.

APPENDIX B. ANOSOV REPRESENTATIONS INTO SEMISIMPLE LIE GROUPS

In this appendix, we develop a more general theory of Anosov representations of geometrically finite Fuchsian groups into a semisimple Lie group \mathbf{G} with respect to a parabolic subgroup P^+ . We extend results of Guichard-Wienhard [16, Prop. 4.3] (see also Guéritaud-Guichard-Wienhard [15, Section 3]) to show that there exists an irreducible representation $\psi : \mathbf{G} \rightarrow \mathrm{SL}(d, \mathbb{R})$ so that a representation into \mathbf{G} is Anosov with respect to P^+ if and only if its composition with ψ is P_1 -Anosov. This will allow us to immediately recover generalizations of all the results we obtained for linear Anosov representations.

For the rest of the section, we will assume that \mathbf{G} is a semisimple Lie group of non-compact type with finite center, denote its Lie algebra by \mathfrak{g} , and let $\mathrm{ad} : \mathfrak{g} \rightarrow \mathfrak{sl}(\mathfrak{g})$ and $\mathrm{Ad} : \mathbf{G} \rightarrow \mathrm{SL}(\mathfrak{g})$ be the adjoint representations.

Fix a parabolic subgroup $P^+ \subset \mathbf{G}$ and an opposite parabolic subgroup $P^- \subset \mathbf{G}$ and let $\mathcal{F}^\pm := \mathbf{G}/P^\pm$ be the associated flag varieties. If $\rho : \Gamma \rightarrow \mathbf{G}$ is a representation, we define the bundles

$$\widehat{B}_\rho^\pm = \Gamma \backslash (\mathrm{U}(\Gamma) \times \mathcal{F}^\pm) \quad \text{and} \quad \widehat{V}_\rho^\pm = \Gamma \backslash (\mathrm{U}(\Gamma) \times T\mathcal{F}^\pm),$$

where $T\mathcal{F}^\pm$ is the tangent bundle of \mathcal{F}^\pm . Observe that V_ρ^\pm is a vector bundle over B_ρ^\pm of rank $\dim(\mathcal{F}^\pm)$.

The geodesic flow on $\mathbf{U}(\Gamma)$ extends to flows on $\mathbf{U}(\Gamma) \times \mathcal{F}^\pm$ and $\mathbf{U}(\Gamma) \times T\mathcal{F}^\pm$ whose action is trivial on the second factor. These in turn descends to flows on \widehat{B}_ρ^\pm and \widehat{V}_ρ^\pm which covers the geodesic flow on $\widehat{\mathbf{U}}(\Gamma)$. We use ϕ_t to denote all of these flows.

We say that a map

$$\xi = (\xi^+, \xi^-) : \Lambda(\Gamma) \rightarrow \mathcal{F}^+ \times \mathcal{F}^-$$

is *transverse* if whenever $x \neq y \in \Lambda(\Gamma)$, the pair $(\xi^+(x), \xi^-(y))$ lies in the unique open \mathbf{G} -orbit in $\mathcal{F}^+ \times \mathcal{F}^-$. Given a transverse, ρ -equivariant, continuous map ξ we define sections

$$\sigma_\xi^\pm : \mathbf{U}(\Gamma) \rightarrow B_\rho^\pm = \mathbf{U}(\Gamma) \times \mathcal{F}^\pm$$

given by $\sigma_\xi^\pm(v) = (v, \xi^\pm(v^\pm))$. Since ξ is ρ -equivariant, σ_ξ^\pm descend to sections $\widehat{\sigma}_\xi^\pm : \widehat{\mathbf{U}}(\Gamma) \rightarrow \widehat{B}_\rho^\pm$.

Definition B.1. A representation $\rho : \Gamma \rightarrow \mathbf{G}$ is *P^\pm -Anosov* if the following hold:

- (1) There exists a ρ -equivariant, continuous, transverse map $\xi = (\xi^+, \xi^-) : \Lambda(\Gamma) \rightarrow \mathcal{F}^+ \times \mathcal{F}^-$.
- (2) For some family of norms $\|\cdot\|$ on the fibers of $\widehat{V}_\rho^- \rightarrow \widehat{B}_\rho^-$, the pullback of the flow ϕ_t , also denoted ϕ_t , is uniformly contracting on $(\widehat{\sigma}_\xi^-)^* \widehat{V}_\rho^-$.

We refer to any such map ξ as the *P^\pm -Anosov limit map* for ρ .

Remark B.2. In the case when $\mathbf{G} = \mathbf{SL}(d, \mathbb{K})$ and $(P^+, P^-) = (P_k, P_{d-k})$, note that $\mathcal{F}^+ = \text{Gr}_k(\mathbb{K}^d)$ and $\mathcal{F}^- = \text{Gr}_{d-k}(\mathbb{K}^d)$. Then for any pair $(F, G) \in \text{Gr}_k(\mathbb{K}^d) \times \text{Gr}_{d-k}(\mathbb{K}^d)$, there is a natural identification $T_F \text{Gr}_k(\mathbb{K}^d) \simeq \text{Hom}(F, G)$ and $T_G \text{Gr}_{d-k}(\mathbb{K}^d) \simeq \text{Hom}(G, F)$. Thus, the pullback bundles $(\widehat{\sigma}_\xi^+)^*(\widehat{V}_\rho^+)$ and $(\widehat{\sigma}_\xi^-)^*(\widehat{V}_\rho^-)$ are canonically identified with $\text{Hom}(\widehat{\Theta}^k, \widehat{\Xi}^{d-k})$ and $\text{Hom}(\widehat{\Xi}^{d-k}, \widehat{\Theta}^k)$ respectively, where $\widehat{\Theta}^k$ and $\widehat{\Xi}^{d-k}$ are the sub-bundles of \widehat{E}_ρ that lift to sub-bundles Θ^k and Ξ^{d-k} of E_ρ with the defining property $\Theta^k|_v = \xi^+(v^+)$ and $\Xi^{d-k}|_v = \xi^-(v^-)$. Thus, in this case, Definition B.1 agrees with Definition 3.1, see Proposition 3.3.

If $\psi : \mathbf{G} \rightarrow \mathbf{SL}(V)$ is a finite-dimensional irreducible representation, we say that ψ *intertwines* (P^+, P^-) and (P_1, P_{d-1}) (where $d = \dim(V)$) if $V = L_0 \oplus W_0$ where L_0 is a line and W_0 is a hyperplane and

$$P^+ = \{g \in \mathbf{G} : \psi(g)(L_0) = L_0\} \quad \text{and} \quad P^- = \{g \in \mathbf{G} : \psi(g)(W_0) = W_0\}.$$

In this case, one may define embeddings $\psi_+ : \mathcal{F}^+ \rightarrow \mathbb{P}(V)$ and $\psi_- : \mathcal{F}^- \rightarrow \text{Gr}_{\dim(V)-1}(V)$ by letting

$$\psi_+(gP^+) = \psi(g)(L_0) \quad \text{and} \quad \psi_-(gP^-) = \psi(g)(W_0).$$

The following result often allows one to reduce the general study of Anosov representations to the study of P_1 -Anosov representations into $\mathbf{SL}(d, \mathbb{R})$. For Anosov representations of word hyperbolic groups the analogous result is due to Guichard-Wienhard [16, Proposition 4.3].

Theorem B.3. *Let \mathbf{G} be a semisimple Lie group with finite center and let P^\pm be a pair of opposite parabolic subgroups. Suppose that $\psi : \mathbf{G} \rightarrow \mathbf{SL}(V)$ is a finite dimensional irreducible representation which intertwines (P^+, P^-) and (P_1, P_{d-1}) (where $d = \dim(V)$). Then a representation $\rho : \Gamma \rightarrow \mathbf{G}$ of a geometrically finite Fuchsian group Γ is P^\pm -Anosov if and only if $\psi \circ \rho$ is P_1 -Anosov.*

Moreover, if $\xi_\rho = (\xi_\rho^+, \xi_\rho^-)$ is a P^\pm -Anosov limit map for ρ , then $\xi_{\psi \circ \rho} = (\psi_+ \circ \xi_\rho^+, \psi_- \circ \xi_\rho^-)$ is the P_1 -Anosov limit map of $\psi \circ \rho$. In particular, the P^\pm -Anosov limit map of ρ is unique.

Intertwining representations are not hard to construct. If \mathfrak{n}^+ is the nilpotent radical of the Lie algebra of P^+ , and $n = \dim \mathfrak{n}^+$, then $\psi(g) = \wedge^n \text{Ad}(g)$ and $V = \text{Span}\{\psi(\mathbf{G})(\wedge^n \mathfrak{n}^+)\} \subset \wedge^n \mathfrak{g}$ intertwines P^\pm and (P_1, P_{d-1}) (where $d = \dim(V)$), see [16, Remark 4.12]. We obtain the following immediate corollary.

Corollary B.4. *Suppose that \mathbf{G} is a semisimple Lie group with finite center and P^\pm is a pair of opposite parabolic subgroups. There exists a finite-dimensional irreducible representation $\psi : \mathbf{G} \rightarrow \mathbf{SL}(V)$ so that a representation $\rho : \Gamma \rightarrow \mathbf{G}$ of a geometrically finite Fuchsian group Γ is P^\pm -Anosov if and only if $\psi \circ \rho$ is P_1 -Anosov.*

Corollary B.4 allows us to generalize our main results about linear Anosov representations into the general setting. As in the $\mathrm{SL}(d, \mathbb{K})$ case, if $\rho : \Gamma \rightarrow \mathbf{G}$ is a representation, let

$$\mathrm{Hom}_{\mathrm{tp}}(\rho) \subset \mathrm{Hom}(\Gamma, \mathbf{G})$$

be the space of representations $\sigma : \Gamma \rightarrow \mathbf{G}$ so that if $\alpha \in \Gamma$ is parabolic, then $\sigma(\alpha)$ is conjugate to $\rho(\alpha)$. Theorem 1.3 becomes:

Corollary B.5. *Suppose that \mathbf{G} is a semisimple Lie group with finite center, P^\pm is a pair of opposite parabolic subgroups of \mathbf{G} and Γ is a geometrically finite Fuchsian group. If $\rho : \Gamma \rightarrow \mathbf{G}$ is P^\pm -Anosov, then there exists an open neighborhood \mathcal{O} of ρ in $\mathrm{Hom}_{\mathrm{tp}}(\rho)$, so that*

- (1) *If $\rho \in \mathcal{O}$, then ρ is P^\pm -Anosov.*
- (2) *There exists $\alpha > 0$ so that if $\rho \in \mathcal{O}$, then its P^\pm -Anosov limit map ξ_ρ is α -Hölder.*
- (3) *If $\{\rho_u\}_{u \in M}$ is an analytic family of representations in \mathcal{O} and $z \in \Lambda(\Gamma)$, then the map from M to $\mathcal{F}^+ \times \mathcal{F}^-$ given by $u \rightarrow \xi_{\rho_u}(z)$ is analytic.*

Theorem 1.2 yields:

Corollary B.6. *Suppose that \mathbf{G} is a semisimple Lie group with finite center, P^\pm is a pair of opposite parabolic subgroups of \mathbf{G} and Γ is a geometrically finite Fuchsian group. A representation $\rho : \Gamma \rightarrow \mathbf{G}$ is P^\pm -Anosov if and only if there exists a ρ -equivariant, transverse, continuous, strongly dynamics preserving map $\xi = (\xi^k, \xi^{d-k}) : \Lambda(\Gamma) \rightarrow \mathcal{F}^+ \times \mathcal{F}^-$. Furthermore, ξ is the P^\pm -Anosov limit map.*

One can also obtain analogues of parts (1) and (2) of Theorem 1.1 where the roles of singular values and eigenvalues are played by roots acting on the Cartan and Jordan projections (see [15, Section 3] for a complete discussion). Part (4) of Theorem 1.1 remains true if we replace $X_d(\mathbb{K})$ with the symmetric space of \mathbf{G} .

Theorem B.3 will be a consequence of Theorem 1.2 and the following dynamical property of Anosov representations.

Lemma B.7. *Suppose $\rho : \Gamma \rightarrow \mathbf{G}$ is a P^\pm -Anosov representation of a geometrically finite Fuchsian group Γ with P^\pm -Anosov limit map $\xi = (\xi^+, \xi^-)$. If $\{\gamma_n\}$ is a sequence in Γ such that $\gamma_n \rightarrow x \in \Lambda(\Gamma)$ and $\gamma_n^{-1} \rightarrow y \in \Lambda(\Gamma)$, then*

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)(F) \rightarrow \xi^+(x)$$

for all $F \in G/P^+$ transverse to $\xi^-(y)$.

Delaying the proof of the lemma we prove Theorem B.3.

Proof of Theorem B.3. We make repeated use of the following observation which follows from [15, Prop 3.5]. We sketch an alternate proof.

Observation B.8. $(F, H) \in \mathcal{F}^+ \times \mathcal{F}^-$ are transverse if and only if $\psi_+(F)$ and $\psi_-(H)$ are transverse.

Sketch of proof. Fix maximal compact subgroups $K_1 \subset \mathbf{G} =: \mathbf{G}_1$ and $K_2 \subset \mathrm{SL}(V) =: \mathbf{G}_2$ so that $\psi(K_1) \subset K_2$. For $j = 1, 2$, we can fix \mathbf{G}_j -invariant Riemannian metrics on $X_j = \mathbf{G}_j/K_j$ so that the map $T_\psi : X_1 \rightarrow X_2$ given by $T_\psi(gK_1) = \psi(g)(K_2)$ is a totally geodesic isometric embedding (see [29, Chapter 2]). Then T_ψ extends to an embedding $T_\psi : X_1(\infty) \hookrightarrow X_2(\infty)$ of the CAT(0)-boundaries.

Fix $(F, H) = (g_+P^+, g_-P^-)$. Let $W_+, W_- \subset X_1(\infty)$ be the interior of the Weyl faces associated to $g_+P^+g_+^{-1}$ and $g_-P^-g_-^{-1}$ respectively, i.e.

$$W_\pm = \{x \in X_1(\infty) : \mathrm{Stab}_{\mathbf{G}}(x) = g_\pm P^\pm g_\pm^{-1}\}.$$

Also, let $\hat{W}_+, \hat{W}_- \subset X_2(\infty)$ denote the interior of the Weyl faces associated to the parabolic subgroups $\psi(g_+)P_1\psi(g_+)^{-1}$ and $\psi(g_-)P_{d-1}\psi(g_-)^{-1}$ respectively. Notice that

$$W_\pm = T_\psi^{-1}(\hat{W}_\pm)$$

since $P^+ = \psi^{-1}(P_1)$ and $P^- = \psi^{-1}(P_{d-1})$.

Next fix a maximal flat $\mathcal{F} \subset X_1$ with $W_+, W_- \subset \mathcal{F}(\infty)$ and fix some $p \in \mathcal{F}$. Then let $s_p : X_1 \rightarrow X_1$ denote the involutive isometry based at p . Then F and H are transverse if and only if $W_+ = s_p(W_-)$. Since $T_\psi : X_1 \rightarrow X_2$ is a totally geodesic isometric embedding, there exists a maximal flat $\hat{\mathcal{F}} \subset X_2$ with $T_\psi(\mathcal{F}) \subset \hat{\mathcal{F}}$. Then $\hat{W}_+, \hat{W}_- \subset \hat{\mathcal{F}}(\infty)$ and if $s_{\hat{p}} : X_2 \rightarrow X_2$ is the involutive isometry based at $\hat{p} := T_\psi(p)$, then $\psi(F)$ and $\psi(H)$ are transverse if and only if $\hat{W}_+ = s_{\hat{p}}(\hat{W}_-)$.

Since

$$T_\psi \circ s_p = s_{\hat{p}} \circ T_\psi$$

and any two distinct interiors of Weyl faces have trivial intersection:

$$\begin{aligned} W_+ = s_p(W_-) &\implies T_\psi(W_+) \subset \hat{W}_+ \cap s_{\hat{p}}(\hat{W}_-) \implies \hat{W}_+ = s_{\hat{p}}(\hat{W}_-) \implies T_\psi(s_p(W_-)) \subset \hat{W}_+ \\ &\implies s_p(W_-) \subset W_+ = T_\psi^{-1}(\hat{W}_+) \implies W_+ = s_p(W_-). \end{aligned}$$

So F and H are transverse if and only if $\psi_+(F)$ and $\psi_-(H)$ are transverse. \square

We first prove the reverse direction of Theorem B.3.

Lemma B.9. *If $\psi \circ \rho$ is P_1 -Anosov, then ρ is P^\pm -Anosov. Moreover, if ξ_ρ is the P^\pm -Anosov limit map of ρ and $\xi_{\psi \circ \rho}$ is the P_1 -Anosov limit map of $\psi \circ \rho$, then $\psi_\pm \circ \xi_\rho^\pm = \xi_{\psi \circ \rho}^\pm$.*

Proof. If γ is a hyperbolic element of Γ , then, since $\psi \circ \rho$ is P_1 -Anosov, $\psi(\rho(\gamma))$ is P_1 -biproximal and $\xi_{\psi \circ \rho}^+(\gamma^+)$ is the attracting eigenline of $\psi(\rho(\gamma))$. Since ψ is irreducible, there exists $F \in \mathcal{F}^+$ such that $\psi_+(F)$ is transverse to the repelling hyperplane of $\psi(\rho(\gamma))$, so

$$\xi_{\psi \circ \rho}^+(\gamma^+) = \lim_{n \rightarrow \infty} (\psi \circ \rho)(\gamma)^n(\psi_+(F)) = \lim_{n \rightarrow \infty} \psi_+(\rho(\gamma)^n(F)) \in \psi_+(\mathcal{F}^+).$$

Hence, $\xi_{\psi \circ \rho}^+(\Lambda(\Gamma)) \subset \psi_+(\mathcal{F}^+)$ because attracting fixed points of hyperbolic elements are dense in $\Lambda(\Gamma)$.

Since ψ_+ is a ψ -equivariant embedding, $\xi^+ = \psi_+^{-1} \circ \xi_{\psi \circ \rho}^+$ is well-defined, continuous and ρ -equivariant. Similarly, $\xi^- = \psi_-^{-1} \circ \xi_{\psi \circ \rho}^-$ is well-defined, continuous and ρ -equivariant. Observation B.8 implies that $\xi = (\xi^+, \xi^-)$ is transverse.

We consider the vector bundles \hat{V}_ρ^- and $\hat{V}_{\psi \circ \rho}^-$ over \hat{B}_ρ^- and $\hat{B}_{\psi \circ \rho}^-$ respectively. Notice that the map ψ_- induces a bundle embedding $\iota_\rho : \hat{V}_\rho^- \hookrightarrow \hat{V}_{\psi \circ \rho}^-$ which intertwines the flows on the two bundles. Since $\psi \circ \rho$ is P_1 -Anosov, there is a continuous family of norms on the fibers of the bundle $\hat{V}_{\psi \circ \rho}^- \rightarrow \hat{B}_{\psi \circ \rho}^-$ such that ψ_t is uniformly contracting on the pullback bundle $(\hat{\sigma}_{\xi_{\psi \circ \rho}^-})^* \hat{V}_{\psi \circ \rho}^-$. Equip the bundle $\hat{V}_\rho^- \rightarrow \hat{B}_\rho^-$ with the pullback of this norm via ι_ρ . Since ι_ρ intertwines the flows, we see that the flow is uniformly contracting on $\sigma_\xi^*(V_\rho^-) = (\iota_\rho)^* \left((\hat{\sigma}_{\xi_{\psi \circ \rho}^-})^* \hat{V}_{\psi \circ \rho}^- \right)$. Therefore, ρ is P^\pm -Anosov with P^\pm -Anosov limit map $\xi = (\xi^+, \xi^-)$. \square

We now prove the forward direction of Theorem B.3.

Lemma B.10. *If ρ is P^\pm -Anosov with P^\pm -Anosov limit maps ξ_ρ , then $\psi \circ \rho$ is P_1 -Anosov with P_1 -Anosov limit map $(\psi_+ \circ \xi_\rho^+, \psi_- \circ \xi_\rho^-)$.*

Proof. Let $\eta = (\eta^+, \eta^-) = (\psi_+ \circ \xi_\rho^+, \psi_- \circ \xi_\rho^-)$. Then η is continuous, $\psi \circ \rho$ -equivariant, and transverse (by Observation B.8). So, by Theorem 1.2, it suffices to show that η is strongly dynamics-preserving.

Consider a sequence $\{\gamma_n\}$ in Γ with $\gamma_n \rightarrow x \in \Lambda(\Gamma)$ and $\gamma_n^{-1} \rightarrow y \in \Lambda(\Gamma)$. Then, by Lemma B.7,

$$\lim_{n \rightarrow \infty} (\psi \circ \rho)(\gamma_n)(\psi_+(F)) = \eta^+(x)$$

for all $F \in \mathcal{F}^+$ transverse to $\eta^-(y)$. Similarly,

$$\lim_{n \rightarrow \infty} (\psi \circ \rho)(\gamma_n)(\psi_-(F)) = \eta^-(x)$$

for all $F \in \mathcal{F}^-$ transverse to $\eta^+(x)$.

Since ψ is irreducible, $\psi_+(\mathcal{F}^+)$ spans V and one can repeat the proof of Corollary 6.3 to show that η is strongly dynamics preserving. \square

It only remains to prove Lemma B.7.

Proof of Lemma B.7. Let \mathfrak{p}^\pm be the Lie algebra of P^\pm . Then there exists a Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, a Cartan subspace $\mathfrak{a} \subset \mathfrak{p}$, and an element $H_0 \in \mathfrak{a}$ so that

$$\mathfrak{p}^\pm = \mathfrak{g}_0 \oplus \bigoplus_{\alpha(\pm H_0) \geq 0} \mathfrak{g}_\alpha$$

where

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}_\alpha$$

is the root space decomposition associated to \mathfrak{a} . Let $\mathfrak{n}^\pm = \bigoplus_{\alpha(\pm H_0) > 0} \mathfrak{g}_\alpha$ and define

$$T : \mathfrak{n}^- \rightarrow \mathcal{F}^+ \quad \text{where} \quad T(X) = e^X P^+$$

The map T have the following properties.

Observation B.11.

- (1) $T(\mathfrak{n}^-) = \{F \in \mathcal{F}^+ : F \text{ is transverse to } P^-\}$.
- (2) If $H \in \mathfrak{a}$, then $e^H \circ T = T \circ \text{Ad}(e^H)$.
- (3) $d(T)_0 : \mathfrak{n}^- \rightarrow T_{P^+} \mathcal{F}^+$ is a linear isomorphism.

Proof. By definition, F is transverse to P^- if and only if $(F, P^-) \in \mathbf{G} \cdot (P^+, P^-)$ if and only if $F = gP^+$ for some $g \in P^-$. By the Langlands decomposition, $P^- = N^-(P^+ \cap P^-)$ where $N^- \subset \mathbf{G}$ is the connected Lie subgroup with Lie algebra \mathfrak{n}^- . Since N^- is nilpotent, $N^- = e^{\mathfrak{n}^-}$. So F is transverse to P^- if and only if $F = e^X P^+$ for some $X \in \mathfrak{n}^-$. This proves part (1).

Part (2) is an immediate consequence of the definition. Part (3) follows from the fact that $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{p}^+$ (as vector spaces) and \mathfrak{p}^+ is the Lie algebra of P^+ . \square

As a consequence of (1) and (2) in Observation B.11, we have the following.

Lemma B.12. *If $\{H_n\}$ is a sequence in \mathfrak{a} with $\lim_{n \rightarrow \infty} \alpha(H_n) = -\infty$ for all $\alpha \in \Sigma$ with $\alpha(H_0) < 0$, then*

$$\lim_{n \rightarrow \infty} e^{H_n}(F) = P^+$$

for all $F \in \mathcal{F}^+$ transverse to P^- .

Proof. By Observation B.11 (1), $F = T(X)$ for some $X \in \mathfrak{n}^-$. Write $X = \sum_{\alpha(H_0) < 0} X_\alpha$, where $X_\alpha \in \mathfrak{g}_\alpha$. Then by Observation B.11 (2),

$$e^{H_n}(F) = T \left(\text{Ad}(e^{H_n}) \left(\sum_{\alpha(H_0) < 0} X_\alpha \right) \right) = T \left(\sum_{\alpha(H_0) < 0} e^{\text{ad}(H_n)} X_\alpha \right) = T \left(\sum_{\alpha(H_0) < 0} e^{\alpha(H_n)} X_\alpha \right).$$

Since $\lim_{n \rightarrow \infty} \alpha(H_n) = -\infty$ for all $\alpha \in \Sigma$ with $\alpha(H_0) < 0$, it follows that

$$\lim_{n \rightarrow \infty} e^{H_n}(F) = T(0) = P^+. \quad \square$$

Let $K \subset \mathbf{G}$ be the maximal compact subgroup with Lie algebra \mathfrak{k} and fix a K -invariant Riemannian metric on \mathcal{F}^+ , and let $|\cdot|$ denote the induced family of norms on the fibers of $T \mathcal{F}^+$.

Let $\xi = (\xi^+, \xi^-)$ be the P^\pm -Anosov limit map for ρ . Then let $\sigma_\xi^+(v) = (v, \xi^+(v^+))$. Since ρ is Anosov, there is a ρ -equivariant family of norms on the fibers of $\mathbf{U}(\Gamma) \times T \mathcal{F}^\pm \rightarrow \mathbf{U}(\Gamma) \times \mathcal{F}^\pm$ and constants $C, c > 0$ such that

$$\|Z\|_{\phi_{-t}(\sigma_\xi^+(v))} \leq C e^{-ct} \|Z\|_{\sigma_\xi^+(v)}$$

for all $t > 0$, $v \in \mathbf{U}(\Gamma)$ and $Z \in T_{\xi^+(v^+)} \mathcal{F}^+$.

Consider an escaping sequence $\{\gamma_n\}$ with $\gamma_n \rightarrow x$ and $\gamma_n^{-1} \rightarrow y$.

Case 1: If $x \neq y$, then γ_n is hyperbolic when n is sufficiently large, $\gamma_n^+ \rightarrow x$, and $\gamma_n^- \rightarrow y$. Furthermore, we can find a bounded sequence $\{v_n\}$ in $\mathbf{U}(\Gamma)$ such that $v_n^\pm = \gamma_n^\pm$, and a bounded sequence $\{g_n\}$ in \mathbf{G} such that

$$g_n(\xi^+(\gamma_n^+), \xi^-(\gamma_n^-)) = (P^+, P^-).$$

Then

$$g_n \rho(\gamma_n) g_n^{-1} P^\pm = P^\pm \quad \text{so} \quad g_n \rho(\gamma_n) g_n^{-1} \in L := P^+ \cap P^-.$$

Notice that

$$\mathfrak{g}_0 \oplus \bigoplus_{\alpha(H_0)=0} \mathfrak{g}_\alpha$$

is a root space decomposition of the Lie algebra of L . Then, using the Cartan decomposition of the reductive group L , there exist $k_{n,1}, k_{n,2} \in K \cap L$ and $H_n \in \mathfrak{a}$ so that

$$g_n \rho(\gamma_n) g_n^{-1} = k_{n,1} e^{H_n} k_{n,2}.$$

Since $\{v_n\}$ is a bounded sequence there exists $C_1 > 1$ such that

$$\frac{1}{C_1} |Z|_{\xi^+(v_n^+)} \leq \|Z\|_{\sigma_\xi^+(v_n)} \leq C_1 |Z|_{\xi^+(v_n^+)}$$

for all $n \geq 1$ and $Z \in T_{\xi^+(v_n^+)} \mathcal{F}^+$. Likewise, there exists $C_2 > 1$ such that

$$\frac{1}{C_2} |Z|_F \leq |g_n(Z)|_{g_n(F)} \leq C_2 |Z|_F$$

for all $n \geq 1$, $F \in \mathcal{F}^+$ and $Z \in T_F \mathcal{F}^+$.

Notice that $\gamma_n^{-1}(v_n) = \phi_{-t_n}(v_n)$ for some sequence $\{t_n\}$ with $t_n \rightarrow \infty$. Since both $k_{n,1}$ and $k_{n,2}$ fix P_+ and $|\cdot|$ is a K -invariant family of norms, it follows that for any $Z \in T_{P^+} \mathcal{F}^+$, we have

$$\begin{aligned} |e^{H_n}(Z)|_{P^+} &= |k_{n,1}^{-1} g_n \rho(\gamma_n) g_n^{-1} k_{n,2}^{-1}(Z)|_{P^+} \leq C_2 |\rho(\gamma_n) g_n^{-1} k_{n,2}^{-1}(Z)|_{\xi^+(v_n^+)} \\ &\leq C_1 C_2 \|\rho(\gamma_n) g_n^{-1} k_{n,2}^{-1}(Z)\|_{\sigma_\xi^+(v_n)} = C_1 C_2 \|g_n^{-1} k_{n,2}^{-1}(Z)\|_{\phi_{-t_n}(\sigma_\xi^+(v_n))} \\ (26) \quad &\leq C_1 C_2 C e^{-ct_n} \|g_n^{-1} k_{n,2}^{-1}(Z)\|_{\sigma_\xi^+(v_n)} \leq C_1^2 C_2 C e^{-ct_n} |g_n^{-1} k_{n,2}^{-1}(Z)|_{\xi^+(v_n^+)} \\ &\leq C_1^2 C_2^2 C e^{-ct_n} |Z|_{P^+}. \end{aligned}$$

By Observation B.11 (3), we know that for any $\alpha \in \Sigma$ such that $\alpha(H_0) < 0$, and any $X \in \mathfrak{g}_\alpha$, there is some $Z \in T_{P^+} \mathcal{F}^+$ such that $d(T)_0(X) = Z$. Then

$$e^{H_n}(Z) = d(e^{H_n} \circ T)_0(X) = \left. \frac{d}{dt} \right|_{t=0} e^{H_n} \circ T(tX).$$

Then by Observation B.11 (2) (see proof of Lemma B.12),

$$\left. \frac{d}{dt} \right|_{t=0} e^{H_n} \circ T(tX) = \left. \frac{d}{dt} \right|_{t=0} T(te^{\alpha(H_n)} X) = e^{\alpha(H_n)} \left. \frac{d}{dt} \right|_{t=0} T(tX) = e^{\alpha(H_n)} Z.$$

Thus, $e^{H_n}(Z) = e^{\alpha(H_n)} Z$, so the inequality (26) implies that

$$\lim_{n \rightarrow \infty} \alpha(H_n) = -\infty$$

whenever $\alpha(H_0) < 0$. Hence, by Lemma B.12,

$$\lim_{n \rightarrow \infty} e^{H_n}(F) = P^+$$

for all $F \in \mathcal{F}^+$ transverse to P^- . Since $g_n(\xi^+(x), \xi^-(y)) \rightarrow (P^+, P^-)$, $k_{n,j} P^\pm = P^\pm$ and $\rho(\gamma_n) = g_n^{-1} k_{n,1} e^{H_n} k_{n,2} g_n$ we then have

$$\lim_{n \rightarrow \infty} \rho(\gamma_n)(F) = \xi^+(x)$$

for all $F \in G/P^+$ transverse to $\xi^-(y)$.

Case 2: If $x = y$, pick $\beta \in \Gamma$ so that $z := \beta^{-1}(x) \neq x$. Then $\gamma_n \beta \rightarrow x$ and $(\gamma_n \beta)^{-1} \rightarrow z \neq x$. By the first case, $\rho(\gamma_n \beta)(F) \rightarrow \xi(x)$ for all $F \in \mathcal{F}^+$ transverse to $\xi^-(z) = \rho(\beta^{-1})\xi^-(x)$. Equivalently, $\rho(\gamma_n)(F) \rightarrow \xi^+(x)$ for all $F \in \mathcal{F}^+$ transverse to $\xi^-(x)$. \square

REFERENCES

- [1] H. Abels, G. Margulis and G. Soifer, “Semigroups containing proximal linear maps,” *Israel J. Math.* **9**(1995), 1–30.
- [2] J. Bochi, R. Potrie and A. Sambarino, “Anosov representations and dominated splittings,” *J.E.M.S.* **21**(2019), 3343–3414.
- [3] B. Bowditch, “Spaces of geometrically finite representations,” *Ann. Acad. Sci. Fenn.* **23**(1998), 389–414.
- [4] C. Brav and H. Thomas, “Thin monodromy in $\mathrm{Sp}(4, \mathbb{R})$,” *Compos. Math.* **150**(2014), 333–343.
- [5] H. Bray, R. Canary, L.-Y. Kao, and G. Martone, “Counting, equidistribution and entropy gaps at infinity with applications to cusped Hitchin representations,” preprint, arXiv:2102.08552.
- [6] M. Bridgeman, R. Canary, and F. Labourie, “Simple length rigidity for Hitchin representations,” *Adv. Math.* **360**(2020).
- [7] M. Bridgeman, R. Canary, F. Labourie and A. Sambarino, “The pressure metric for Anosov representations,” *G.A.F.A.* **25**(2015), 1089–1179.
- [8] M. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, 1999.
- [9] J.P. Burelle and N. Treib, “Schottky presentations of positive representations,” preprint, arXiv:1807.05286.
- [10] C.I. Byrnes and M.A. Gauger, “Decidability criteria for the similarity problem, with applications to the moduli of linear dynamical systems,” *Adv. Math.* **25**(1977), 59–90.
- [11] M. Crampon and L. Marquis “Finitude géométrique en géométrie de Hilbert,” *Ann. Inst. Fourier (Grenoble)* **64** (2014), 2299–2377.
- [12] S. Filip, “Uniformization of some weight 3 variations of Hodge structure, Anosov representations, and Lyapunov exponents,” preprint, arXiv:2110.07533.
- [13] S. Filip and C. Fougerson, “A cyclotomic family of thin hypergeometric monodromy groups in $\mathrm{Sp}(4, \mathbb{R})$,” preprint, arXiv:2106.09181.
- [14] V. Fock and A. Goncharov, “Moduli spaces of local systems and higher Teichmüller theory,” *Publ. Math. I.H.E.S.* **103**(2006), 1–211.
- [15] F. Guéritaud, O. Guichard, F. Kassel and A. Wienhard, “Anosov representations and proper actions,” *Geom. Top.* **21**(2017), 485–584.
- [16] O. Guichard and A. Wienhard, “Anosov representations: Domains of discontinuity and applications,” *Invent. Math.* **190**(2012), 357–438.
- [17] N. Hitchin, “Lie groups and Teichmüller space,” *Topology* **31**(1992), 449–473.
- [18] M. Kapovich and B. Leeb, “Relativizing characterizations of Anosov subgroups, I,” preprint, arXiv:1807.00160.
- [19] M. Kapovich, B. Leeb and J. Porti, “Anosov subgroups: Dynamical and geometric characterizations,” *Eur. Math. J.* **3**(2017), 808–898.
- [20] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, 1995.
- [21] S. Kim, S.P. Tan and T. Zhang, “Weakly positive representations,” preprint, arXiv:2008.02424.
- [22] F. Labourie, “Anosov flows, surface groups and curves in projective space,” *Invent. Math.* **165**(2006), 51–114.
- [23] F. Labourie and G. McShane, “Cross ratios and identities for higher Thurston theory,” *Duke Math. J.* **148**(2009), 279–345.
- [24] J. Loftin and T. Zhang, “Coordinates on the augmented moduli space of convex $\mathbb{R}\mathbb{P}^2$ structures,” *Jour. L.M.S.* **104**(2021), 1930–1972.
- [25] G. Lusztig, “Total positivity in reductive groups,” in *Lie theory and geometry*, Birkhäuser, 1994, 531–568.
- [26] L. Marquis, “Espaces des modules des surfaces projectives proprement convexes de volume fini,” *Geom. Top.* **14**(2010), 2103–2149.
- [27] L. Marquis, “Surface projective proprement convexe de volume fini,” *Ann. Inst. Four.* **62**(2012), 325–392.
- [28] H. Masur, “Extension of the Weil-Petersson metric to the boundary of Teichmüller space,” *Duke Math. J.* **43**(1976), 623–635.
- [29] G.D. Mostow, *Strong rigidity of locally symmetric spaces*, Princeton University Press, 1973.
- [30] A. Sambarino, “Infinitesimal Zariski closures of positive representations,” preprint, arXiv:2012.10276.
- [31] M. Shub, *Global Stability of Dynamical Systems*, Springer-Verlag, 1987.
- [32] Z. Sun, A. Wienhard and T. Zhang, “Flows on the $\mathrm{PGL}(V)$ -Hitchin component,” *G.A.F.A.* **30**(2020), 588–692.
- [33] K. Tsouvalas, “Anosov representations, strongly convex cocompact groups and weak eigenvalue gaps,” preprint, arXiv:2008.04462.
- [34] F. Zhu, “Relatively dominated representations,” *Ann. Inst. Fourier*, to appear, arxiv:1912.13152.
- [35] F. Zhu, “Relatively dominated representations from eigenvalue gaps and limit maps,” preprint, arXiv:2102.10611.
- [36] T. Zhang and A. Zimmer, “Regularity of limit sets of Anosov representations,” preprint, arXiv:1903.11021.

UNIVERSITY OF MICHIGAN

NATIONAL UNIVERSITY OF SINGAPORE

UNIVERSITY OF WISCONSIN-MADISON