

Igor V. Dolgachev*

Quartic surfaces with icosahedral symmetry

DOI 10.1515/advgeom-2017-0040. Received 23 September, 2016; revised 9 April, 2017

Abstract: We study smooth quartic surfaces in \mathbb{P}^3 which admit a group of projective automorphisms isomorphic to the icosahedron group.

Keywords: Quartic surface, icosahedron group.

2010 Mathematics Subject Classification: 14J10

Communicated by: I. Coskun

1 Introduction

Let \mathfrak{A}_5 be the icosahedron group isomorphic to the alternating group in 5 letters. Starting from Platonic solids, it appears as an omnipresent symmetry group in geometry. In this article, complementing papers [4] and [5], we discuss families of smooth quartic surfaces in \mathbb{P}^3 that admit the group \mathfrak{A}_5 as its group of projective symmetries.

It follows from loc. cit. that any smooth quartic surface S with a faithful action of \mathfrak{A}_5 belongs to one of the two pencils of invariant quartic surfaces. One of them arises from a linear irreducible 4-dimensional representation of \mathfrak{A}_5 and was studied in great detail by K. Hashimoto in [10]. It contains a double quadric and four surfaces with 5, 10, 10, or 15 ordinary double points. The other pencil arises from a faithful linear representation of the binary icosahedron group $2.\mathfrak{A}_5$. As was shown in [4] and [5], it contains two singular surfaces singular along its own rational normal cubic and two surfaces with 10 ordinary double points. In this paper we show first that one of the surfaces with 10 ordinary double points from Hashimoto's pencil can be realized as a Cayley quartic symmetroid defined by a \mathfrak{A}_5 -invariant web W of quadrics (this result was independently obtained by S. Mukai). We also show that the Steinerian surface of this web parametrizing singular points of singular quadrics in the web coincides with one of two smooth members of the second pencil that admits a larger group of projective symmetries isomorphic to \mathfrak{S}_5 . To see this we give an explicit equation of the second pencil. We also show that the apolar linear system of quadrics to the web W contains two invariant rational normal curves that give rise to two rational plane sextics with symmetry \mathfrak{A}_5 discovered by R. Winger [20], [21].

It is my pleasure to thank I. Cheltsov, B. van Geemen, K. Hulek and S. Mukai for their help in collecting the known information about the subject of this paper. I would like also to thank the referee for careful reading of the manuscript and useful remarks.

2 Two irreducible 4-dimensional representations

Let E be a 4-dimensional linear space over an algebraically closed field \mathbb{k} of characteristic $\neq 2, 3, 5$ and let $|E| \cong \mathbb{P}^3$ be the projective space of lines in E . Assume that E is a non-trivial projective linear representation space for $\mathfrak{A}_5 \cong \text{PSL}(2, \mathbb{F}_5)$ in $|E|$. Then it originates either from a linear representation of \mathfrak{A}_5 in E , or from a linear representation of its central double extension $2.\mathfrak{A}_5 \cong \text{SL}(2, \mathbb{F}_5)$, the *binary icosahedral group*. Thus E is either an irreducible representation, or contains a one-dimensional trivial representation, or decomposes into the sum of two irreducible two-dimensional representations.

*Corresponding author: Igor V. Dolgachev, Department of Mathematics, University of Michigan, 525 E. University Av., Ann Arbor, MI, 49109, USA, email: idolga@umich.edu

Proposition 1. *Suppose there exists a smooth quartic surface S in $|E|$ which is invariant with respect to a non-trivial projective representation of \mathfrak{A}_5 . Then E is an irreducible representation of $G = \mathfrak{A}_5$ or $G = 2 \cdot \mathfrak{A}_5$.*

Proof. Suppose E has a one-dimensional trivial summand. It is known that an element of order 3 acting on a smooth quartic surface has 6 isolated fixed points; see [18]. A glance at the character tables of the groups shows that an element g of order 3 in G in its action on the 3-dimensional summand has three different eigenvalues, one of them is equal to 1. This shows that g has a pointwise fixed line in $|E|$ and two isolated fixed points. The line intersects the quartic at 4 points, so all the fixed points are accounted for. In particular, one of them is fixed with respect to the whole group \mathfrak{A}_5 . However, the assumption on the characteristic implies that G acts faithfully on the tangent space of S at this point, and the group \mathfrak{A}_5 has no non-trivial 2-dimensional linear representations.

Now we consider case when E decomposes into the sum of two irreducible 2-dimensional representations. This could happen only when $G = 2 \cdot \mathfrak{A}_5$. In this case G has two invariant lines in $|E|$, hence the union of two invariant sets of at most 4-points on S . The known possible orders of subgroups of \mathfrak{A}_5 shows that this is impossible. \square

So, we are dealing with projective representations of the group \mathfrak{A}_5 coming from an irreducible 4-dimensional representation W_4 of \mathfrak{A}_5 or from an irreducible faithful 4-dimensional linear representation U_4 of $2 \cdot \mathfrak{A}_5$.

It is known that the group $2 \cdot \mathfrak{A}_5$ has two irreducible 2-dimensional representations V and V' , both self-dual but one is obtained from the other by composition with an outer automorphism of \mathfrak{A}_5 . The center acts as the minus identity. Since it acts as the identity on the symmetric squares S^2V or S^2V' , the groups \mathfrak{A}_5 , $2 \cdot \mathfrak{A}_5$ admit two 3-dimensional representations, both of them are irreducible and self-dual, they differ from each other by an outer automorphism.

The group \mathfrak{A}_5 acts in $|S^2V^\vee|$ via the Veronese map of $|V| \rightarrow |S^2V|$ of its natural action on $|V| \cong \mathbb{P}^1$. The restriction of the representation S^2V of \mathfrak{A}_5 to its subgroup H of order 10 isomorphic to the dihedral group D_{10} has a trivial one-dimensional summand. It defines a fixed point of H in its action in the plane $|S^2V|$. Its orbit of \mathfrak{A}_5 consists of six points, called by F. Klein the *fundamental points*. The linear system of plane cubics is invariant with respect to \mathfrak{A}_5 and defines an irreducible subrepresentation of $S^3(S^2V)^\vee$ of dimension 4. The image of the plane under the map defined by the linear system of cubics is isomorphic to the *Clebsch diagonal cubic surface* representing the unique isomorphism class of a nonsingular cubic surface with the automorphism group isomorphic to \mathfrak{S}_5 ; see [9], 9.5.4. The representation W_4 is isomorphic to the restriction of the *standard irreducible representation* of \mathfrak{S}_5 realized in the hyperplane $x_1 + \dots + x_5 = 0$ in k^5 .

The representation U_4 is realized in the third symmetric power S^3V of V . It is self-dual and isomorphic to S^3V' . The projective representation of \mathfrak{A}_4 in $|U_4|$ is obtained via the Veronese map $|V| \rightarrow |S^3V|$ from the natural action of \mathfrak{A}_5 on $|V|$.

Using the character table for G one obtains the following.

Proposition 2. *Let S^4W^\vee and S^4U_4 be the fourth symmetric powers of W_4 and of U_4 , and let $(\)^G$ denote the subspace of G -invariant elements. Then*

$$\dim(S^4W^\vee)^G = \dim(S^4U_4)^G = 2.$$

Thus we have two pencils of invariant quartic polynomials $(S^4W_4)^{\mathfrak{A}_5}$ and $(S^4U_4)^{2 \cdot \mathfrak{A}_5}$, so our quartic surface S is a member of one of them.

3 The pencil $|(S^4W_4^\vee)^{\mathfrak{A}_5}|$

Since W_4 is the restriction of the standard representation of \mathfrak{S}_5 , the invariant theory says that the space $(S^4W_4)^{\mathfrak{A}_5}$ is isomorphic to the linear space of symmetric polynomials of degree 4 in variables x_1, \dots, x_5 and the discriminant anti-symmetric polynomial. Since the degree of the latter is larger than 4, we obtain that

$$(S^4W_4)^{\mathfrak{A}_5} = (S^4W_4)^{\mathfrak{S}_5}.$$

Thus any \mathfrak{A}_5 -invariant pencil of quartic surfaces in $|W_4|$ consists of the quartics given by equations $s_4 + \lambda s_2^2 = s_1 = 0$, where the s_i are elementary symmetric functions. It will be more convenient to rewrite this equation in terms of power symmetric functions:

$$x_1^4 + \dots + x_5^4 - t(x_1^2 + \dots + x_5^2)^2 = x_1 + \dots + x_5 = 0 \tag{3.1}$$

where $t = (2\lambda + 1)/2$.

Computing the partial derivatives, we find that the \mathfrak{S}_5 -orbits of singular points are represented by points with coordinates $(1, 1, -1, -1, 0)$, or $(1, -1, 0, 0, 0)$, or $(2, 2, 2, -3, -3)$, or $(1, 1, 1, 1, -4)$ corresponding to the parameters $t = 1/4, 1/2, 7/30, 13/20$, respectively. This gives the following.

Proposition 3. *The surface S_t given by equation (3.1) is nonsingular if and only if $t \neq \frac{1}{2}, \frac{1}{4}, \frac{7}{30}, \frac{13}{20}$. If $t = 1/4$, it has 15 singular points, if $t = 1/2, 7/30$, it has 10 singular points, if $t = 13/20$, it has 5 singular points. Each singular point is an ordinary node.*

Example 1. The surface $S_{1/4}$ is the intersection of the *Castelnuovo–Richmond–Igusa quartic threefold* (see [9], 9.4.4) given by the equations

$$x_1^4 + \dots + x_6^4 - \frac{1}{4}(x_1^2 + \dots + x_6^2)^2 = x_1 + \dots + x_6 = 0 \tag{3.2}$$

in \mathbb{P}^5 and the hyperplane $x_6 = 0$. Its singular points are the intersection of the hyperplane with the fifteen double lines of the threefold.

Example 2. If $t = 1/2$, then $\lambda = 0$ and the surface can be rewritten by the equations $\sum_{i=1}^5 \frac{1}{x_i} = x_1 + \dots + x_5 = 0$ in \mathbb{P}^4 . We recognize the equation of the Hessian surface of the Clebsch diagonal cubic surface C_3 . Its 10 nodes are the vertices of the Sylvester pentahedron. Its edges lie on the surface and together with the nodes form the Desargues symmetric configuration (10_3) ; see [11], III §19. By definition of the Hessian surface, $S_{1/2}$ is a *quartic symmetroid* (see [9], 4.2.6), the locus of points in $|E|$ such that the polar quadric of the cubic surface C_3 is singular. This symmetric determinantal representation of $S_{1/2}$ is defined by a linear map of \mathfrak{S}_5 -representations $W_4 \rightarrow S^2W_4^\vee$. It is the polar map associated with the cubic surface (see [9]). It allows one to rewrite the equation of the Hessian of the Clebsch cubic surface as the symmetric determinant:

$$\det \begin{pmatrix} L - x_1 & L & L & L \\ L & L - x_2 & L & L \\ L & L & L - x_3 & L \\ L & L & L & L - x_4 \end{pmatrix} = 0,$$

where $L = x_1 + x_2 + x_3 + x_4$.

It is known that the surface $S_{1/2}$ is isomorphic to the K3-cover of an Enriques surface X with $\text{Aut}(X) \cong \mathfrak{S}_5$ (of type VI in Kondo’s classification [16]). The covering involution is defined by the standard Cremona transformation $(x_1, \dots, x_5) \mapsto (x_1^{-1}, \dots, x_5^{-1})$.

Example 3. Assume that $t = 7/30$. Projecting from the node $q_0 = (2, 2, 2, -3, -3)$, we get the equation of $S_{7/30}$ as the double plane

$$\begin{aligned} w^2 &= (xy + xz + yz)^2(x + y + z)^2 - 3(x - y)^2(x - z)^2(y - z)^2 \\ &= 111x^2y^2z^2 + 80(x^3y^2z + x^3yz^2 + x^2y^3z + x^2yz^3 + xy^3z^2 + xy^2z^3) \\ &\quad + 13(x^4y^2 + x^4z^2 + y^4z^2 + x^2z^4 + y^2z^4 + x^2y^4) + 10(x^4yz + x^3y^3 + x^3z^3 + xy^4z + xyz^4 + y^3z^3). \end{aligned}$$


The branch curve B is the union of two cubic curves intersecting at 9 points

$$[1, 0, 0], [0, 1, 0], [0, 0, 1], [2, -1, 2], [-1, 2, 2], [2, 2, -1], [-2, 1, 1], [1, -2, 1], [1, 1, -2].$$

It is well-known and goes back to A. Cayley (see a modern exposition in [8]) that this implies that the surface $S_{7/30}$ is a quartic symmetroid. We will return to this example in the next section.

Example 4. Let \mathcal{S}_3 be the Segre cubic primal (see [9], 9.4.4) given by the equations $x_1^3 + \dots + x_6^3 = x_1 + \dots + x_6 = 0$. It is isomorphic to the image of a rational map $f : \mathbb{P}^3 \dashrightarrow \mathbb{P}^4$ given by the linear system of quadrics through 5 points p_1, \dots, p_5 in general linear position. The images of the lines $\langle p_i, p_j \rangle$ are the 10 nodes on \mathcal{S}_3 . Consider the surface S in \mathcal{S}_3 given by the additional equation $x_1^2 + \dots + x_6^2 = 0$. Obviously, it has \mathfrak{S}_6 -symmetry. The pre-image of S under the map f is a quartic surface in \mathbb{P}^3 with 5 nodes at p_1, \dots, p_5 . One can make the map f to be \mathfrak{S}_5 -invariant by viewing \mathbb{P}^3 as the hyperplane $y_0 + \dots + y_5 = 0$ in \mathbb{P}^4 and choosing the points p_1, \dots, p_5 to be the points in the hyperplane with coordinates $[1, 0, 0, 0, -1], [0, 1, 0, 0, -1],$ etc. The group \mathfrak{S}_5 acts naturally in \mathbb{P}^3 by permuting the 5 points. The restriction of this representation to \mathfrak{A}_5 is the projectivization of the linear representation space W_4 . Via this action, the map f becomes a \mathfrak{S}_5 -invariant birational map from \mathbb{P}^3 to the Segre cubic \mathcal{C}_3 . Thus $f^{-1}(S)$ is a 5-nodal quartic in $|W_4|$ with \mathfrak{S}_5 -symmetry.

Note that the action of \mathfrak{S}_5 on the Segre cubic primal is closely related to its known action on the del Pezzo surface of degree 5 via the following commutative diagram explained in [19], Proposition 4.7:



Here χ is a flop, φ and φ_0^+ are small contractions to the Segre cubic primal, \mathcal{D} is a del Pezzo surface of degree 5 and φ is a \mathbb{P}^1 -bundle.

The pencil (3.1) was intensively studied by K. Hashimoto in [10]. So it is appropriate to refer to it as the Hashimoto pencil. In particular, Hashimoto computed the lattice $T(X_t)$ of transcendental cycles of a minimal nonsingular model X_t of a singular member S_t of the pencil.

Theorem 1. Assume that $\mathbb{k} = \mathbb{C}$. For any singular member of the Hashimoto pencil, the lattice $T(X_t)$ is of rank 2 and is given by the matrix

$$\begin{pmatrix} 4 & 0 \\ 0 & 10 \end{pmatrix} \text{ for } t = \frac{1}{2}, \quad \begin{pmatrix} 4 & 1 \\ 1 & 4 \end{pmatrix} \text{ for } t = \frac{1}{4}, \quad \begin{pmatrix} 4 & 2 \\ 2 & 16 \end{pmatrix} \text{ for } t = \frac{7}{30}, \quad \begin{pmatrix} 6 & 0 \\ 0 & 20 \end{pmatrix} \text{ for } t = \frac{13}{20}.$$

The transcendental lattice of a generic member of the pencil is of rank 3 and is given by the matrix

$$\begin{pmatrix} 4 & 1 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & -20 \end{pmatrix}.$$

Remark 1. It is known that the quartic surface defined by the equation

$$F = x^4 + y^4 + z^4 + w^4 + 12xyzw = 0 \tag{3.3}$$

admits a group of projective automorphisms isomorphic to $2^4 \cdot \mathfrak{S}_5$ (see [17]). According to S. Mukai [18], the subgroup $2^4 \cdot \mathfrak{A}_5$ is isomorphic to the Mathieu group M_{20} and is realized as one of the maximal finite groups of symplectic automorphisms of a complex K3 surface. The equation defining the surface is an invariant of the Heisenberg group \mathcal{H}_2 acting in its Schrödinger 4-dimensional irreducible representation (see [3]). The linear space of invariant quartic polynomials is 5-dimensional and it has a basis that consists of the polynomials

$$p_0 = x^4 + y^4 + z^4 + w^4, \quad p_1 = x^2w^2 + y^2z^2, \quad p_2 = x^2z^2 + y^2w^2, \quad p_3 = x^2y^2 + z^2w^2, \quad p_4 = xyzw.$$

Let

$$\begin{aligned} M_1 &= -2p_0 + 24p_4, \\ M_2 &= p_0 - 6(p_1 + p_2 + p_3), \\ M_3 &= p_0 + 6(-p_1 + p_2 + p_3), \\ M_4 &= p_0 + 6(p_1 - p_2 + p_3), \\ M_5 &= -2p_0 - 24p_4, \\ M_6 &= p_0 + 6(p_1 + p_2 - p_3) \end{aligned}$$

be a spanning set of this linear space. The fifth of them defines the surface from (3.3). We propose to call the polynomials M_i the *Maschke quartic polynomials* (not to be confused with the Maschke octic polynomial from [2]). It is shown in [17], p. 505, that they satisfy the equations

$$\left(\sum_{i=1}^6 M_i^2\right)^2 - 4 \sum_{i=1}^6 M_i^4 = \sum_{i=1}^6 M_i = 0.$$

We recognize the equations of the Castelnuovo–Richmond–Igusa quartic threefold from Example 1. Thus for any Maschke polynomial M_i , the surface $V(M_i)$ is a Galois 2^4 -cover of the pre-image of a coordinate hyperplane section of the quartic threefold isomorphic to the surface $S_{1/4}$ from the Hashimoto pencil. This shows the appearance of $2^4 \cdot \mathfrak{S}_5$ in the group of projective automorphisms of $V(M_i)$.

It was communicated to me by Bert van Geemen that the projective transformations defined by the matrices

$$M = \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \\ -1 & 1 & -1 & 1 \\ -1 & -1 & -1 & -1 \end{pmatrix} \quad \text{and} \quad N = \begin{pmatrix} 1 & -1 & -i & -i \\ -1 & 1 & -i & -i \\ i & i & 1 & -1 \\ i & i & -1 & 1 \end{pmatrix} \tag{3.4}$$

define automorphisms of orders 5 and 2 that generate a subgroup of automorphisms of $V(F)$ that splits the extension $2^4 \cdot \mathfrak{S}_5$. This shows that the Maschke quartic surface $V(M_i)$ admits \mathfrak{S}_5 and hence \mathfrak{A}_5 as groups of projective automorphisms. By computing the traces of M and N we find that \mathfrak{S}_5 originates from its linear standard irreducible 4-dimensional representation, and the surface must be isomorphic to a member of the Hashimoto pencil. According to computations made by Bert van Geemen, the surface $V(M_1)$ corresponds to the parameter $t = \frac{3}{20}(3-i)$. Note that according to S. Mukai the transcendental lattice of the Maschke surface is given by the diagonal 2×2 -matrix with the diagonal entries 4 and 40. So it is different from the transcendental lattice of a general member of the pencil.

4 The pencil $|(S^4 \mathbb{U}_4^\vee)^{\mathfrak{A}_5}|$

Recall that the linear representation space \mathbb{U}_4 of $G = 2 \cdot \mathfrak{A}_5$ is isomorphic to the space $S^3 \mathbb{V}$. Since there is only one isomorphism class of irreducible faithful 4-dimensional representations of G , we have an isomorphism $\mathbb{U}_4 \cong S^3 \mathbb{V} \cong S^3 \mathbb{V}'$.

Let e_1, e_2 be a basis in \mathbb{V} and let (u, v) be the coordinates in \mathbb{V} with respect to this basis, i.e. the dual basis of (e_1, e_2) in \mathbb{V}^\vee . The spaces $S^d \mathbb{V}$ and $S^d \mathbb{V}^\vee$ have natural monomial bases $(e_1^d, e_1^{d-1} e_2, \dots, e_2^d)$ and $(u^d, u^{d-1} v, \dots, v^d)$, respectively. The polarization isomorphism

$$S^d \mathbb{V}^\vee \rightarrow (S^d \mathbb{V})^\vee$$

assigns to $u^{d-i} v^i$ the linear function on $S^d \mathbb{V}$ that takes the value $\frac{1}{d!} (d-i)! i!$ on $e_1^{d-i} e_2^i$ and zero on all other monomials. This shows that the basis $((\binom{d}{i}) u^{d-i} v^i)_{i=0, \dots, d}$ is the dual basis of $(e_1^d, e_1^{d-1} e_2, \dots, e_2^d)$. Thus any binary form $f \in S^d \mathbb{V}^\vee$ can be written as

$$f = \sum_{i=0}^d \binom{d}{i} a_i u^{d-i} v^i, \tag{4.1}$$

so that (a_0, \dots, a_d) are the natural coordinates in the space $S^d\mathbb{U}^\vee$. Although this notation is widely used in the invariant theory, we switch to the basis $((\binom{d}{i}e_1^{d-i}e_2^i)$ of $S^d\mathbb{V}$ to get the dual basis formed by the monomials $u^{d-i}v^i$. So we drop the binomial coefficients in (4.1). This helps us to obtain formulas that agree with those given in Klein's book.

Let us clarify the coordinate-free definition of the Veronese map

$$v_d : \mathbb{V} \rightarrow S^d\mathbb{V}.$$

It is defined by assigning to a vector $\alpha e_1 + \beta e_2$ the linear function $f \mapsto f(\alpha, \beta)$ on $S^d\mathbb{V}^\vee = (S^d\mathbb{V})^\vee$. It follows that

$$v_d(\alpha e_1 + \beta e_2) = \sum_{i=0}^d \binom{d}{i} \alpha^{d-i} \beta^i e_1^{d-i} e_2^i = (\alpha e_1 + \beta e_2)^d.$$

In coordinates, this is the map

$$(u, v) \mapsto (u^d, u^{d-1}v, \dots, uv^{d-1}, v^d). \tag{4.2}$$

Passing to the projective space, we get the map

$$v_d : |\mathbb{V}| \rightarrow |S^d\mathbb{V}|$$

that is given by the complete linear system $|S^d\mathbb{V}^\vee| = |\mathcal{O}_{|\mathbb{V}|}(d)|$. The image R_d of this map is a *Veronese curve* of degree d , or a *rational normal curve* of degree d . If we re-denote the coordinates $u^{d-i}v^i$ by (x_0, \dots, x_d) , a hyperplane $V(\sum_{i=0}^d a_i x_i)$ intersects R_d along the closed subscheme that is isomorphic, under the Veronese map v_d , to the closed subscheme $V(\sum_{i=0}^d a_i u^{d-i}v^i)$ of $|\mathbb{V}|$.

Dually, we have the Veronese map

$$v_d^* : \mathbb{V}^\vee \rightarrow S^d\mathbb{V}^\vee$$

which assigns to a linear function $l = au + bv \in \mathbb{V}^\vee$ the linear function $v_d(l) \in S^d\mathbb{V}^\vee$ that takes on $e_1^{d-i}e_2^i$ the value equal to $a^{d-i}b^i$. It follows that

$$v_d^*(\alpha u + \beta v) = \sum_{i=0}^d \binom{d}{i} \alpha^{d-i} \beta^i u^{d-i} v^i = (\alpha u + \beta v)^d.$$

So we get a familiar picture: points of $|S^d\mathbb{V}^\vee|$ are non-zero binary forms of degree d up to proportionality, and points of the Veronese curve are powers of linear forms, up to proportionality.

In coordinates, the dual Veronese map is given by

$$(e_1, e_2) \mapsto (e_1^d, \dots, \binom{d}{i} e_1^{d-i} e_2^i, \dots, e_2^d).$$

The image R_d^* of the corresponding map

$$v_d^* : |\mathbb{V}^\vee| = \mathbb{P}(\mathbb{V}) \rightarrow |S^d\mathbb{V}^\vee| = \mathbb{P}(S^d\mathbb{V})$$

is the *dual Veronese curve* of degree d . The duality can be clarified more explicitly. For any point $x \in R_d$, one can consider the *osculating hyperplane* at x , the unique hyperplane in $|S^d\mathbb{V}|$ that intersects R_d at one point x with multiplicity d . The dual Veronese curve R_d^* in the dual space $|S^d\mathbb{V}^\vee|$ is the locus of osculating hyperplanes.

One can use the isomorphism $|\mathbb{V}^\vee| \rightarrow |\mathbb{V}|$ defined by assigning to a linear function $l = au + bv \in \mathbb{V}^\vee$ its zero $V(l) = [-b, a] \in |\mathbb{V}|$. In other terms, it is defined by the exterior product pairing $\mathbb{V} \times \mathbb{V} \rightarrow \wedge^2 \mathbb{V} \cong \mathbb{k}$. Thus we have two Veronese maps

$$v_d : |\mathbb{V}| \rightarrow |S^d\mathbb{V}| \quad \text{and} \quad v_d^* : |\mathbb{V}| \rightarrow |S^d\mathbb{V}^\vee|$$

with the images R_d and R_d^* . The map of $|\mathbb{V}| \rightarrow |S^d\mathbb{V}^\vee|$ is given, in coordinates, by

$$(u, v) \rightarrow ((-1)^d e_2^d, \dots, (-1)^{d-i} \binom{d}{i} e_1^i e_2^{d-i}, \dots, e_1^d).$$

Let $\rho : G \rightarrow GL(\mathbb{V})$ be a linear representation of a group G . By functoriality, it defines a linear representation $S^d(\rho) : G \rightarrow GL(S^d\mathbb{V})$. The dual linear representation $\rho^* : G \rightarrow GL(\mathbb{V}^\vee)$ defines a representation $S^d(\rho^*) : G \rightarrow GL(S^d\mathbb{V}^\vee)$. It follows from the polarization isomorphism that the representations $S^d(\rho)$ and $S^d(\rho^*)$ are dual to each other.

After fixing these generalities, it is easy to describe irreducible linear representations of $G = 2.\mathfrak{A}_5$. We start with the 2-dimensional representations \mathbb{V} and \mathbb{V}^\vee . We choose a basis e_1, e_2 in \mathbb{V} and its dual basis (u, v) in \mathbb{V}^\vee as above to assume that the group preserves the volume forms $e_1 \wedge e_2$ and $u \wedge v$. Thus in these bases we represent the matrices of the representation by unimodular matrices. According to [15], p. 213, the group $2.\mathfrak{A}_5$ is generated by the transformations S, T, U of orders 5, 4, 4, respectively, and its representation in \mathbb{V} is given in terms of coordinates as follows:

$$\begin{aligned} S : (u, v) &\mapsto (\varepsilon^3 u, \varepsilon^2 v), \\ T : (u, v) &\mapsto \frac{1}{\sqrt{5}}(-cu + dv, du + cv), \\ U : (u, v) &\mapsto (-v, u), \end{aligned}$$

where $\varepsilon = e^{2\pi i/5}$ and $c = \varepsilon - \varepsilon^{-1}, d = \varepsilon^2 - \varepsilon^{-2}$. We have

$$\lambda = \frac{c}{d} = \varepsilon + \varepsilon^{-1} + 1 = \frac{1 + \sqrt{5}}{2},$$

which is the golden ratio. It satisfies $\lambda^2 = \lambda + 1$. Note that the trace of S is equal to $\varepsilon^3 + \varepsilon^2 = \lambda^2 - 2 = \lambda - 1 = \frac{-1 + \sqrt{5}}{2}$. This is denoted by b_5 in [7], so we can identify this representation with the one given by the character χ_6 . The representation \mathbb{V}' is given by the same formulas as above, where ε is replaced with ε^2 . The trace of S becomes $b_5^* := -\lambda$.

Next we consider the 3-dimensional irreducible representations realized in $S^2\mathbb{V}$ and $S^3\mathbb{V}'$. In the basis $(e_1^2, 2e_1e_2, e_2^2)$ of \mathbb{V} , the first representation is given by the matrices

$$S : \begin{pmatrix} \varepsilon & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \varepsilon^{-1} \end{pmatrix}, \quad T : \frac{1}{\sqrt{5}} \begin{pmatrix} \varepsilon + \varepsilon^4 & 2 & \varepsilon^2 + \varepsilon^3 \\ 1 & 1 & 1 \\ \varepsilon^2 + \varepsilon^3 & 2 & \varepsilon + \varepsilon^4 \end{pmatrix}, \quad U : \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Note that Klein uses slightly different coordinates $(A_0, A_1, A_2) = (-uv, v^2, -u^2)$, so his matrices are slightly different.

The second irreducible 3-dimensional representation $(S^2\mathbb{V})'$ is obtained by replacing ε with ε^2 . The invariant theory for the icosahedron group \mathfrak{A}_5 in this representation is discussed in Klein’s book (see also [9], 9.5.4).

The linear representation \mathbb{U}_4 of $2.\mathfrak{A}_5$ realized in $S^3\mathbb{V}$ is given by the matrices

$$\begin{aligned} S &: \begin{pmatrix} \varepsilon^4 & 0 & 0 & 0 \\ 0 & \varepsilon^3 & 0 & 0 \\ 0 & 0 & \varepsilon^2 & 0 \\ 0 & 0 & 0 & \varepsilon \end{pmatrix}, \\ T &: \frac{d^3}{5\sqrt{5}} \begin{pmatrix} -\lambda^3 & 3\lambda^2 & -3\lambda & 1 \\ \lambda^2 & -2\lambda + \lambda^3 & 1 - 2\lambda^2 & \lambda \\ -\lambda & 1 - 2\lambda^2 & 2\lambda - \lambda^3 & \lambda^2 \\ 1 & 3\lambda & 3\lambda^2 & \lambda^3 \end{pmatrix} = \frac{d^3}{5\sqrt{5}} \begin{pmatrix} -(1 + 2\lambda) & 3(\lambda + 1) & -3\lambda & 1 \\ \lambda + 1 & 1 & -2\lambda - 1 & \lambda \\ -\lambda & -2\lambda - 1 & -1 & \lambda + 1 \\ 1 & 3\lambda & 3(\lambda + 1) & 2\lambda + 1 \end{pmatrix}, \\ U &: \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

The dual representation is given by the same formulas, where ε is replaced with ε^2 .

We will use the coordinates

$$(x_0 = u^3, x_1 = u^2v, x_2 = uv^2, x_3 = v^3).$$

Let $\mathcal{N}_1 \subset S^2\mathbb{U}_4^\vee$ be the linear space of quadratic forms such that $|\mathcal{N}_1|$ is the linear system of quadrics with the base locus equal to the Veronese curve R_3 . Obviously, it is an irreducible summand of $S^2\mathbb{U}_4^\vee$. It is generated by the quadratic forms

$$q_1 = x_0x_3 - x_1x_2, \quad q_2 = x_0x_2 - x_1^2, \quad q_3 = x_1x_3 - x_2^2.$$

Under the transformation S they are multiplied by $1, \varepsilon, \varepsilon^4$, respectively. The trace of S is equal to $1 + \varepsilon + \varepsilon^4 = \lambda$. Thus we can identify the space \mathcal{N}_1 with the linear representation $S^2\mathbb{V}$. Let \mathcal{N}_2^* be the linear space of quadratic forms in the dual space \mathbb{U}_4^\vee such that the linear system of quadrics $|\mathcal{N}_2^*|$ has the base locus equal to the dual Veronese curve R_3^* . It is spanned by the quadratic forms

$$q'_1 = 9\xi_0\xi_3 - \xi_1\xi_2, \quad q'_2 = 3\xi_0\xi_2 - \xi_1^2, \quad q'_3 = 3\xi_1\xi_3 - \xi_2^2,$$

where $\xi_0 = e_1^3, \xi_1 = e_1^2e_2, \xi_2 = e_1e_2^2, \xi_3 = e_2^3$ are the dual coordinates. The representation \mathcal{N}_2^* is isomorphic to $S^2\mathbb{V}'$. Consider the dual space $(\mathcal{N}_2^*)^\perp \subset S^2\mathbb{U}_4^\vee$ of apolar quadratic forms. It is spanned by the quadratic forms

$$p_1 = x_0x_3 + 9x_1x_2, \quad p_2 = 2x_0x_2 + 3x_1^2, \quad p_3 = 2x_1x_3 + 3x_2^2, \quad p_4 = x_0^2, \quad p_5 = x_3^2, \quad p_6 = x_0x_1, \quad p_7 = x_2x_3.$$

The linear span is a 7-dimensional summand of $S^2\mathbb{U}_4$. Computing the character of $S^2\mathbb{U}^\vee$, we find the decomposition

$$S^2\mathbb{U}_4^\vee \cong S^2\mathbb{U}_4 \cong \mathbb{W}_4 \oplus S^2\mathbb{V} \oplus S^2\mathbb{V}'. \tag{4.3}$$

This shows that

$$(\mathcal{N}_2^*)^\perp = \mathbb{W}_4 \oplus S^2\mathbb{V}'.$$

Observe that the quadratic forms p_i are eigenvectors of S with eigenvalues $1, \varepsilon, \varepsilon^4, \varepsilon^3, \varepsilon^2, \varepsilon^2, \varepsilon^3$, respectively. Since S has trace -1 on \mathbb{W}_4 , we find that the summand V_1 of $(\mathcal{N}_2^*)^\perp$ isomorphic to \mathbb{W}_4 is spanned by eigenvectors with eigenvalues $\varepsilon, \varepsilon^2, \varepsilon^3, \varepsilon^4$. Under the transformation T , the forms (p_1, \dots, p_7) are transformed to $(-p_1, p_3, p_2, p_5, p_4, -p_7, -p_6)$. Now we use the following known fact due to T. Reye (see [14], Lemma 4.3).

Theorem 2 (T. Reye). *A general 6-dimensional linear space L of quadrics in $\mathbb{P}^3 = |E|$ contains precisely two nets $\mathcal{N}_1, \mathcal{N}_2$ with the base loci equal to Veronese curves C_1, C_2 of degree 3. The dual space L^\perp of apolar quadrics in $|E^\vee|$ contains 10 quadrics Q_i with one-dimensional singular locus ℓ_i . Each line ℓ_i is a common secant of C_1, C_2 .*

It follows from the lemma that the summand of $S^2\mathbb{U}_4^\vee$ isomorphic to $S^2\mathbb{V}'$ is isomorphic to the linear space \mathcal{N}_2 such that the base locus of the net of quadrics $|\mathcal{N}_2|$ is a rational normal cubic curve. It is easy to see that this is possible only if \mathcal{N}_2 is generated by

$$r_1 = x_0x_3 + 9x_1x_2, \quad r_2 = x_0^2 + ax_2x_3, \quad r_3 = x_3^2 - ax_0x_1,$$

where $a^2 = 9$. Since the image of the point $[0, 1, 0, 0]$ on the curve under the transformation T must be on the curve, we check that $a = 3$. Thus the base locus is a rational normal cubic curve with parametric equations

$$|\mathbb{U}| \rightarrow |\mathbb{U}_4|, \quad (u, v) \mapsto (9u^2v, 27v^3, -u^3, 27uv^2). \tag{4.4}$$

Note that one can avoid using Reye's Theorem by using decomposition (4.3) and the facts there exist no \mathfrak{A}_5 -invariant lines nor \mathfrak{A}_5 -invariant conics, and that \mathfrak{A}_5 -orbits are of length ≤ 8 .

Now we are ready to describe the pencil $|S^4\mathbb{U}_4^\vee|$ of invariant quartics. We take as generators of the pencil the tangential ruled surfaces of the invariant rational normal curves defined by the nets $|\mathcal{N}_1|$ and $|\mathcal{N}_2|$. They are obviously invariant with respect to the action of G in $|\mathbb{W}_4|$. The net $|\mathcal{N}_1|$ defines a map from $|\mathbb{W}_4|$ to $|\mathcal{N}_1^\vee| = |S^2\mathbb{V}^\vee|$ whose image is an invariant conic, the *fundamental conic*. Its equation in Klein's coordinates A_0, A_1, A_2 is $A_0^2 + A_1A_2 = 0$. The equation of the dual conic in the dual coordinates A'_0, A'_1, A'_2 in the dual plane $|S^2\mathbb{U}^\vee|$ is

$$A'_0{}^2 + 4A'_1A'_2 = 0.$$

It follows that the equation of the tangential ruled surface must be of the form $Q_0^2 + 4Q_1Q_2$, where $Q_i = 0$ are equations of quadrics from the net $|\mathcal{N}_1|$. We know that A_0, A_1, A_2 are eigenvectors for the transformation S acting in $|S^2\mathbb{U}|$ with eigenvalues $1, \varepsilon, \varepsilon^{-1}$. Thus A'_0, A'_1, A'_2 are eigenvectors for the action of S in $S^2\mathbb{U}^\vee$ with the eigenvalues $1, \varepsilon^{-1}, \varepsilon$. Therefore our quadrics Q_0, Q_1, Q_2 are also eigenvectors for the action of S in $S^2\mathbb{W}_4^\vee$ with eigenvalues $1, \varepsilon^{-1}, \varepsilon$. We find that $(Q_0, Q_1, Q_2) = (q_1, \lambda q_3, \lambda' q_2)$ for some constants λ, λ' . Since the equation is invariant with respect to the transformation U , and $A'_1 \mapsto -A'_2$, we must have $\lambda = -\lambda'$. We find the condition on λ that guarantees that the equation $Q_0^2 - \lambda^2 Q_1 Q_2 = 0$ is the equation of the tangential ruled surface of the Veronese curve $C_1 = R_3$.

At each point $[1, t_0, t_0^2, t_0^3]$ the tangent line to the Veronese curve R_3 is spanned by the vector $[0, 1, 2t_0, 3t_0^2]$. Plugging in the parametric equation $s(1, t_0, t_0^2, t_0^3) + r(0, 1, 2t_0, 3t_0^2)$ of the tangent line in the equation $q_1^2 + c^2 q_2 q_3 = 0$, we obtain that $c^2 - 4 = 0$. Thus the equation of the quartic tangential ruled surface is

$$S_1 : (x_0x_3 - x_1x_2)^2 - 4(x_0x_2 - x_1^2)(x_1x_3 - x_2^2) = x_0^2x_3^2 - 6x_0x_1x_2x_3 + 4x_0x_3^2 + 4x_1x_3^3 - 3x_1^2x_2^2 = 0. \tag{4.5}$$

The second invariant quartic is the tangential ruled surface of the rational normal curve C_2 defined parametrically in (4.4). The net of quadrics containing this curve is G -equivariantly isomorphic to $S^2\mathbb{V}'$. It defines the map from $|\mathbb{U}_4|$ to $|(S^2\mathbb{V}')^\vee|$. The invariant conic in this space is the same conic as in the previous case. Similarly to this case we get the equation

$$(x_0x_3 + 9x_1x_2)^2 + d^2(x_0^2 + 3x_2x_3)(x_3^2 - 3x_0x_1) = 0. \tag{4.6}$$

The parametric equation of the tangent line to the base curve C_2 is

$$s(9t_0^2, 27, -t_0^3, 27t_0) + r(6t_0, 0, -t_0^2, 9) = 0.$$

Plugging this in the equation (4.6), we obtain $d^2 + 4 = 0$ and the equation of the second tangential surface is

$$S_2 : (x_0x_3 + 9x_1x_2)^2 - 4(x_0^2 + 3x_2x_3)(x_3^2 - 3x_0x_1) = 3(4x_0^3x_1 - x_0^2x_3^2 + 18x_0x_1x_2x_3 + 27x_1^2x_2^2 - 4x_2x_3^3) = 0.$$

So our pencil $|(S^4\mathbb{U}_4^\vee)^{2 \cdot \mathfrak{A}_5}|$ can be explicitly written in the form $\lambda F_1 + \mu F_2 = 0$, where

$$F_1 = x_0^2x_3^2 - 6x_0x_1x_2x_3 + 4x_0x_3^2 + 4x_1x_3^3 - 3x_1^2x_2^2 \quad \text{and} \\ F_2 = 4x_0^3x_1 - x_0^2x_3^2 + 18x_0x_1x_2x_3 + 27x_1^2x_2^2 - 4x_2x_3^3.$$

Consider the linear transformation

$$K : (x_0, x_1, x_2, x_3) \mapsto \left(\sqrt{3}x_2, \frac{1}{\sqrt{3}}x_0, -\frac{1}{\sqrt{3}}x_3, \sqrt{3}x_1 \right). \tag{4.7}$$

We have $K^2 = U \in G$, and the group generated by K and G is isomorphic to $2 \cdot \mathfrak{S}_5$. We immediately check that it transforms

$$x_0x_3 + 9x_1x_2 \mapsto 3(-x_0x_3 + x_1x_2), \quad x_0^2 + 3x_2x_3 \mapsto 3(x_1^2 - x_0x_2), \quad x_3^2 - 3x_0x_1 \mapsto 3(x_2^2 - x_1x_3).$$

This shows that $K(S_2) = S_1$, more precisely, we get $K(F_2) = 3F_1$. Thus the surfaces

$$S_3 = V(3F_1 + F_2) = V(x_0^2x_3^2 + 2x_0^3x_1 + 9x_1^2x_2^2 - 2x_2x_3^3 + 6x_0x_3^2 + 6x_1^3x_3) \quad \text{and} \tag{4.8} \\ S_4 = V(F_2 - 3F_1) = V(-x_0^2x_3^2 + 9x_0x_1x_2x_3 - 3x_0x_3^2 - 3x_1^3x_3 + 9x_1^2x_2^2 + x_0^3x_1 - x_2x_3^3).$$

are \mathfrak{S}_5 -invariant. One checks, using MAPLE, that they are nonsingular.

The following result is proven in [5].

Proposition 4. *The pencil generated by the quartic surfaces S_1 and S_2 contains two more singular members; each of them has 10 ordinary double points. The base curve B of the pencil is an irreducible curve of degree 16 with 24 singular points, and each of them is an ordinary cusp. It also contains a unique \mathfrak{A}_5 -orbit of 20 points.*

It is easy to see these two orbits of singular points of B . Each of them is the intersection of one of the rational normal cubic curves C_1 and C_2 with a nonsingular surface from the pencil. If we take the latter to be the surface S_3 or S_4 , we obtain that the union of the two orbits is invariant with respect to \mathfrak{S}_5 .

In the next section we will be able to see explicitly the unique orbit of 20 points.

5 An \mathfrak{A}_5 -invariant web of quadrics in $|\mathbb{U}_4|$

Consider the linear $2.\mathfrak{A}_5$ -equivariant map

$$\varphi : \mathbb{W}_4 \rightarrow S^2\mathbb{U}_4^\vee \quad (5.1)$$

defined by the decomposition (4.3). The corresponding map of the projective spaces defines a web of quadrics in $|\mathbb{U}_4|$. Recall that some of its attributes are the determinantal surface $D(\varphi)$ parametrizing singular quadrics in the web, and the Steinerian surface $\text{St}(\varphi)$ parametrizing singular points of singular quadrics from the web.

Let us find the equation of the discriminant surface. We choose a basis of $\varphi(\mathbb{W}_4)$ formed by the quadrics

$$Q_1 = 2x_1x_3 + 3x_2^2, \quad Q_2 = x_0^2 - 2x_2x_3, \quad Q_3 = x_3^2 + 2x_0x_1, \quad Q_4 = 3x_1^2 + 2x_0x_2.$$

They are transformed under the representation of $2.\mathfrak{A}_5$ (with the center acting trivially) as follows

$$\begin{aligned} S : (Q_1, Q_2, Q_3, Q_4) &\mapsto (\varepsilon^4 Q_1, \varepsilon^3 Q_2, \varepsilon^2 Q_3, \varepsilon Q_4), \\ T : \begin{pmatrix} Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{pmatrix} &\mapsto c \begin{pmatrix} -\lambda & 1 & \lambda^2 & \lambda \\ 1 & \lambda & -\lambda & \lambda^2 \\ \lambda^2 & -\lambda & \lambda & 1 \\ \lambda & \lambda^2 & 1 & -\lambda \end{pmatrix} \begin{pmatrix} Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{pmatrix}, \\ U : (Q_1, Q_2, Q_3, Q_4) &\mapsto (Q_4, Q_3, Q_2, Q_1), \end{aligned}$$

where c is some constant which will not be of concern for us. Choose a basis of $\varphi(\mathbb{W}_4) \subset S^2\mathbb{U}_4^\vee$ spanned by the following quadratic forms:

$$\begin{aligned} Q'_1 &= \varepsilon^4 Q_1 - \varepsilon^3 Q_2 - \varepsilon^2 Q_3 + \varepsilon Q_4, \\ Q'_2 &= \varepsilon^3 Q_1 - \varepsilon Q_2 - \varepsilon^4 Q_3 + \varepsilon^2 Q_4, \\ Q'_3 &= \varepsilon^2 Q_1 - \varepsilon^4 Q_2 - \varepsilon Q_3 - \varepsilon^3 Q_4, \\ Q'_4 &= \varepsilon Q_1 - \varepsilon^2 Q_2 - \varepsilon^3 Q_3 + \varepsilon^4 Q_4. \end{aligned} \quad (5.2)$$

The symmetric matrix defining the quadratic form $y_1 Q'_1 + y_2 Q'_2 + y_3 Q'_3 + y_4 Q'_4$ is equal to the matrix

$$A(y_1, y_2, y_3, y_4) = \begin{pmatrix} a_{11} & a_{12} & a_{13} & 0 \\ a_{21} & a_{22} & 0 & a_{24} \\ a_{31} & 0 & a_{33} & a_{34} \\ 0 & a_{42} & a_{43} & a_{44} \end{pmatrix},$$

where

$$\begin{aligned} a_{11} &= -(\varepsilon^3 y_1 + \varepsilon y_2 + \varepsilon^4 y_3 + \varepsilon^2 y_4), \\ a_{12} = a_{21} &= -(\varepsilon^2 y_1 + \varepsilon^4 y_2 + \varepsilon y_3 + \varepsilon^3 y_4), \\ a_{13} = a_{31} &= \varepsilon y_1 + \varepsilon^2 y_2 + \varepsilon^3 y_3 + \varepsilon^4 y_4, \\ a_{22} &= 3(\varepsilon y_1 + \varepsilon^2 y_2 + \varepsilon^3 y_3 + \varepsilon^4 y_4), \\ a_{24} = a_{42} &= \varepsilon^4 y_1 + \varepsilon^3 y_2 + \varepsilon^2 y_3 + \varepsilon y_4, \\ a_{33} &= 3(\varepsilon^4 y_1 + \varepsilon^3 y_2 + \varepsilon^2 y_3 + \varepsilon y_4), \\ a_{34} = a_{43} &= \varepsilon^3 y_1 + \varepsilon y_2 + \varepsilon^4 y_3 + \varepsilon^2 y_4, \\ a_{44} &= -(\varepsilon^2 y_1 + \varepsilon^4 y_2 + \varepsilon y_3 + \varepsilon^3 y_4). \end{aligned}$$

Plugging in these expressions and computing the determinant, we obtain (using MAPLE) the equation of the discriminant surface, where $y_5 = -y_1 - y_2 - y_3 - y_4$,

$$\det(A(y_1, y_2, y_3, y_4)) = 25 \left(\sum_{i=1}^5 y_i^4 - \sum_{1 \leq i < j \leq 5} y_i y_j^3 + \sum_{1 \leq i < j < k \leq 5} y_i y_j y_k^2 - 3 \sum_{1 \leq i < j < k < l \leq 5} y_i y_j y_k y_l \right),$$

where $y_5 = -(y_1 + y_2 + y_3 + y_4)$. We can rewrite this in terms of power functions to obtain the equation of the discriminant surface in the form

$$30 \sum_{i=1}^5 y_i^4 - 7 \left(\sum_{i=1}^5 y_i^2 \right)^2 = y_1 + y_2 + y_3 + y_4 + y_5 = 0. \tag{5.3}$$

This is the equation of the 10-nodal surface $S_{7/30}$ from the Hashimoto pencil. (This identification of the discriminant surface with the surface $S_{7/30}$ was also confirmed to me by S. Mukai.)

Remark 2. The 10 nodes of the symmetric discriminant quartic (also known as a *Cayley symmetroid quartic*) correspond to the singular lines of 10 reducible quadrics in the web. According to A. Coble [6], p. 250, they are the 10 common secants of the rational normal curves C_1 and C_2 .

Let us now compute the equation of the Steinerian surface of the web of quadrics defined by the map (5.1). If we choose the basis of $\varphi(W^4)$ given by (5.2), then the equation of the Steinerian surface is given by the determinant of the matrix with columns $Q'_1 \cdot \mathbf{x}, Q'_2 \cdot \mathbf{x}, Q'_3 \cdot \mathbf{x}, Q'_4 \cdot \mathbf{x}$, where we identify Q'_i with the associated symmetric matrix and \mathbf{x} is the column of the coordinates x_0, x_1, x_2, x_3 in \mathbb{U}_4 :

$$A = \begin{pmatrix} -\varepsilon^3 x_0 - \varepsilon^2 x_1 + \varepsilon x_2 & \varepsilon^4 x_3 - \varepsilon^2 x_0 + 3\varepsilon x_1 & 3\varepsilon^4 x_2 + \varepsilon^3 x_3 + \varepsilon x_0 & \varepsilon^4 x_1 + \varepsilon^3 x_2 - \varepsilon^2 x_3 \\ -\varepsilon^4 x_1 + \varepsilon^2 x_2 - \varepsilon x_0 & -\varepsilon^4 x_0 + \varepsilon^3 x_3 + 3\varepsilon^2 x_1 & 3\varepsilon^3 x_2 + \varepsilon^2 x_0 + \varepsilon x_3 & -\varepsilon^4 x_3 + \varepsilon^3 x_1 + \varepsilon x_2 \\ -\varepsilon^4 x_0 + \varepsilon^3 x_2 - \varepsilon x_1 & 3\varepsilon^3 x_1 + \varepsilon^2 x_3 - \varepsilon x_0 & \varepsilon^4 x_3 + \varepsilon^3 x_0 + 3\varepsilon^2 x_2 & \varepsilon^4 x_2 + \varepsilon^2 x_1 - \varepsilon x_3 \\ \varepsilon^4 x_3 - \varepsilon^3 x_1 - \varepsilon^2 x_0 & 3\varepsilon^4 x_2 - \varepsilon^3 x_0 + \varepsilon x_3 & \varepsilon^4 x_1 + \varepsilon^2 x_3 + 3\varepsilon x_2 & -\varepsilon^3 x_3 + \varepsilon^2 x_2 + \varepsilon x_1 \end{pmatrix}^t$$

Computing the determinant of the matrix A , we find that the equation of the Steinerian surface coincides with the equation of the surface S_4 from (4.8).

The Steinerian surface S_4 contains 10 lines, the singular lines of 10 reducible members of the web corresponding to singular points of the discriminant surface $S_{7/30}$. These are of course the 10 common secants of the two rational normal curves C_1 and C_2 . The unique orbit of 20 points from the same proposition is of course the intersection points of the 10 common secants with the curves C_1 and C_2 .

Remark 3. Recall that a minimal nonsingular model of a Cayley quartic symmetroid is isomorphic to the K3-cover of an Enriques surface. In turn, the Enriques surface is isomorphic to a Reye congruence of bidegree $(7, 3)$ in the Grassmannian $G(2, 4)$; see [8]. This applies to our surface S_4 . The embedding $j : S_4 \hookrightarrow |\mathbb{U}_4|$ defines an invertible sheaf $\mathcal{L}_1 = j^* \mathcal{O}_{|\mathbb{U}_4|}(1)$. The birational morphism from $s : S_4 \rightarrow S_{7/30} \subset |W_4|$ whose image is the quartic symmetroid $S_{7/30}$ that inverts the rational map from the discriminant surface to the Steinerian surface defines another invertible sheaf $\mathcal{L}_2 = s^* \mathcal{O}_{|W_4|}(1)$. It is known that the covering involution of S_4 preserves the tensor product $\mathcal{L}_1 \otimes \mathcal{L}_2$. Since both maps are \mathfrak{A}_5 -equivariant, we obtain that the action of \mathfrak{A}_5 on $S_{7/30}$ descends to an action on the Enriques surface.

The following is a list of open questions:

- Find the values of the parameters corresponding to the 10-nodal members. Are these surfaces determinantal? What is the transcendental lattice of its minimal nonsingular member?
- Find more facts about the \mathfrak{S}_5 -invariant surface S_3 . Is it determinantal? What is its transcendental lattice?
- Find the transcendental lattice of the general member of the pencil.

6 The catalecticant quartic surface

There is another view of the Cayley symmetroid quartic surface $S_{7/30}$. First we have the decomposition of the linear representations of \mathfrak{A}_5

$$S^6 \mathbb{V}^\vee \cong S^6 \mathbb{V} \cong W_4 \oplus S^2 \mathbb{V}.$$

Comparing this with (4.3), we find an isomorphism

$$S^2(\mathbb{U}_4)^\vee \cong S^6 \mathbb{V} \oplus S^2 \mathbb{V}'.$$

The summand $S^6\mathbb{V}$ is isomorphic to the image of $S^2(\mathbb{U}_4)^\vee$ under the restriction to the Veronese curve R_6 of degree 3. The kernel $S^2\mathbb{V}'$ is isomorphic to the (linear) space of quadrics vanishing on the Veronese curve. Recall that each $f \in S^{2d}\mathbb{V}^\vee$ defines a linear map $a : S^d\mathbb{V} \rightarrow S^d\mathbb{V}^\vee$ (the *apolarity map*). We view a basis of $S^d\mathbb{V}$ as partial derivatives of the coordinates (u, v) in $S^d\mathbb{V}$ and apply the differential operator $f_d(\frac{\partial}{\partial u}, \frac{\partial}{\partial v})$ to f to obtain a binary form of degree d in u, v . In these coordinates, the determinant of the map is a polynomial of degree $d + 1$ in coefficients of the form f , called the *catalecticant* (see [9], 1.4.1). A general zero of this polynomial is a binary form of degree $2d$ that can be written as a sum of less than the expected number (equal to d) of powers of linear forms. In the projective space $|S^{2d}\mathbb{V}^\vee|$ this corresponds to the variety of d -secant subspaces of dimension $d - 1$ of the Veronese curve R_{2d} . In our case where $d = 3$ and the basis in $S^6\mathbb{V}^\vee$ is taken as in (4.1), we get a quartic polynomial

$$\text{Cat} = \det \begin{pmatrix} a_0 & a_1 & a_2 & a_3 \\ a_1 & a_2 & a_3 & a_4 \\ a_2 & a_3 & a_4 & a_5 \\ a_3 & a_4 & a_5 & a_6 \end{pmatrix}.$$

The zeros of this polynomial in $|S^6\mathbb{V}^\vee|$ is the variety $\text{Tri}(R_6)$ of trisecant planes of the Veronese curve R_6 . Let $S = \text{Tri}(R_6) \cap |\mathbb{W}_4|$; this is a quartic surface in $|\mathbb{W}_4|$. Let us see that it coincides with the Cayley symmetroid $S_{7/30}$ studied in the previous section. In coordinates, a direct computation shows that the binary form $f = \sum a_s \binom{6}{s} t_0^s t_1^{6-s} \in S^6\mathbb{V}^\vee$ corresponds to the quadric

$$Q = a_0y_0^2 + a_2y_1^2 + a_4y_2^2 + a_6y_3^2 + 2a_1y_0y_1 + 2a_2y_0y_2 + 2a_3y_0y_3 + 2a_5y_2y_3 + 2a_4y_3y_1 + 2a_3y_1y_2.$$

The condition that $\text{Cat}(f) = 0$ becomes the condition that $\text{Discr}(Q) = 0$. This shows that the catalecticant quartic becomes isomorphic to the discriminant quartic in $\mathbb{P}^6 = \mathbb{P}(N^\perp)$. The singular locus of the variety $\text{Tri}(R_6)$ is of degree 10 and it is isomorphic to the secant variety of R_6 . Intersecting it with $|\mathbb{W}_4|$ we obtain 10 singular points of our symmetroid $S_{7/30}$.

Note that another model of $S^6\mathbb{V}$ is the space of *harmonic cubics* used in [12] in $|S^2\mathbb{V}|$ with respect to the dual of the fundamental conic.

7 \mathfrak{A}_5 -invariant rational plane sextic

Let \mathcal{N}_1 and \mathcal{N}_2 be the nets of quadrics with base loci rational normal curves C_1 and C_2 defined by the parametric equations (4.2) and (4.4). Restricting $|\mathcal{N}_2|$ to C_1 we obtain a map $C_1 \rightarrow |\mathcal{N}_2^\vee| \cong |S^2\mathbb{V}'^\vee|$. We identify the plane $|S^2\mathbb{V}'^\vee|$ with the plane $|S^2\mathbb{V}'|$ via the \mathfrak{A}_5 -invariant conic. Using the basis of \mathcal{N}_2 formed by the quadrics $V(x_0x_3 + 9x_1x_2)$, $V(x_0^2 + 3x_2x_3)$ and $V(x_3^2 - 3x_0x_1)$, the map $\mathbb{P}^1 \rightarrow C_1 \rightarrow |S^2\mathbb{V}'^\vee|$ is given by

$$(u, v) \mapsto (z_0, z_1, z_2) = (10u^3v^3, u^6 + 3uv^5, v^6 - 3u^5v).$$

Recall that the quadratic polynomials r_1, r_2, r_3 defining the basis are eigenvectors of S with eigenvalues $(1, \varepsilon^3, \varepsilon^2)$ and hence proportional to Klein’s dual coordinates (A'_0, A'_1, A'_2) . Since they are also transformed as $(r_1, r_2, r_3) \mapsto (-r_1, r_3, r_2)$ under U , they are equal to $(c'A'_0, cA'_1, -cA'_2)$ for some constants c, c' . Also, we know that the dual conic $A_0'^2 + 4A_1'A_2' = 0$ parameterizes the singular quadrics in the net, and hence the equation of S_2 shows that we may assume that $c' = \pm 1, c = 2$. We noticed before that in our coordinates u, v , we have to choose $c' = -1$. Thus

$$(z_0, z_1, z_2) = (-A'_0, 2A'_1, -2A'_2).$$

The polarity isomorphism $|S^2\mathbb{U}| \rightarrow |S^2\mathbb{U}'^\vee|$ defined by the conic $A_0^2 + A_1A_2 = 0$ gives $(A_0, A_1, A_2) = (-2A'_0, A'_2, A'_1)$. Thus, in the Klein coordinates, the image of $\mathbb{P}^1 \rightarrow C_1 \rightarrow |S^2\mathbb{U}'^\vee| \rightarrow |S^2\mathbb{U}'|$ is equal to the curve Γ_1 with parametric equation

$$\Gamma_1 : (u, v) \mapsto (A_0, A_1, A_2) = (-5u^3v^3, -v^6 + 3u^5v, u^6 + 3uv^5).$$

This agrees with the parametric equation of an \mathfrak{A}_5 -invariant rational sextic curve found by R. Winger in [20], [21]. He uses slightly different coordinates $(x, y, z) = (A_1, A_2, A_0)$.

The pencil of \mathfrak{A}_5 -invariant sextics is spanned by the triple conic and the union of the six fundamental lines:

$$\mu(xy + z^2)^3 + \lambda z(x^5 + y^5 + z^5 + 5x^2y^2z - 5xyz^3 + z^5).$$

According to R. Winger [21], the curve Γ_1 is a unique rational curve in this pencil with its 10 nodes forming an orbit of \mathfrak{A}_5 . It corresponds to the parameters $(\lambda : \mu) = (27, 5)$. The equation of Γ_1 becomes

$$27z(x^5 + y^5) + 5x^3y^3 + 150x^2y^2z^2 - 120xyz^4 + 32z^6 = 0.$$

The action of \mathfrak{A}_5 in the dual plane $|S^2\mathbb{U}^\vee|$ has an orbit of 6 points. The corresponding lines in the original plane $|S^2\mathbb{U}|$ are the 6 lines with equations

$$A_0 = 0, \quad A_0 + \varepsilon^v A_1 + \varepsilon^{-v} A_2 = 0, \quad v = 0, \dots, 4.$$

Each of these *fundamental lines* intersects the curve Γ_1 at 2 points with multiplicity 3. The 12 intersection points are the images of the orbit of \mathfrak{A}_5 acting in \mathbb{P}^1 that can be taken as the vertices of the icosahedron. They are the zeros of the polynomial

$$\Phi_{12} = uv(v^{10} + 11u^5v^5 - u^{10}).$$

The 10 pairs of branches of the singular points of Γ_1 correspond to the \mathfrak{A}_5 -orbit of 20 points in \mathbb{P}^1 . They are the zeros of the polynomial

$$\Phi_{20} = u^{20} + v^{20} + 288(u^{15}v^5 - u^5v^{15} - 494u^{10}v^{10}).$$

The dual curve Γ_1^* of Γ_1 is a rational curve of degree 10 with parametric equation

$$(x^*, y^*, z^*) = (-10u^7v^3 - 5u^2v^8, 5u^8v^2 - 10u^3v^7, u^{10} - 14u^5v^5 - v^{10}).$$

Its equation can be found in [12], p. 83. Note that, via the fundamental conic, we can identify the dual planes $|S^2\mathbb{V}|$ and $|S^2\mathbb{V}^\vee|$. Thus we have the second rational sextic curve Γ_2 and its second dual curve Γ_2^* . It is a rational curve Γ_2^* of degree 10 with parametric equation

$$(x, y, z) = (-10(u^7v^3 - u^2v^8), 10(u^8v^2 - 2u^3v^7), u^{10} - 14u^5v^5 - v^{10}).$$

The pair of these curves corresponds to the pair of rational normal curves C_1, C_2 . This pairing of rational normal sextics is discussed in detail in [6], Chapter 4.

Since the line $z = 0$ intersects Γ_1 at two points with multiplicity 3, the Plücker formulas show that Γ_1^* , and hence also Γ_2^* , has six 4-fold multiple points. They are the fundamental points of \mathfrak{A}_5 . (Note that there is another \mathfrak{A}_5 -invariant rational curve of degree 10 with 36 double points; we will not be concerned with it.)

Remark 4. The blow-up of the 10 nodes of Γ_1 (or of Γ_2) is a rational *Coble surface* C with $|-K_C| = \emptyset$ but $|-2K_C| \neq \emptyset$ and consists of the proper transform of the sextic Γ_1 . It inherits the \mathfrak{A}_5 -symmetry of Γ_1 . It is known that a Coble surface with an irreducible anti-bicanonical curve is a degenerate member of a pencil of Enriques surfaces. As I was informed by S. Mukai, there is a pencil of Enriques surfaces containing \mathfrak{A}_5 in its automorphism group (acting linearly in its Fano embedding in \mathbb{P}^5) that contains among its members the Enriques surfaces with K3-covers $S_{1/2}$ and $S_{7/30}$ as well as the Coble surface C .

Remark 5. The parametric equation of the \mathfrak{A}_5 -invariant rational sextic Γ_1 appears in [13], p. 122. It is shown there that the curve Γ_1 is equal to the intersection of the tangent lines at the origin of the modular family of elliptic quintic curves in \mathbb{P}^4 with the eigenplane of the negation involution. It is also discussed in [1], pp. 751–752, where it is shown that there is a $(3 : 1)$ -map from the modular Bring’s curve $X_0(2, 5)$ (isomorphic to the intersection of the Clebsch diagonal cubic surface with the quadric $x_1^2 + \dots + x_5^2 = 0$) to the normalization of the curve Γ_1 that coincides with the forgetting map $X_0(2, 5) \rightarrow X(5)$. The twelve cusps of Γ_1 are its intersection points with the fundamental conic.

References

- [1] W. Barth, K. Hulek, R. Moore, Degenerations of Horrocks-Mumford surfaces. *Math. Ann.* **277** (1987), 735–755. MR901715 Zbl 0663.14026
- [2] G. Bini, B. van Geemen, Geometry and arithmetic of Maschke’s Calabi–Yau three-fold. *Commun. Number Theory Phys.* **5** (2011), 779–820. MR2905137 Zbl 1253.14035
- [3] B. L. Cerchiai, B. van Geemen, From qubits to E_7 . *J. Math. Phys.* **51** (2010), 122203, 25. MR2779172 Zbl 1314.81044
- [4] I. Cheltsov, V. Przyjalkowski, C. Shramov, Quartic double solids with icosahedral symmetry. *Eur. J. Math.* **2** (2016), 96–119. MR3454093 Zbl 06551210
- [5] I. Cheltsov, C. Shramov, Two rational nodal quartic 3-folds. *Q. J. Math.* **67** (2016), 573–601. MR3609847 Zbl 1370.14013
- [6] A. B. Coble, *Algebraic geometry and theta functions*, volume 10 of *American Mathematical Society Colloquium Publications*. Amer. Math. Soc. 1982. MR733252 Zbl 55.0808.02
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, *Atlas of finite groups*. Oxford Univ. Press 1985. MR827219 Zbl 0568.20001
- [8] F. R. Cossec, Reye congruences. *Trans. Amer. Math. Soc.* **280** (1983), 737–751. MR716848 Zbl 0541.14034
- [9] I. V. Dolgachev, *Classical algebraic geometry. A modern view*. Cambridge Univ. Press 2012. MR2964027 Zbl 1252.14001
- [10] K. Hashimoto, Period map of a certain $K3$ family with an \mathfrak{S}_5 -action. *J. Reine Angew. Math.* **652** (2011), 1–65. MR2787353 Zbl 1213.14070
- [11] D. Hilbert, S. Cohn-Vossen, *Anschauliche Geometrie*. Springer 1932. MR0430930 Zbl 58.0597.01
- [12] N. Hitchin, Vector bundles and the icosahedron. In: *Vector bundles and complex geometry*, volume 522 of *Contemp. Math.*, 71–87, Amer. Math. Soc. 2010. MR2681722 Zbl 1227.14036
- [13] K. Hulek, Projective geometry of elliptic curves. *Astérisque* no. **137** (1986), 143. MR845383 Zbl 0602.14024
- [14] K. Hulek, K. Ranestad, Abelian surfaces with two plane cubic curve fibrations and Calabi–Yau threefolds. In: *Complex analysis and algebraic geometry*, 275–316, de Gruyter 2000. MR1760881 Zbl 1078.14057
- [15] F. Klein, *Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade*. Reprint of the 1884 original. Birkhäuser, Teubner 1993. MR1315530 Zbl 0803.01037
- [16] S. Kondō, Enriques surfaces with finite automorphism groups. *Japan. J. Math. (N.S.)* **12** (1986), 191–282. MR914299 Zbl 0616.14031
- [17] H. Maschke, Ueber die quaternäre, endliche, lineare Substitutionsgruppe der Borchardt’schen Moduln. *Math. Ann.* **30** (1887), 496–515. MR1510459 Zbl 19.0140.01
- [18] S. Mukai, Finite groups of automorphisms of $K3$ surfaces and the Mathieu group. *Invent. Math.* **94** (1988), 183–221. MR958597 Zbl 0705.14045
- [19] Y. G. Prokhorov, Fields of invariants of finite linear groups. In: *Cohomological and geometric approaches to rationality problems*, 245–273, Birkhäuser 2010. MR2605172 Zbl 1204.14007
- [20] R. M. Winger, Self-Projective Rational Sextics. *Amer. J. Math.* **38** (1916), 45–56. MR1506275 JFM 46.0952.02
- [21] R. M. Winger, On the invariants of the ternary icosahedral group. *Math. Ann.* **93** (1925), 210–216. MR1512233 JFM 51.0120.01