

## 1 Connected topological spaces

**Definition 1.1. (Disconnected spaces; connected spaces).** A topological space  $(X, \mathcal{T})$  is *disconnected* if there exist disjoint nonempty open subsets  $A$  and  $B$  in  $X$  such that  $X = A \cup B$ . We call the sets  $A$  and  $B$  a *separation* of  $X$ . If no separation of  $X$  exists, then  $X$  is called *connected*.

A subset  $S$  of  $X$  is said to be *connected* if it is connected when viewed with the subspace topology  $(S, \mathcal{T}_S)$ . This means ...

**Example 1.2.** Determine which of the following topological spaces are connected.

(a)  $X = \mathbb{R}$ ,  $\mathcal{T} = \{(-\infty, a) \mid a \in \mathbb{R}\} \cup \{\mathbb{R}, \emptyset\}$

(b)  $X = \{a, b, c, d\}$ ,  $\mathcal{T} = \{\emptyset, \{a, b\}, \{c\}, \{a, b, c\}, \{d\}, \{a, b, d\}, \{c, d\}, \{a, b, c, d\}\}$

### In-class Exercises

1. Show that the following topological spaces (with the Euclidean metric) are disconnected.

(a)  $\{\frac{1}{n} \mid n \in \mathbb{N}\}$

(b)  $(0, 1) \cup \{5\}$

(c)  $\mathbb{Q}$

2. (a) Give an example of a connected topological space  $X$ , and a subset  $S \subseteq X$  that is disconnected.

(b) Give an example of a disconnected topological space  $X$ , and a subset  $S \subseteq X$  that is connected.

3. Prove that a topological space  $(X, \mathcal{T})$  is disconnected if and only if there is subset  $A$ , with  $\emptyset \subsetneq A \subsetneq X$ , that is both open and closed.

4. Consider  $\{0, 1\}$  as a topological space with the discrete topology. Show that a topological space  $(X, \mathcal{T})$  is disconnected if and only if there is a continuous **surjective** function  $X \rightarrow \{0, 1\}$ .

5. Prove the following (often useful) lemma:

**Lemma.** Let  $X$  be a topological space, and let  $A, B$  be a separation of  $X$ . Let  $S \subseteq X$ . If  $S$  is connected, then  $S \subseteq A$  or  $S \subseteq B$ .

6. Let  $X$  be a topological space, and let  $A_i, i \in I$ , be a collection of subsets of  $X$ . Suppose that  $A_i$  is connected for each  $i$ .

(a) Show by example that the union  $\bigcup_{i \in I} A_i$  may be disconnected.

(b) Suppose that  $\bigcap_{i \in I} A_i \neq \emptyset$ . Show that the union  $\bigcup_{i \in I} A_i$  is connected.

7. **(Optional)**. Consider  $\mathbb{R}$  with the Euclidean metric. Which of the following subsets are connected?

$$\{x \in \mathbb{R} \mid d(x, 1) < 1 \text{ or } d(x, -1) < 1\}$$

$$\{x \in \mathbb{R} \mid d(x, 1) \leq 1 \text{ or } d(x, -1) < 1\}$$

$$\{x \in \mathbb{R} \mid d(x, 1) \leq 1 \text{ or } d(x, -1) \leq 1\}$$

8. **(Optional)**. Consider the set  $X = \{a, b, c, d\}$ . For which of the following topologies  $\mathcal{T}$  is the topological space  $(X, \mathcal{T})$  connected?

(a)  $\mathcal{T} = \{\emptyset, \{c\}, \{a, b\}, \{a, b, c\}, \{a, b, c, d\}\}$

(b)  $\mathcal{T} = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{b, c\}, \{a, b, c\}, \{a, d\}, \{a, b, d\}, \{a, b, c, d\}\}$

9. **(Optional)**. Consider the following topologies on  $\mathbb{R}$ . Which of these topological spaces are connected?

(a) indiscrete topology

(f)  $\mathcal{T} = \{\mathbb{R}, \{0, 1\}, \{0\}, \{1\}, \emptyset\}$

(b) discrete topology

(g)  $\mathcal{T} = \{(a, \infty) \mid a \in \mathbb{R}\} \cup \{\emptyset\} \cup \{\mathbb{R}\}$

(c) Euclidean topology

(h)  $\mathcal{T} = \{A \mid A \subseteq \mathbb{R}, 0 \in A\} \cup \{\emptyset\}$

(d) cofinite topology

(i)  $\mathcal{T} = \{A \mid A \subseteq \mathbb{R}, 0 \notin A\} \cup \{\mathbb{R}\}$

(e)  $\mathcal{T} = \{\mathbb{R}, (0, 1), \emptyset\}$

(j)  $\mathcal{T} = \{A \mid A \subseteq \mathbb{R}, 1 \in A\} \cup \{\emptyset\}$

10. **(Optional)**. Let  $(X, \mathcal{T})$  be a topological space. Let  $\{A_n \mid n \in \mathbb{Z}\}$  be a family of connected subspaces of  $X$  such that  $A_n \cap A_{n+1} \neq \emptyset$  for every  $n$ . Prove  $\bigcup_{n \in \mathbb{Z}} A_n$  is connected.
11. **(Optional)**. Let  $(X, \mathcal{T})$  be a topological space. Let  $\{A_n \mid n \in \mathbb{N}\}$  be a family of connected subspaces in  $X$  such that  $A_{n+1} \subseteq A_n$  for every  $n \in \mathbb{N}$ . Is  $\bigcap_{n \in \mathbb{N}} A_n$  necessarily connected?
12. **(Optional)**. Let  $(X, \mathcal{T})$  be a topological space, and let  $A, B \subseteq X$ . Suppose  $A \cup B$  and  $A \cap B$  are connected. Prove that if  $A$  and  $B$  are both closed or both open, then  $A$  and  $B$  are connected.