## 1 Compact topological spaces

Recall the definition of an open cover:

**Definition 1.1. (Open covers; open subcovers.)** Let  $(X, \mathcal{T})$  be a topological space. A collection  $\{U_i\}_{i \in I}$  of open subsets of X is an *open cover* of X if  $X = \bigcup_{i \in I} U_i$ . In other words, every point in X lies in the set  $U_i$  for some  $i \in I$ .

A sub-collection  $\{U_i\}_{i \in I_0}$  (where  $I_0 \subseteq I$ ) is an open subcover (or simply subcover) if  $X = \bigcup_{i \in I_0} U_i$ . In other words, every point in X lies in some set  $U_i$  in the subcover.

**Definition 1.2. (Compact spaces; compact subspaces.)** We say that a topological space  $(X, \mathcal{T})$  is *compact* if **every** open cover of X has a finite subcover.

A subset  $A \subseteq X$  is called *compact* if it is compact with respect to the subspace topology. This means ...

**Example 1.3.** Let  $(X, \mathcal{T})$  be a finite topological space. Then X is compact.

**Example 1.4.** Let X be a topological space with the indiscrete topology. Then X is compact.

**Example 1.5.** Let X be an infinite topological space with the discrete topology. Then X is **not** compact.

## **In-class Exercises**

- 1. (a) Let X be a set with the cofinite topology. Prove that X is compact.
  - (b) Let X = (0, 1) with the topology induced by the Euclidean metric. Show that X is not compact.
- 2. Suppose that  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  are topological spaces, and  $f : X \to Y$  is a continuous map. Show that, if X is compact, then f(X) is a compact subspace of Y. In other words, the continuous image of a compact set is compact.
- 3. Prove that a closed subset of a compact space is compact.
- 4. (a) Let X be a topological space with topology induced by a metric d. Prove that any compact subset A of X is bounded.
  - (b) Suppose that  $(X, \mathcal{T})$  is a **Hausdorff** topological space. Prove that any compact subset A of X is closed in X.
  - (c) Consider  $\mathbb{Q}$  with the Euclidean metric. Show that the subset  $(-\pi,\pi) \cap \mathbb{Q}$  of  $\mathbb{Q}$  is closed and bounded, but not compact.
- 5. (Optional). Determine which of the following topologies on  $\mathbb{R}$  are compact.
  - Any topology T consisting of only finitely many sets.
    T = {(a,∞) | a ∈ ℝ} ∪ {Ø} ∪ {ℝ}
    T = {A | A ⊆ ℝ, 0 ∈ A} ∪ {Ø}
  - the discrete topology  $\mathcal{T} = \{A \mid A \subseteq \mathbb{R}, \ 0 \notin A\} \cup \{\mathbb{R}\}$
- 6. (Optional). Consider  $\mathbb{R}$  with the topology  $\mathcal{T} = \{A \mid A \subseteq \mathbb{R}, 0 \notin A\} \cup \{\mathbb{R}\}$ . Give necessary and sufficient conditions for a subset  $C \subseteq \mathbb{R}$  to be compact.
- 7. (Optional). Let X be a nonempty set, and let  $x_0$  be a distinguished element of X. Let

 $\mathcal{T} = \{ A \subseteq X \mid x_0 \notin A \text{ or } X \setminus A \text{ is finite } \}.$ 

- (a) Show that  $\mathcal{T}$  defines a topology on X.
- (b) Verify that  $(X, \mathcal{T})$  is Hausdorff.
- (c) Verify that  $(X, \mathcal{T})$  is compact.

This exercise shows that **any** nonempty set X admits a topology making it a compact Hausdorff topological space.

- 8. (Optional). Let  $K_1 \supseteq K_2 \supseteq \cdots$  be a descending chain of nonempty, closed, compact sets. Then  $\bigcap_{n \in \mathbb{N}} K_n \neq \emptyset$ .
- 9. (Optional). Let X be a topological space, and let  $A, B \subseteq X$  be compact subsets.
  - (a) Suppose that X is Hausdorff. Show that  $A \cap B$  is compact.
  - (b) Show by example that, if X is not Hausdorff,  $A \cap B$  need not be compact. *Hint:* Consider  $\mathbb{R}$  with the topology  $\{U \mid U \subseteq \mathbb{R}, 0, 1 \notin U\} \cup \{\mathbb{R}\}.$
- 10. (Optional). Suppose that  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  are topological spaces, and  $f : X \to Y$  is a closed map (this means that f(C) is closed for every closed subset  $C \subseteq X$ ). Suppose that Y is compact, and moreover that  $f^{-1}(y)$  is compact for every  $y \in Y$ . Prove that X is compact.