

Terms and concepts covered: Singular homology is a homotopy invariant. Singular homology is functorial. Homology of S^n and the Brouwer fixed point theorem for D^n . Diagram chases; the short five lemma. Relative chains, relative cycles, relative boundaries. Relative homology groups. Long exact sequence of a pair. "A short exact sequence of chain complexes induces a long exact sequence of homology groups". Excision theorem. $H_n(X, A) \cong \tilde{H}_n(X/A)$ for a good pair (X, A) .

Corresponding reading: Hatcher Ch 2.1, Exact sequences and excision, Ch 2.A Homology and fundamental group.

Warm-up questions

(These warm-up questions are optional, and won't be graded.)

1. Let

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

be a short exact sequence of abelian groups. The group B is called an *extension* of C by A . The question of classifying the possible groups B given the groups A and C is called the *extension problem*. Solve the extension problem for the following short exact sequences.

$$0 \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow ? \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

$$0 \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow ? \longrightarrow \mathbb{Z}/3\mathbb{Z} \longrightarrow 0$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow ? \longrightarrow \mathbb{Z} \longrightarrow 0$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow ? \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

$$0 \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow ? \longrightarrow \mathbb{Z} \longrightarrow 0$$

2. We proved in class that $\tilde{H}_k(S^n) = \begin{cases} \mathbb{Z}, & k = n \\ 0, & k \neq n. \end{cases}$

(a) Conclude that S^n and S^m are not homotopy equivalent unless $n = m$.

(b) Prove that \mathbb{R}^n and \mathbb{R}^m are not homeomorphic.

Hint: Consider a hypothetical homeomorphism $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and its restriction to $\mathbb{R}^n \setminus \{0\}$.

3. Let $A \subseteq X$ be spaces. Recall that we defined $C_n(X, A)$ as the quotient of singular chain groups $C_n(X)/C_n(A)$.

(a) Explain why we can identify $C_n(X, A)$ with free abelian group of singular n -chains $\Delta^n \rightarrow X$ with image not fully contained in A .

(b) Explain why we did not define $C_n(X, A)$ as the subgroup of $C_n(X)$ spanned by singular n -chains $\Delta^n \rightarrow X$ with image not fully contained in A .

Hint: Is this a sub-chain complex? What happens when you restrict ∂ ?

4. Let X be a space, and $*$ $\in X$.

(a) Show that

$$H_n(X, X) = 0 \quad \text{for all } n.$$

(b) Show that

$$H_n(X, \emptyset) \cong H_n(X) \quad \text{for all } n.$$

(c) Show that

$$H_n(X, *) \cong \tilde{H}_n(X) \quad \text{for all } n.$$

5. Suppose that (X, A) is a good pair of spaces, and that A is contractible. Use the long exact sequence of a pair to prove that

$$\tilde{H}_n(X) \cong \tilde{H}_n(X/A) \quad \text{for all } n.$$

6. Let $f : (X, A) \rightarrow (Y, B)$ be a map of pairs (Assignment Problem 2). Verify that the induced map on relative homology is functorial.
7. Let $f : X \rightarrow Y$ be a nullhomotopic map. Show that the induced map $f_* : \tilde{H}_n(X) \rightarrow \tilde{H}_n(Y)$ is zero for all n . What are the induced maps $f_* : H_n(X) \rightarrow H_n(Y)$?

Assignment questions

(Hand these questions in!)

1. In this problem, we will prove one of the foundational results of homological algebra. You may consult Hatcher p116-118 as much as you would like while you write your proof, but you should make a good-faith effort to attempt the diagram chases on your own!

Slogan: A short exact sequence of chain complexes gives rise to a long exact sequence of homology groups.

The complete theorem statement is as follows.

Theorem (The LES on homology of an SES of chain complexes). Let (A_*, d^A) , (B_*, d^B) and (C_*, d^C) be chain complexes. Let

$$0 \longrightarrow A_* \xrightarrow{i_\#} B_* \xrightarrow{j_\#} C_* \longrightarrow 0$$

be a short exact sequence of these chain complexes. In other words, $i_\#$ and $j_\#$ are chain maps such that the sequence of group homomorphisms

$$0 \longrightarrow A_n \xrightarrow{i_n} B_n \xrightarrow{j_n} C_n \longrightarrow 0$$

is exact for all n . Then there exists a long exact sequence on homology groups

$$\cdots \longrightarrow H_n(A_*) \xrightarrow{i_*} H_n(B_*) \xrightarrow{j_*} H_n(C_*) \xrightarrow{\delta} H_{n-1}(A_*) \xrightarrow{i_*} H_{n-1}(B_*) \xrightarrow{j_*} H_{n-1}(C_*) \longrightarrow \cdots$$

The map δ is called the *connecting homomorphism*.

Remark: This result is called the *Zig-Zag Lemma* or *Snake Lemma*, due to the shape of the connecting homomorphism when the result is formulated as in this diagram.

Remark: Implausibly, a construction of the connecting homomorphism appeared at the beginning of a 1980 romantic comedy. You can watch the clip [here](#). The film was nominated for a Razzie Award for Worst Screenplay.

- (a) The following commutative diagram shows our short exact sequence of chain complexes.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \cdots & \longrightarrow & A_{n+1} & \xrightarrow{d_{n+1}^A} & A_n & \xrightarrow{d_n^A} & A_{n-1} \longrightarrow \cdots \\
 & & \downarrow i_\# & & \downarrow i_\# & & \downarrow i_\# \\
 \cdots & \longrightarrow & B_{n+1} & \xrightarrow{d_{n+1}^B} & B_n & \xrightarrow{d_n^B} & B_{n-1} \longrightarrow \cdots \\
 & & \downarrow j_\# & & \downarrow j_\# & & \downarrow j_\# \\
 \cdots & \longrightarrow & C_{n+1} & \xrightarrow{d_{n+1}^C} & C_n & \xrightarrow{d_n^C} & C_{n-1} \longrightarrow \cdots \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Given a cycle $c \in C_n$, explain how to construct a cycle $\delta(c)$ in A_{n-1} , and verify that your map is well-defined as a map on homology $\delta : H_n(C_*) \rightarrow H_{n-1}(A_*)$.

(b) Verify that δ is a group homomorphism.

(c) Verify that the sequence

$$\cdots \longrightarrow H_n(A_*) \xrightarrow{i_*} H_n(B_*) \xrightarrow{j_*} H_n(C_*) \xrightarrow{\delta} H_{n-1}(A_*) \xrightarrow{i_*} H_{n-1}(B_*) \xrightarrow{j_*} H_{n-1}(C_*) \longrightarrow \cdots$$

is exact.

(d) Given a space X , let $\tilde{C}_*(X)$ denote the augmented singular chain complex of X . For space $A \subseteq X$, let $i : A \rightarrow X$ denote the inclusion map. Verify that the following is a short exact sequence of chain complexes

$$0 \longrightarrow \tilde{C}_*(A) \xrightarrow{i_\#} \tilde{C}_*(X) \xrightarrow{j_\#} C_*(X, A) \longrightarrow 0$$

where $j_\#$ is the quotient map $C_*(X) \rightarrow C_*(X)/C_*(A) = C_*(X, A)$. Deduce the existence of the long exact sequence of a pair.

(e) Verify the following comment of Hatcher (p117):

If a class $[\alpha] \in H_n(X, A)$ is represented by a relative cycle α , then $\delta[\alpha]$ is the class of the cycle $\partial\alpha$ in $H_{n-1}(A)$.

For this reason, in the long exact sequence of a pair the connecting homomorphism is sometimes called the *boundary map*.

2. (a) **Definition (Map of pairs).** Let $A \subseteq X$ and $B \subseteq Y$ be spaces. A *map of pairs* $(X, A) \rightarrow (Y, B)$ is a continuous map $f : X \rightarrow Y$ such that $f(A) \subseteq B$.

Verify that a map of pairs $f : (X, A) \rightarrow (Y, B)$ induces a chain map $f_\# : C_*(X, A) \rightarrow C_*(Y, B)$. It follows that it induces a map on homology (see Warm-up Problem 6).

(b) Let $B \subseteq A \subseteq X$ be spaces. Show that there is a short exact sequence of chain complexes

$$0 \longrightarrow C_*(A, B) \longrightarrow C_*(X, B) \longrightarrow C_*(X, A) \longrightarrow 0$$

(c) Write down the corresponding long exact sequence on homology groups. It is called the *long exact sequence of a triple*.

(d) By modifying our proof of the analogous statement for absolute homology, we can prove the following. (You do not need to check this).

Proposition (Homotopic maps of pairs induce the same map on relative homology).

Suppose two maps of pairs $f, g : (X, A) \rightarrow (Y, B)$ are homotopic through maps of pairs.

Then for all n , the induced maps $f_*, g_* : H_n(X, A) \rightarrow H_n(Y, B)$ are equal.

Let $B \subseteq A \subseteq X$, and suppose that A deformation retracts to B . Show that $H_n(A, B) = 0$ for all n , and use the long exact sequence of a triple to conclude that $H_n(X, B) \cong H_n(X, A)$ for all n .

Remark: We use this result in the proof that $H_n(X, A) \cong \tilde{H}_n(X/A)$ when (X, A) is a good pair.

Remark: We will not prove the following property, but it is useful to know:

Proposition (Naturality of a map of pairs). Let $f : (X, A) \rightarrow (Y, B)$ be a map of pairs. Then the long exact sequence of a pair is *natural* in the sense that the following diagram commutes.

$$\begin{array}{ccccccccccc} \cdots & \longrightarrow & H_n(A) & \xrightarrow{i_*} & H_n(X) & \xrightarrow{j_*} & H_n(X, A) & \xrightarrow{\delta} & H_{n-1}(A) & \longrightarrow & \cdots \\ & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* & & \downarrow f_* & & \\ \cdots & \longrightarrow & H_n(B) & \xrightarrow{i_*} & H_n(Y) & \xrightarrow{j_*} & H_n(B, Y) & \xrightarrow{\delta} & H_{n-1}(B) & \longrightarrow & \cdots \end{array}$$

3. In this problem, we will complete our proof of the following theorem.

Theorem ($H_1(X) \cong \pi_1(X, x_0)^{ab}$). Let X be path-connected space with basepoint x_0 . There is a surjective group homomorphism

$$\begin{aligned} h : \pi_1(X, x_0) &\longrightarrow H_1(X) \\ [\gamma] &\longmapsto \text{singular 1-chain } \gamma \end{aligned}$$

whose kernel is the commutator subgroup of $\pi_1(X, x_0)$. In particular,

$$H_1(X) \cong \pi_1(X, x_0)^{ab}.$$

Last week, we proved that the Hurewicz map h is a group homomorphism, that h surjects, and that the kernel of h contains the commutator subgroup of $\pi_1(X, x_0)$. It remains to show that the commutator subgroup contains $\ker(h)$.

You may read Hatcher 2.A and other relevant sections while you write your solutions.

- (a) Read the three paragraphs in Hatcher before Proposition 2.6, starting at the bottom of p108 with “Cycles in singular homology are defined algebraically, but they can be given a somewhat more geometric interpretation . . .”

Explain in your own words the construction K_ξ in the case $n = 2$, and why “elements of $H_2(X)$ are represented by maps of closed oriented surfaces into X .”

- (b) Let $[\gamma] \in \pi_1(X, x_0)$ be a loop such that $h([\gamma]) = 0$. Our goal is to show that $[\gamma]$ is in the commutator subgroup of $\pi_1(X, x_0)$. Since $h([\gamma]) = 0$, the loop γ (viewed as a 1-cycle) is the boundary of some 2-chain $\sum_i n_i \sigma_i$. Let $\tau_{i,j}$ denote the singular 1-simplices defined by

$$\partial \sigma_i = \tau_{i,0} - \tau_{i,1} + \tau_{i,2}$$

thus

$$\gamma = \partial \left(\sum_i n_i \sigma_i \right) = \sum_{i,j} (-1)^j n_i \tau_{i,j}.$$

Explain why we can group all the 1-simplices into cancelling pairs, with one remaining term $\tau_{i,j}$ equal to γ .

- (c) Explain why we can associate the chain $\sum_i n_i \sigma_i$ to a map $\sigma : K \rightarrow X$ from a 2-dimensional Δ -complex K with one boundary loop corresponding to γ .
- (d) Explain how we can use the homotopy extension property (Homework 3 Problem 4(d)) to homotope σ (rel the edge corresponding to γ) so that all of its vertices map to x_0 .
- (e) Deduce that we obtain a new map $\sigma' : K \rightarrow X$ associated to a new 2-chain $\sum_i n_i \sigma'_i$ with

$$\gamma = \partial \left(\sum_i n_i \sigma'_i \right) = \sum_{i,j} (-1)^j n_i \tau'_{i,j}$$

and every 1-simplex $\tau'_{i,j}$ a loop based at x_0 .

- (f) Let $[[\tau'_{i,j}]]$ denote the image of $\tau'_{i,j}$ in $\pi_1(X, x_0)^{ab}$. Explain why $\sum_{i,j} (-1)^j n_i [[\tau'_{i,j}]]$ is zero in $\pi_1(X, x_0)^{ab}$. *Hint:* The boundary of a 2-simplex is zero in π_1 .
- (g) Conclude that γ is zero in $\pi_1(X, x_0)^{ab}$, hence, $[\gamma]$ is in the commutator subgroup. This concludes our proof of the Hurewicz theorem for π_1 .
- (h) Let $f : (X, x_0) \rightarrow (Y, f(x_0))$ be a continuous map of topological spaces. Verify that the following diagram commutes,

$$\begin{array}{ccc} \pi_1(X, x_0) & \xrightarrow{f_*} & \pi_1(Y, f(x_0)) \\ \downarrow h & & \downarrow h \\ H_1(X) & \xrightarrow{f_*} & H_1(Y) \end{array}$$

and explain how this allows us to compute the map induced by f on H_1 using the map induced by f on π_1 .

Remark: The commutativity of this diagram is the statement that the Hurewicz homomorphism is a *natural transformation* from the functor π_1 to the functor H_1 [redefined to be a functor on $\underline{\text{Top}}^*$].

4. (a) Prove the following proposition.

Proposition (Homology of a wedge sum). Let $\{X_\alpha\}$ be a collection of topological spaces, with basepoint $x_\alpha \in X_\alpha$ such that (X_α, x_α) is a good pair for each α . Let $\bigvee_\alpha X_\alpha$ be the wedge sum formed by identifying the basepoints x_α , and let $i_\alpha : X_\alpha \rightarrow \bigvee_\alpha X_\alpha$ be the inclusion map. Then for each n there is an isomorphism on homology

$$\bigoplus_\alpha (i_\alpha)_* : \bigoplus_\alpha \tilde{H}_n(X_\alpha) \xrightarrow{\cong} \tilde{H}_n\left(\bigvee_\alpha X_\alpha\right)$$

Hint: Consider the pair $(\bigsqcup_\alpha X_\alpha, \bigsqcup_\alpha \{x_\alpha\})$.

- (b) State the homology groups of the following spaces. No justification needed.

- (i) $\bigvee_k S^1$
- (ii) a once-punctured torus
- (iii) $S^1 \vee S^2 \vee S^3 \vee S^\infty$
- (iv) the wedge sum of $\mathbb{R}P^2$ and a Mobius band
- (v) \mathbb{R}^2 with two points identified

5. (a) Verify that $S^1 \times S^1$ and $S^1 \vee S^1 \vee S^2$ have isomorphic homology groups in every degree.
 (b) Verify that $S^1 \times S^1$ and $S^1 \vee S^1 \vee S^2$ are not homotopy equivalent. Conclude that, although homology groups are homotopy invariants, they are not complete invariants.
 (c) Show that the universal covers of $S^1 \times S^1$ and $S^1 \vee S^1 \vee S^2$ do not have the same homology groups.

6. **Definition (Local homology groups).** Let X be a space. The *local homology groups* at a point $x \in X$ are the groups $H_n(X, X \setminus \{x\})$.

- (a) Let X be a space and let $x \in X$ be a point such that $\{x\}$ is closed in X . Use excision to show that, if U is any neighbourhood of x , then

$$H_n(X, X \setminus \{x\}) \cong H_n(U, U \setminus \{x\}).$$

Hence the groups only depend on the local topology of X near x .

- (b) Explain why a homeomorphism $f : X \rightarrow Y$ must induce isomorphisms

$$f_* : H_n(X, X \setminus \{x\}) \xrightarrow{\cong} H_n(Y, Y \setminus \{f(x)\}).$$

Conclude that the local homology groups must be equal if two spaces are locally homeomorphic at a pair of points.

- (c) Prove the following theorem.

Theorem (Invariance of dimension). Let $U \subseteq \mathbb{R}^n$ and $V \subseteq \mathbb{R}^m$ be nonempty open subsets. If U and V are homeomorphic, then $m = n$.

Hint: $H_n(U, U \setminus \{x\}) = H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\})$. Consider the long exact sequence of the pair $(\mathbb{R}^n, \mathbb{R}^n \setminus \{x\})$.

Wellbeing

(This section is completely optional. This is a nudge to prioritize your wellbeing during the pandemic.)

1. **(Health comes first)**. Make your health and happiness goals a priority this week.
2. **(Sleep quality)**. This week, re-evaluate your sleep habits. Getting enough sleep, and getting quality sleep, can be a challenge during the semester. Is it worth adjusting your schedule, or your priorities, in order to free more time to sleep? Could you take steps to improve the quality of your sleep? Some standard advice is to consider the following factors:
 - the comfort of your sleep environment – including light, temperature, noise, and bedding
 - use of screens or other blue-light sources within an hour of bedtime
 - exercise – enough during the day; not too much within two hours of bedtime
 - use of substances, including caffeine and alcohol
 - your bedtime routine
Experts recommend establishing a consistent bedtime routine, and (for those who need to unwind) incorporate relaxing activities like a warm bath, soothing music, low-stress reading, or meditation. Harness 'the power of ritual' to cue to your body it is time to sleep.

You can find more suggestions from the CDC's "Tips for better sleep".

There are resources online for managing stress and worrying. If your sleep (or life) quality is being impacted by a more serious problem such as severe anxiety, grief, depression, physical pain or discomfort, or another health concern, please be proactive and seek support. Some UM services are listed on our course webpage.