

**Terms and concepts covered:**  $n$ -simplex; vertices, subsimplices, and faces; boundary and interior of a simplex;  $\Delta$ -complex; chain complex,  $n$ -chains, exactness, homology groups; boundary homomorphisms, cycles and boundaries, simplicial homology groups, homology classes.

**Corresponding reading:** Hatcher Chapter 2, Introduction and Section 2.1, “ $\Delta$ -Complexes” and “Simplicial Homology”.

## Warm-up questions

(These warm-up questions are optional, and won't be graded.)

1. **Definition (Convex).** A subset  $C$  in Euclidean space is *convex* if it contains the line segment connecting any pair of its points.

**Definition (Convex combination, convex hull).** Let  $X$  be a subset of Euclidean space. A *convex combination* of points in  $X$  is a sum of the form

$$t_1x_1 + t_2x_2 + \cdots + t_nx_n \quad \text{such that } x_i \in X, t_i \in \mathbb{R}, t_i \geq 0, t_1 + \cdots + t_n = 1.$$

The *convex hull* of  $X$  is the set of all convex combinations of points in  $X$ .

- (a) Prove that the convex hull of  $X$  is the minimal (under inclusion) convex subset containing  $X$ .
  - (b) Prove that the convex hull of  $X$  is the intersection of all convex subsets containing  $X$ .
2. Let  $\Delta^n = [v_0, v_1, \dots, v_n]$  be the standard  $n$ -simplex,

$$\Delta^n = \left\{ (t_0, t_1, \dots, t_n) \in \mathbb{R}^{n+1} \mid t_i \geq 0, \sum_i t_i = 1 \right\}.$$

Show that the convex hull of any  $(k + 1)$  of its vertices is canonically homeomorphic to a  $k$ -simplex. Conclude that it therefore makes sense topologically (as well as combinatorially) to call this subspace a  $k$ -dimensional subsimplex.

3. Let  $\Delta^n = [v_0, v_1, \dots, v_n]$  be an  $n$ -dimensional simplex. For each  $k \leq n$ , how many  $k$ -dimensional subsimplices does  $\Delta^n$  have?
4. Describe the canonical  $\Delta$ -complex structure on an  $n$ -simplex. What is its  $k$ -skeleton?
5. In this question, we will find another way to coordinatize an  $n$ -simplex. Let

$$\Delta_*^0 = \{0\},$$

$$\Delta_*^n = \{(s_1, \dots, s_n) \in \mathbb{R}^n \mid 0 \leq s_1 \leq s_2 \leq \dots \leq s_n \leq 1\}$$

- (a) Draw  $\Delta_*^n$  for  $n = 0, 1, 2, 3$ .

- (b) Recall that we defined the standard simplex  $\Delta^n = \{(t_0, t_1, \dots, t_n) \in \mathbb{R}^{n+1} \mid t_i \geq 0, \sum_i t_i = 1\}$ . Show that  $\Delta^n$  is homeomorphic to  $\Delta_*^n$  via the map


$$s_i = t_0 + t_1 + \cdots + t_{i-1}.$$

6. Consider our coordinatization of the  $n$ -simplex,

$$\Delta_*^n = \{(s_1, \dots, s_n) \in \mathbb{R}^n \mid 0 \leq s_1 \leq s_2 \leq \dots \leq s_n \leq 1\}.$$

Prove that the boundary  $\partial\Delta^n$  and the open simplex  $\mathring{\Delta}^n$  are indeed the boundary and interior of  $\Delta^n$ , respectively, in the usual sense of point-set topology, when  $\Delta^n$  is viewed as the subset  $\Delta_*^n$  of  $\mathbb{R}^n$ .

7. (a) Verify that an  $n$ -simplex (as a topological space) is homeomorphic to a closed  $n$ -ball.  
 (b) Verify that a  $\Delta$ -complex structure on a space  $X$  is, in particular, a CW complex structure.
8. (a) Which of our standard CW complex structures on the spheres  $S^1$  and  $S^2$  are  $\Delta$ -complex structures?  
 (b) Is our standard CW complex structure on  $\mathbb{R}P^2$  a  $\Delta$ -complex structure?
9. Choose your preferred name for the  $\partial$  symbol.
10. Let  $X$  be either of the  $\Delta$ -complexes shown below. Let  $C_n(X)$  denote the associated  $n$ th simplicial chain group, and let  $\partial_n : C_n(X) \rightarrow C_{n-1}(X)$  be the boundary map.



- (a) We choose the total orderings of the vertices  $v_1, v_2, v_3, v_4$  and  $u_1, u_2, u_3, u_4$ , respectively, for the two complexes. Explain how this determines an order on the vertices of every simplex. Label the edges of each complex with the appropriate direction.
- (b) Compute the boundary (that is, the image under  $\partial_n$ ) of the following  $n$ -chains.
- $2[v_1, v_2, v_3]$
  - $[v_1, v_2, v_3] + [v_2, v_3, v_4]$
  - $[v_1, v_2, v_3] - [v_2, v_3, v_4]$
  - $[u_1, u_2] - [u_1, u_3] + [u_2, u_3]$
  - $[u_1, u_2] - [u_1, u_3] + [u_2, u_3] + [u_3, u_4]$
- (c) Explain for each calculation how this boundary relates to your intuitive geometric understanding of “boundary”.
11. Let  $X$  be a  $\Delta$ -complex. Let  $C_n(X)$  denote the  $n$ th simplicial chain group, and let  $\partial_n : C_n(X) \rightarrow C_{n-1}(X)$  be the boundary map.
- Verify that  $\partial_n \circ \partial_{n+1} = 0$ .
  - Give a geometric interpretation of the equation  $\partial_n \circ \partial_{n+1} = 0$ , in the spirit of “a boundary has no boundary”.
12. (a) Let  $u, v$  be vertices in a simplicial complex joined by an edge. What is the relationship between the (oriented) edge  $[u, v]$  and the (oriented) edge  $[v, u]$ ? What is the relationship between the 1-chains  $[u, v]$  and  $-[v, u]$ , and the relationships between their boundaries?
- (b) Let  $\Delta_n = [v_0, v_1, \dots, v_n]$  be an  $n$ -simplex, and  $\tau$  a permutation in  $S_{n+1}$ . Show that

$$\partial_n([v_{\tau(0)}, v_{\tau(1)}, \dots, v_{\tau(n)}]) = \begin{cases} \partial_n([v_0, v_1, \dots, v_n]) & \text{if } \tau \text{ is an even permutation} \\ -\partial_n([v_0, v_1, \dots, v_n]) & \text{if } \tau \text{ is an odd permutation.} \end{cases}$$

Conclude that our ordering of the vertices does matter in our computation of the differential—different orders result in different signs—but order does not matter up to even permutations.

13. Let  $(C_*, d_*)$  be a chain complex, and suppose it is exact at every point  $C_n$ . Such sequences are called *exact sequences*. What is the homology of  $(C_*, d_*)$ ?

14. Let  $(C_*, d_*)$  be a chain complex supported in degree  $n$ , that is,

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow C_n \rightarrow 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0.$$

What is the homology of  $(C_*, d_*)$ ?

15. Let  $(C_*, d_*)$  be a chain complex.


(a) Suppose that the differential  $d_n$  is identically zero for some  $n$ .

$$\cdots \xrightarrow{d_{n+1}} C_n \xrightarrow{d_n=0} C_{n-1} \xrightarrow{d_{n-1}} \cdots$$

Show that  $H_{n-1} = \ker(d_{n-1})$ , and  $H_n = C_n/\text{im}(d_{n+1})$ .

(b) Suppose the differential  $d_n$  is identically zero for every  $n$ . Show that  $H_n = C_n$  for every  $n$ .

16. Compute the simplicial homology of the disjoint union of  $n$  points.  
 17. Compute the simplicial homology of a 1-simplex.  
 18. Compute the simplicial homology of  $S^1$  with each of the following  $\Delta$ -complex structures, with the given orientations of the edges.



19. Compute the simplicial homology groups of the wedge  $\bigvee_k S^1$  of  $k$  circles.  
 20. Let  $X$  be a  $\Delta$ -complex, and  $C_n(X)$  its  $n^{\text{th}}$  simplicial chain group.  
 (a) Show that  $C_0(X) = \ker(\partial_0)$ , so  $C_0(X)$  is the group of 0-cycles. Conclude that, topologically, a 0-cycle is any linear combination of vertices of  $X$ .  
 (b) Show that two vertices in  $C_0(X)$  are homologous exactly if they are connected via a path of edges in  $X$ .  
 (c) Conclude that  $H_0(X)$  consists of formal sums of equivalence classes of vertices of  $X$ , where two vertices are equivalent if they are in the same path-component of  $X$ .  
 (d) Explain the sense in which  $H_0(X)$  "is" the free abelian group on the path components of  $X$ .  
 21. (a) Let  $(C_*, d_*)$  be a chain complex. Explain why, if the  $n^{\text{th}}$  homology group  $H_n$  has rank  $N$ , then the  $n^{\text{th}}$  chain group  $C_n$  must have had rank at least  $N$ .  
 (b) Let  $X$  be a space. We will show that the homology groups are homeomorphism invariants (in fact, homotopy invariants). Explain why, if the  $n^{\text{th}}$  simplicial homology group  $H_n(X)$  has rank  $N$ , then any  $\Delta$ -complex structure on  $X$  must have at least  $N$  simplices of dimension  $n$ .  
 22. Review the structure theorem for finitely generated abelian groups.

## Assignment questions

(Hand these questions in!)

- For each of the following spaces, define a  $\Delta$ -complex structure on the space, and compute its simplicial homology groups.
  - a 2-simplex
  - $S^2$
  - a torus
  - a Mobius band

- Definition (Morphism of chain complexes).** A morphism  $f_*$  of chain complexes or chain map from  $(C_*, \partial_*)$  to  $(D_*, \delta_*)$  is a sequence of group homomorphisms  $f_n : C_n \rightarrow D_n$  making the following diagram commute.

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & C_{n+1} & \xrightarrow{\partial_{n+1}} & C_n & \xrightarrow{\partial_n} & C_{n-1} & \xrightarrow{\partial_{n-1}} & \dots \\
 & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} & & \\
 \dots & \longrightarrow & D_{n+1} & \xrightarrow{\delta_{n+1}} & D_n & \xrightarrow{\delta_n} & D_{n-1} & \xrightarrow{\delta_{n-1}} & \dots
 \end{array}$$

- Verify that a morphism  $f_*$  of chain complexes induces well-defined group homomorphisms on the homology groups

$$f_n : H_n(C_*) \rightarrow H_n(D_*)$$

for every  $n$ .

- Definition (Quasi-isomorphism).** A morphism of chain complexes  $f_* : (C_*, \partial_*) \rightarrow (D_*, \delta_*)$  is a *quasi-isomorphism* if the maps induced on homology are all isomorphisms.

Give an example of a quasi-isomorphism of chain complexes where at least one map  $f_n$  is not an isomorphism.

- (Homomorphisms of free abelian groups).** Let  $A$  be an  $n \times n$  integer matrix, viewed as  $\mathbb{Z}$ -linear map from  $\mathbb{Z}^n$  to  $\mathbb{Z}^n$ .
  - Suppose that  $A$  has rank  $n$ . Prove that the kernel of  $A$  is trivial. (Note: Here we mean 'rank' in the usual sense from linear algebra, for example, it is the rank of  $A$  when  $A$  is viewed as a matrix with entries in  $\mathbb{Q}$ ).
  - Show by example that, even if  $A$  has rank  $n$ , it need not be surjective.
  - The *cokernel* of a map of abelian groups is the quotient of its codomain by its image. Prove or find a counterexample: if the map  $A$  has rank  $n$ , then the cokernel of  $A$  must be finite.

## Wellbeing

(This section is completely optional. This is a nudge to prioritize your wellbeing during the pandemic.)

- (Health comes first).** Make your health and happiness goals a priority this week.
- (Cardio exercise).** This week, if you're able, get your heart rate up! Start at an exercise intensity appropriate for your current fitness level, and be mindful of your knees and joints. Go outside if you can. Even if you cannot, consider jumping rope, jumping jacks, push-ups, burpees, squats, high-knees, moving planks, lunges – or other exercise that requires minimal space or equipment. Or, find a workout video that suits your level and taste, like interval training, kickboxing, or dance.

Aerobic exercise has benefits for physical health, immune function, cognitive performance, and mood.