

# 1 Simplicial Group Actions

These notes follow Bredon “Compact transformation groups” Chapter 3.1.

**Definition I.** Let  $X$  be an (abstract) simplicial complex and  $G$  a group. An action of  $G$  on  $X$  is called a *simplicial group action* if each element  $g \in G$  acts by a simplicial map  $g : X \rightarrow X$ .

Our goal is to make sense of the quotient of a simplicial group action.

**Exercise 1. (Bonus)** Let  $X$  be an abstract simplicial complex with a simplicial action by a group  $G$ . Show that the action of  $G$  induces an action of  $G$  on the barycentric subdivision of  $\text{sd}(X)$  of  $X$ , and that these actions coincide on the geometric realizations under the natural homeomorphism  $|X| \cong |\text{sd}(X)|$ .

**Definition II.** Let  $X$  be an abstract simplicial complex with vertex set  $V(X)$  and simplices  $S(X)$ . Suppose  $X$  admits a simplicial action by a group  $G$ . Define the quotient to be the abstract simplicial complex  $X/G$  defined as follows. A vertex  $v^* \in V(X/G)$  is a  $G$ -orbit of a vertices in  $V(X)/G$ . A collection of vertices  $\{v_0^*, \dots, v_p^*\}$  span a simplex in  $S(X/G)$  if at least one choice of representatives  $\{v_0, \dots, v_p\}$  of the orbits span a simplex in  $S(X)$ . The simplex  $\{v_0, \dots, v_p\}$  of  $X$  is called a simplex *over* the simplex  $\{v_0^*, \dots, v_p^*\}$  of  $X/G$ .

**Exercise 2. (Bonus)** Let  $X$  be an abstract simplicial complex with a simplicial action by a group  $G$ . Verify that the abstract simplicial structure defined on  $X/G$  is in fact a valid abstract simplicial structure.

There is a natural simplicial map

$$\begin{aligned} X &\longrightarrow X/G \\ v &\longmapsto v^* \end{aligned}$$

that gives rise to a simplicial map  $|X| \rightarrow |X/G|$ . This map is  $G$ -equivariant with respect to the trivial  $G$ -action on  $|X/G|$ , therefore the map factors through the orbit space  $|X|/G$  via continuous maps

$$\begin{array}{ccc} |X| & & \\ \downarrow & \searrow & \\ |X|/G & \dashrightarrow & |X/G| \end{array}$$

For a simplex  $\{v_0, \dots, v_p\}$  of  $X$ , and a point  $(\sum_{i=0}^p t_i v_i)$  in the simplex, consider the  $G$ -orbit of this point in  $|X|/G$ . The map  $|X|/G \rightarrow |X/G|$  is defined as follows.

$$\begin{aligned} |X|/G &\longrightarrow |X/G| \\ G \cdot \left( \sum_{i=0}^p t_i v_i \right) &\longmapsto \sum_{i=0}^p t_i (G \cdot v_i) = \sum_{i=0}^p t_i v_i^* \end{aligned}$$

**Exercise 3. (Bonus)**

(a) Verify that the map  $X \rightarrow X/G$  is simplicial, and  $G$ -equivariant with respect to the trivial action of  $G$  on  $X/G$ .

(b) Verify that the induced map  $|X| \rightarrow |X/G|$  factors through  $|X|/G$ , and the factorization has the formula claimed.

Unfortunately, in general, this map  $|X|/G \rightarrow |X/G|$  need not be a homeomorphism, or a homotopy equivalence, as we see in the following exercise.

**Exercise 4.** Consider the simplicial set  $R$  with vertices  $V(R) = \{n\}_{n \in \mathbb{Z}}$ , and simplices  $S(R)$  consisting of the vertices and all edges of the form  $\{n, n + 1\}$ . We can identify its geometric realization with the simplicial structure on the real line  $\mathbb{R}$  that has a vertex at each integer point.

- (a) Consider the group  $G = \mathbb{Z}$ ; for notational clarity we denote its elements  $g_m$  for  $m \in \mathbb{Z}$ . Consider the action of  $\mathbb{Z}$  on  $R$  by

$$\begin{aligned} g_m : V(R) &\longrightarrow V(R) \\ g_m \cdot n &\longmapsto m + n \end{aligned}$$

Verify that these are well-defined simplicial maps, and describe the action on the geometric realization  $|R|$ .

- (b) Describe the simplicial complex  $R/G$ . Describe the spaces  $|R/G|$  and  $|R|/G$ , and the map  $|R|/G \rightarrow |R/G|$ .
- (c) Now consider the barycentric subdivision of  $\text{sd}(R)$  of  $R$ , with the induced action of  $G$ . Is  $\text{sd}(R)/G$  isomorphic to  $R/G$ ? Again compare the spaces  $|\text{sd}(R)/G|$  and  $|\text{sd}(R)|/G$ .
- (d) Barycentrically subdivide a second time. Compare  $|\text{sd}^2(R)/G|$  and  $|\text{sd}^2(R)|/G$ .

**Definition III.** Let  $X$  be an abstract simplicial complex with a simplicial action of a group  $G$ . The action is called *regular* if the group  $G$  and all of its subgroups  $H \subseteq G$  satisfy the following Condition  $(\mathcal{R})$ :

- $(\mathcal{R})$  Suppose  $h_0, h_1, \dots, h_p$  are elements of  $H$ . Suppose that  $(v_0, v_1, \dots, v_p)$  and  $(h_0 \cdot v_0, h_1 \cdot v_1, \dots, h_p \cdot v_p)$  are  $p$ -tuples of (not necessarily distinct) vertices of  $X$  such that  $\{v_0, v_1, \dots, v_p\}$  and  $\{h_0 \cdot v_0, h_1 \cdot v_1, \dots, h_p \cdot v_p\}$  are simplices of  $X$ . Then there exists some  $h \in H$  such that  $h \cdot v_i = h_i \cdot v_i$  for all  $i$ .

In this case, we call  $X$  a *regular  $G$ -complex*.

**Theorem IV.** Let  $X$  is a  $G$ -complex and  $H \subseteq G$  a subgroup, and  $X/H$  the quotient complex. Then  $|X|/H$  has a natural simplicial structure, such that the map  $|X|/H \rightarrow |X/H|$  is a simplicial isomorphism.

**Exercise 5.** This exercise will establish Theorem IV.

- (a) Suppose that  $X$  is a regular  $G$ -complex, and let  $\sigma = \{v_0^*, \dots, v_p^*\}$  be a simplex of  $X/G$ . Show that the set of simplices of  $X$  over  $\sigma$  form a  $G$ -orbit of  $p$ -simplices.
- (b) Verify that the map  $|X|/G \rightarrow |X/G|$  is one-to-one and onto.
- (c) **(Bonus)** Verify that the topology on  $|X|/G$  agrees with the weak topology on  $|X/G|$ .

We will prove that, if a group  $G$  acts simplicially on an abstract simplicial complex  $X$ , then the induced action of  $G$  of the two-fold barycentric subdivision  $\text{sd}^2(X)$  of  $X$  is a regular action. To prove this, we introduce a weaker Condition  $(\mathcal{S})$  on a group action,

- $(\mathcal{S})$  For any  $g \in G$  and simplex  $\sigma$  of  $X$ , the element  $g$  fixes the intersection  $\sigma \cap (g \cdot \sigma)$  pointwise.

Equivalently,

- $(\mathcal{S}')$  For any  $g \in G$ , if vertices  $v$  and  $g \cdot v$  are contained in the same simplex, then  $v = (g \cdot v)$ .

- Exercise 6.** (a) Verify that Condition  $(\mathcal{R})$  implies Condition  $(\mathcal{S}')$ .  
(b) Verify that Conditions  $(\mathcal{S})$  and  $(\mathcal{S}')$  are equivalent.

**Theorem V.** *Let  $X$  be an abstract simplicial complex, and  $G$  a group.*

- *If  $G$  acts simplicially on  $X$ , then the induced action of  $G$  on the barycentric subdivision  $\text{sd}(X)$  of  $X$  satisfies Condition  $(\mathcal{S})$ .*
- *If  $G$  acts on  $X$  by a simplicial action satisfying Condition  $(\mathcal{S})$ , then the induced action of  $G$  on the barycentric subdivision  $\text{sd}(X)$  of  $X$  satisfies Condition  $(\mathcal{R})$ .*

*In particular, if  $G$  acts simplicially on  $X$ , then the induced action of  $G$  on the twice-iterated barycentric subdivision  $\text{sd}^2(X)$  of  $X$  is a regular group action in the sense of Definition III.*

**Exercise 7.** Prove Theorem V.

**Exercise 8. (Bonus)** Suppose that a simplicial action of  $G$  on  $X$  satisfies Condition  $(\mathcal{S})$ . Show that  $|X^G| \cong |X|^G$ . Here  $X^G$  denotes the subcomplex of simplices fixed pointwise by  $G$ , and  $|X|^G$  the subspace of  $|X|$  fixed pointwise by the induced action of  $G$ .