

MASTERCLASS EXERCISES

JENNY WILSON, LECTURE 1

INTRODUCTION TO THE TITS BUILDINGS

Let \mathbb{F} be a field, and V an \mathbb{F} -vector space. Recall that the *Tits building* $\mathcal{T}(V)$ is the geometric realization of the poset of proper nonzero vector subspaces of V under inclusion. We sometimes write $\mathcal{T}_n(\mathbb{F})$ for $\mathcal{T}(\mathbb{F}^n)$.

With the following exercises, we can extend the definition of the Tits building to PIDs.

- (1) Let R be a PID, and $U \subseteq R^n$ be an R -submodule. Show that the following conditions are equivalent. If these conditions hold, we say that U is *split* or that U is a *summand* of R^n .
 - (i) There exist an R -submodule C such that $R^n = U \oplus C$.
 - (ii) There exists a basis for U that extends to a basis for R^n .
 - (iii) Any basis for U extends to a basis for R^n .
 - (iv) The quotient R^n/U is torsion-free.
- (2) Let R be a PID. An element $v = (r_1, r_2, \dots, r_n) \in R^n$ is called *unimodular* if the ideal generated by (r_1, r_2, \dots, r_n) is R . In other words, the gcd of (r_1, r_2, \dots, r_n) is a unit.
 - (a) Show that a nonzero element $v \in R^n$ spans a direct summand of R^n if and only if it is unimodular.
 - (b) Show that $v \in R^n$ is an element of a basis for R^n if and only if it is unimodular.
 - (c) Give an example of a PID R and two unimodular vectors in R^n that can never both be elements of the same basis.

For R a PID and V a free R -module, we may further define $\mathcal{T}(V)$ to be the poset of proper nonzero summands of V under inclusion.

- (3) Let R be a PID, and let U, V be summands of R^n .
 - (a) Show that $U \cap V$ is always a summand of R^n .
 - (b) Show that $U + V$ need not be a summand of R^n .
- (4) Let $U \subseteq R^n$ be a direct summand, and let $W \subseteq U$. Show that W is a summand of U if and only if it is a summand of R^n .

In the next exercises we will that this ostensible generalizations of the Tits buildings to PIDs in fact reduces to the field case.

- (5) Let R be a PID and $F(R)$ be its field of fractions.
 - (a) Show that there is a bijection between R -submodule summands of R^n and $F(R)$ -vector subspaces of $F(R)^n$, given by the following correspondence. Identify $F(R)^n$ with $R^n \otimes_R F(R)$.

$$\begin{aligned} \{\text{summands of } R^n\} &\longleftrightarrow \{\text{subspaces of } F(R)^n\} \\ U &\longmapsto U \otimes_R F(R) \\ V \cap R^n &\longleftarrow V \end{aligned}$$

- (b) Verify that this bijection induces an isomorphism of posets of submodules under inclusion.
- (c) Conclude that $\mathcal{T}_n(R)$ can be canonically identified with $\mathcal{T}_n(F(R))$.

COXETER COMPLEXES AND BUILDINGS

The Tits buildings, as the name suggests, are examples of *buildings*. In this exercise we will define a building and verify that the Tits buildings satisfy the axioms.

To define a building, we first need the notion of a Coxeter complex.

Definition (Coxeter system). A *Coxeter system* is a group W (the *Coxeter group*) along with a distinguished generating set $S = \{s_1, s_2, \dots, s_n\} \subseteq W$ such that the corresponding presentation has the form

$$W = \langle s_1, s_2, \dots, s_n \mid (s_i s_j)^{m_{i,j}} = 1 \rangle, \quad m_{i,i} = 1, \quad 2 \leq m_{i,j} \leq \infty \text{ for } i \neq j.$$

Coxeter groups are abstract generalizations of *reflection groups*. Notably for our purposes, the symmetric group S_{n+1} is a Coxeter group with generators the simple transpositions $s_i = (i \ i+1)$ and associated presentation

$$S_{n+1} = \langle s_1, s_2, \dots, s_n \mid s_i^2, (s_i s_{i+1})^3, (s_i s_j)^2 \text{ for } |i-j| > 1 \rangle.$$

Given a Coxeter system (W, S) , a *standard subgroup* of W is any subgroup W_J generated by a subset J of S . A *standard coset* is a coset wW_J for $w \in W$ and W_J a standard coset.

Definition (Coxeter complex). Given a Coxeter system (W, S) , consider the poset $P_{W,S}$ of proper standard cosets under reverse inclusion. One way to define the *Coxeter complex* $X_{W,S}$ associated to (W, S) is as follows. The p -simplices of $X_{W,S}$ are indexed by standard cosets wW_J with $|J| = n - p - 1$, and assembled in such a way that the geometric realization of $P_{W,S}$ is the barycentric subdivision of $X_{W,S}$. In other words, the poset of cells of $X_{W,S}$ under inclusion is precisely $P_{W,S}$.

- (6) (a) Sketch the Coxeter complex for the symmetric groups S_2 and S_3 .
 (b) Describe the standard cosets in the symmetric group S_{n+1} .
 (c) Show that the Coxeter complex of S_{n+1} can be identified with the flag complex of nonempty subsets of the set $\{1, 2, \dots, n, n+1\}$.
 (d) Show that the Coxeter complex of the symmetric group S_{n+1} can be identified with the boundary of a barycentrically-subdivided n -simplex. Conclude that the Coxeter complex is topologically a sphere S^{n-1} .

Definition (Building). A *building* is a simplicial complex Δ that can be written as a union of subcomplexes Σ , called *apartments*, that satisfy the following axioms.

(B0) Each apartment Σ is a Coxeter complex.

(B1) For any two simplices $A, B \in \Delta$, there is an apartment Σ containing both of them.

(B2) If Σ and Σ' are two apartments containing A and B , then there is an isomorphism $\Sigma \rightarrow \Sigma'$ fixing A and B pointwise.

Top-dimensional simplices are called *chambers*, and codimension-one simplices are *panels*.

Condition **(B2)** is equivalent to the following.

(B2') Let Σ and Σ' be two apartments containing a simplex C that is a chamber of Σ . Then there is an isomorphism $\Sigma \xrightarrow{\cong} \Sigma'$ fixing every simplex of $\Sigma \cap \Sigma'$.

Let V be a vector space over \mathbb{Q} . Recall that the *Tits buildings* $\mathcal{T}(V)$ is the geometric realization of the poset of proper nonzero vector subspaces of V .

Further recall that a *frame* for V is a decomposition $V = L_1 \oplus L_2 \oplus \dots \oplus L_n$ of V as a direct sum of 1-dimensional subspaces L_i . For each frame $L = \{L_1, L_2, \dots, L_n\}$ for V , we define an apartment A_L

to be the full subcomplex of $\mathcal{T}(V)$ on vertices corresponding to direct sums of all proper nonempty subsets of $\{L_1, L_2, \dots, L_n\}$.

In the next exercise we will verify that the Tits building $\mathcal{T}(V)$, along with the system of apartments $\{A_L \mid L \text{ a frame for } V\}$, is a building.

- (7) (a) Suppose V is n -dimensional. Show that an apartment A_L is isomorphic to the Coxeter complex associated to S_n . Conclude that axiom **(B0)** holds.
- (b) Given a flag $0 \subsetneq V_1 \subsetneq V_2 \subsetneq \dots \subsetneq V_p \subsetneq V$, let's say that frame $L = \{L_1, L_2, \dots, L_n\}$ is *compatible* with this flag if every subspace V_i is a direct sum of lines L_j . Show that, given any two flags in V , there is a frame that is compatible with both of them. Use this result to conclude that **(B1)** holds.
(Despite being “just” elementary linear algebra, this exercise is not entirely trivial! See Abramenko–Brown “Buildings” Section 4.3.)
- (c) Verify that axiom **(B2')** holds.
Hint: Given a chamber in Σ corresponding to a complete flag $0 \subsetneq V_1 \subsetneq V_2 \subsetneq \dots \subsetneq V_{n-1} \subsetneq V$, construct an explicit isomorphism to the Coxeter complex using the function

$$\begin{aligned} \phi : \Sigma &\longrightarrow \{\text{subsets of } [n]\} \\ U &\longmapsto \{i \mid \dim(U \cap V_i) < \dim(U \cap V_{i+1})\} \end{aligned}$$

Observe that this isomorphism depends only on the chamber and not on Σ .

SOLOMON–TITS

The goal of this section is to give a proof of the Solomon–Tits theorem, which states that the Tits building $\mathcal{T}_n(K)$ is homotopy equivalent to a wedge of spheres of dimension $(n - 2)$.

Fix a field K and a positive integer n . Recall that the Tits building $\mathcal{T}_n(K)$ is the geometric realization of the poset of proper nonzero subspaces of K^n under inclusion. Explicitly, $\mathcal{T}_n(K)$ is a simplicial complex defined as follows. The vertices of $\mathcal{T}_n(K)$ are proper nonzero subspaces of K^n . A collection of vertices span a simplex precisely when they form a flag.

- (8) Fix a field K . Verify that, when $n = 1$, the building $\mathcal{T}_n(K)$ is empty, and when $n = 2$, the building $\mathcal{T}_n(K)$ is a discrete set of points, that is, a wedge of 0-spheres.
- (9) Draw the Tits building for $K = \mathbb{Z}/2\mathbb{Z}$ and $n \leq 3$. Can you see explicitly that it is homotopy equivalent to a wedge of spheres?

To prove the Solomon–Tits theorem, we will use a method sometimes called “discrete Morse theory”. I first learned this proof from Bestvina’s notes “PL Morse Theory”.

Definition (Realizations and links). For a poset T , write $|T|$ for its geometric realization. For $t \in T$, we write $\text{Lk}_T(t)$ for the link of t in T ,

$$\text{Lk}_T(t) = \{s \in T \mid s < t \text{ or } s > t\}.$$

We write $\text{Lk}_T^\uparrow(t)$ for the subposet

$$\text{Lk}_T^\uparrow(t) = \{s \in T \mid s > t\}$$

and we write $\text{Lk}_T^\downarrow(t)$ for the subposet

$$\text{Lk}_T^\downarrow(t) = \{s \in T \mid s < t\}.$$

- (10) Verify that $|\text{Lk}_T(t)| = |\text{Lk}_T^\uparrow(t)| \overset{*}{\text{join}} |\text{Lk}_T^\downarrow(t)|$.

The following result is our key lemma.

Lemma (Discrete Morse Theory). Let T be a poset with $T = X_0 \cup T_1 \cup \cdots \cup T_m$ as sets. Let $X_k = X_0 \cup T_1 \cup \cdots \cup T_k$. Suppose the following:

- (i) $|X_0|$ is contractible.
- (ii) For $i \geq 1$ then any pair $s, t \in T_i$ of distinct elements are not comparable.
- (iii) For $i \geq 1$ and $t \in T_i$,

$$|\mathrm{Lk}_T(t) \cap X_{i-1}| \simeq \bigvee S^{d-1} \quad \text{or} \quad |\mathrm{Lk}_T(t) \cap X_{i-1}| \simeq *.$$

Then $|T|$ is $(d-1)$ -connected. In particular, if $|\mathrm{Lk}_T(t) \cap X_{i-1}| \simeq \bigvee S^{d-1}$ for at least one i and t , then $|T|$ is homotopy equivalent to a wedge of d -spheres. Otherwise, $|T|$ is contractible.

- (11) Prove the Discrete Morse Theory lemma.
- (12) Fix a field K and $n \geq 2$. Let T be the poset of nonzero proper subspaces of K^n , so $\mathcal{T}_n(K)$ is defined to be $|T|$. Assume by induction that $\mathcal{T}_m(K) \simeq \bigvee S^{m-2}$ for all $m < n$; we proved the base case in Exercise (8). Fix a line L in K^n .
 - Let X_0 be the subposet of T on vertices V such that $L \subseteq V$.
 - For $i = 1, \dots, n-1$, let T_i be the set of subspaces V of K^n

$$T_i = \{V \subseteq K^n \mid \dim(V) = i, L \not\subseteq V\}$$

- (a) Verify that $|X_0| \subseteq |T|$ is the star on the vertex L , and hence contractible.
- (b) Verify that for fixed i , distinct elements in T_i are not comparable.
- (c) Suppose $1 \leq i \leq (n-2)$. Show that, for $V \in T_i$, the subspace $(V+L)$ is a cone point of $|\mathrm{Lk}_T(V) \cap X_{i-1}|$. (Why did we need the assumption $i \leq (n-2)$?)
- (d) Verify that, for $i = n-1$ and $V \in T_i$,

$$|\mathrm{Lk}_T(V) \cap X_{i-1}| \simeq \bigvee S^{n-3}.$$

Hint: Compare $|\mathrm{Lk}_T(V) \cap X_{i-1}|$ to $\mathcal{T}_{n-1}(K)$.

- (e) Use the Discrete Morse Theory lemma to conclude that $\mathcal{T}_n(K) \simeq \bigvee S^{n-2}$.

There are other elegant approaches to computing the homotopy type of $\mathcal{T}_n(K)$. For example, see Abramenko–Brown “Buildings” Section 4.12 for an approach using the theory of shellability. In principle, these proofs can also be used to describe a generating sets for the reduced homology of $\mathcal{T}_n(K)$.

THE SHARBLY RESOLUTION

Let R be a PID, and let $\mathrm{St}_n(R)$ be the associated Steinberg module,

$$\mathrm{St}_n(R) := \tilde{H}_{n-2}(\mathcal{T}_n(R); \mathbb{Z}).$$

By Exercise 5, we can identify $\mathcal{T}_n(R)$ with $\mathcal{T}_n(\mathbb{F})$, where \mathbb{F} is the field of fractions of R .

In this section we will construct a resolution of the Steinberg module due to Lee–Szczerba (1976), which has since been named the *Sharbly resolution*. To quote Lee–Szczerba (Section 4):

In theory, one should be able to use [the Sharbly resolution] to compute the groups $H_q(\mathrm{SL}_n(R); \mathrm{St}_n(R))$ for $q \geq 0$. However, because of the size of [the terms of the resolution], this is impractical except when $q = 0$.

The proof of the Sharbly resolution is both a significant historical development, and also an instructive application of some of the techniques of the field. The construction will use the *Acyclic Covering*

Lemma (See Brown “Cohomology of Groups” Section VII Lemma 4.4). Let X be a CW complex, and suppose X is the union of a family of nonempty subcomplexes

$$X = \bigcup_{\alpha \in J} X_\alpha.$$

The *nerve* N of the family X_α is the abstract simplicial complex with vertex set J and such that a finite subset $\sigma \subseteq J$ spans a simplex if and only if the intersection $X_\sigma = \bigcap_{\alpha \in \sigma} X_\alpha$ is nonempty.

Lemma (Acyclic Covering Lemma). Suppose a CW complex X is a union of subcomplexes X_α such that every non-empty intersection $X_{\alpha_0} \cap X_{\alpha_1} \cap \dots \cap X_{\alpha_p}$ is acyclic. Then $H_*(X) \cong H_*(N)$, where N is the nerve of the cover.

- (13) **(Bonus).** Use the Mayer–Vietoris spectral sequence to prove the Acyclic Covering Lemma. See Brown “Cohomology of Groups” Section VII.4.

Let R be a PID. Recall that an element $v = (r_1, r_2, \dots, r_n) \in R^n$ is called *unimodular* if the ideal generated by (r_1, r_2, \dots, r_n) is R .

$$\mathcal{S}_q = \mathcal{S}_q(R^n) = \{(n+q) \times n \text{ matrices } A = (a_{i,j}) \text{ over } R \mid (a_{i,1}, \dots, a_{i,n}) \text{ is unimodular for all } i\}$$

$$\mathcal{P}_q = \mathcal{P}_q(R^n) = \{A \in \mathcal{S}_q \mid \text{each } n \times n \text{ submatrix has determinant } 0\}$$

These sets have actions of $\text{GL}_n(R)$ by right multiplication. Let $C(\mathcal{S}_q)$ and $C(\mathcal{P}_q)$ denote the free abelian groups on \mathcal{S}_q and \mathcal{P}_q , respectively, and let $C_q(R^n) = C(\mathcal{S}_q)/C(\mathcal{P}_q)$ be the quotient; it is a right $\mathbb{Z}[\text{GL}_n(R)]$ -module.

Our goal is to prove the following, Lee–Szczarba Theorem 3.1.

Theorem (The Sharbly resolution). There is an epimorphism $\phi : C_0(R^n) \rightarrow \text{St}_n(R)$ of right $\mathbb{Z}[\text{GL}_n(R)]$ -modules so that

$$\longrightarrow C_q(R^n) \longrightarrow C_{q-1}(R^n) \longrightarrow \dots \longrightarrow C_0(R^n) \xrightarrow{\phi} \text{St}_n(R) \longrightarrow 0$$

is a free resolution of $\text{St}_n(R)$ by $\mathbb{Z}[\text{GL}_n(R)]$ -modules.

- (14) (a) Let K be the simplicial complex whose vertices are the unimodular elements of R^n , and whose simplices are all finite nonempty subsets of vertices. Show that K is contractible.
 (b) Use the long exact sequence of a pair to show that, for a subcomplex $L \subseteq K$,

$$H_q(K, L) \cong \tilde{H}_{q-1}(L).$$

- (c) Let $L \subseteq K$ be the subcomplex consisting of all simplices with the property that all of their vertices lie in a proper direct summand of R^n . Let $\{H_i \mid i \in I\}$ be the set of direct summands of R^n of rank $(n-1)$. Let K_i be the full subcomplex of L with vertices lying in H_i . Show that $\{K_i \mid i \in I\}$ is an acyclic covering of L in the sense of the Acyclic Covering Lemma.

Hint: First argue that, since R is a PID, a nonempty intersection of summands H_i must be a direct summand of R^n isomorphic to R^r for some $0 < r < n$.

- (d) Let N be the nerve of the cover $\{K_i \mid i \in I\}$. Use the Acyclic Covering Lemma to deduce that $H_q(L) \cong H_q(N)$ for all $q \geq 0$.
 (e) Recall that \mathbb{F} denotes the field of fractions of R . Let $\{W_j \mid j \in J\}$ denote the set of hyperplanes in \mathbb{F}^n . Let T_j be the subcomplex of the Tits building $\mathcal{T}_n(\mathbb{F})$ consisting of all simplices with W_j as a vertex. Show that $\{T_j \mid j \in J\}$ is an acyclic covering of L in the sense of the Acyclic Covering Lemma.
 (f) Let \tilde{N} be the nerve of the cover $\{T_j \mid j \in J\}$. Use the Acyclic Covering Lemma to deduce that $H_q(\mathcal{T}_n(\mathbb{F})) \cong H_q(\tilde{N})$ for all $q \geq 0$.

(g) Show that the mapping

$$\begin{aligned} \{\text{summands of } R^n\} &\longrightarrow \{\text{subspaces of } \mathbb{F}^n \cong R^n \otimes_R \mathbb{F}\} \\ H &\longmapsto H \otimes_R \mathbb{F} \end{aligned}$$

defines a simplicial isomorphism of N onto \tilde{N} .

(h) We have proved the following isomorphisms. Verify that they are $\text{GL}_n(R)$ -equivariant.

$$\begin{aligned} H_q(K, L) &\cong \tilde{H}_{q-1}(L) \\ &\cong \tilde{H}_{q-1}(N) \\ &\cong \tilde{H}_{q-1}(\tilde{N}) \\ &\cong \tilde{H}_{q-1}(\mathcal{T}_n(\mathbb{F})) \end{aligned}$$

(i) Verify that the $(n-2)$ -skeleton of L coincides with the $(n-2)$ -skeleton of K , so

$$C_q(K, L) = 0 \quad \text{for } q \leq n-2.$$

(j) Using the Solomon–Tits result that $\mathcal{T}_n(\mathbb{F}) \simeq \bigvee S^{n-2}$, deduce that there is an exact sequence

$$\cdots \longrightarrow C_{q+n}(K, L) \longrightarrow C_{q+n-1}(K, L) \longrightarrow \cdots \longrightarrow C_{n-1}(K, L) \longrightarrow \text{St}_n(R) \longrightarrow 0.$$

(k) Show that there are $\text{GL}_n(R)$ -equivariant isomorphisms of chain complexes

$$\begin{aligned} C_{q+n-1}(K) &\cong C(\mathcal{S}_q) \\ C_{q+n-1}(L) &\cong C(\mathcal{P}_q) \\ C_{q+n-1}(K, L) &\cong C_q(R^n) \end{aligned}$$

(l) Conclude the existence of the Sharbly resolution.

(m) Show that differentials in the Sharbly resolution are induced by the map

$$\begin{aligned} \mathcal{S}_q &\longrightarrow \mathcal{S}_{q-1} \\ A &\longmapsto \sum (-1)^i d_i(A) \end{aligned}$$

where d_i is the map that deletes the i^{th} row of the matrix A .

(15) Prove that the Sharbly resolution is free.

Hint: First verify that $\text{GL}_n(R)$ acts freely on the set of $(n+q) \times n$ matrices A satisfying

- each row of A is unimodular,
- some $(n \times n)$ -submatrix of A has non-vanishing determinant.

(16) Give a geometric interpretation of the map $\phi : C_0(R^n) \rightarrow \text{St}_n(R)$. Conclude that the Steinberg module is generated by apartment classes.

Let R be a Euclidean ring with a multiplicative Euclidean norm. Using the Sharbly resolution, Lee–Szczarba went on to prove that the $\text{SL}_n(R)$ -coinvariants of $C_0(R^n)$ vanish, which implies that

$$H_0(\text{SL}_n(R); \text{St}_n(R)) = 0.$$

By virtual Bieri–Eckmann duality, this then implies that the rational cohomology of $\text{SL}_n(R)$ vanishes in its virtual cohomological dimension.

(17) **(Bonus).** Let R be a Euclidean ring with a multiplicative Euclidean norm. Prove directly that $\text{SL}_n(R)$ -coinvariants of $C_0(R^n)$ vanish. See Lee–Szczarba Theorem 4.1.

ASH–RUDOLPH

Let $\mathcal{T}_n(\mathbb{Q})$ be the Tits building on the rational vector space \mathbb{Q}^n . Let A_L denote the apartment associated to a frame $L = \{L_1, L_2, \dots, L_n\}$ for \mathbb{Q}^n . Recall the following.

Definition (Integral apartment). Let $L = \{L_1, L_2, \dots, L_n\}$. The frame L (and the apartment A_L) are called *integral* if

$$(L_1 \cap \mathbb{Z}^n) \oplus (L_2 \cap \mathbb{Z}^n) \oplus \dots \oplus (L_n \cap \mathbb{Z}^n) = \mathbb{Z}^n.$$

(18) Consider \mathbb{Q}^2 . Verify that the apartment corresponding to the frame

$$\left\{ \mathbb{Q} \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbb{Q} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

is integral, but the apartment corresponding to the frame

$$\left\{ \mathbb{Q} \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \mathbb{Q} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

is not integral.

(19) Develop a determinant condition for verifying whether or not a frame is integral.

Let $L = \{L_1, L_2, \dots, L_n\}$ be a frame, and let A_L be the associated apartment in the Tits building. Recall from Exercises (6) and (7) that A_L is an $(n-2)$ -sphere, specifically, it is simplicially isomorphic to the barycentric subdivision of the boundary of an $(n-1)$ -simplex.

- (20) (a) A permutation $\sigma \in S_n$ acts on L by $L_i \mapsto L_{\sigma(i)}$. Show that σ induces a simplicial isomorphism $A_L \rightarrow A_L$. Show that this isomorphism is orientation-preserving if σ is even, and orientation-reversing if σ is odd.
- (b) The apartment A_L represents a homology class $[A_L] \in H_{n-2}(\mathcal{T}_n(\mathbb{Q}))$. Explain why, to make the sign of $[A_L]$ well-defined, we must order the frame L up to sign. Moreover, the symmetric group $S_n \subseteq \mathrm{GL}_n(\mathbb{Z})$ acts on $[A_L]$ by sign. (Some sources leave L unordered and $[A_L]$ only defined up to sign).

Ash–Rudolph (1979) proved the following.

Theorem (Ash–Rudolph). The homology group $H_{n-2}(\mathcal{T}_n(\mathbb{Q}))$ is generated by integral apartment classes

$$\left\{ [A_L] \mid \begin{array}{l} L \text{ an integral frame for } \mathbb{Q}^n, \\ \text{ordered up to sign} \end{array} \right\}.$$

(21) (a) Find an element of $\mathrm{SL}_n(\mathbb{Z})$ that interchanges the two lines

$$\left\{ \mathbb{Q} \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbb{Q} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}.$$

(b) Show that there is **no** matrix in $\mathrm{SL}_n(\mathbb{Z})$ that interchanges the two lines

$$\left\{ \mathbb{Q} \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \mathbb{Q} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}.$$

(c) Explain why this non-existence result was an obstacle in our first lecture to computing $H_0(\mathrm{SL}_n(\mathbb{Z}); \mathrm{St}_n(\mathbb{Q}) \otimes_{\mathbb{Z}} \mathbb{Q})$, and explain why it is resolved by Ash–Rudolph.

(22) **(Bonus)**. Again consider \mathbb{Q}^2 . Write the apartment class corresponding to the frame

$$\left\{ \mathbb{Q} \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \mathbb{Q} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

as a linear combination of integral apartment classes.

This theorem of Ash–Rudolph offers a simpler proof of the earlier theorem of Lee–Szczarba.

(23) (a) Let C_0 be the free abelian group on the symbols $[L]$ with $L = \{L_0, L_1, \dots, L_n\}$ a frame for R^n , L ordered up to sign, subject to the relation

$$\sigma \cdot [L] = (-1)^{\text{sign}(\sigma)} [L] \quad \text{for } \sigma \in S_n.$$

Show that the $\text{SL}_n(\mathbb{Z})$ -coinvariants of C_0 vanish.

(b) Deduce that $H_0(\text{SL}_n(\mathbb{Z}); \text{St}_n(\mathbb{Q})) = 0$.

(c) Deduce that $H^{vcd}(\text{SL}_n(\mathbb{Z}); \mathbb{Q}) = 0$.

JENNY WILSON, LECTURE 2

SIMPLICIAL METHODS AND A LEMMA OF QUILLEN

Recall the definition of the link of a simplex in a simplicial complex.

Definition (Link). Let X be a simplicial complex, and let $\sigma = [s_0, \dots, s_p]$ be a simplex in X . The *link* $\text{Link}_X(\sigma)$ of σ is the subcomplex of X of simplices

$$\{[t_0, \dots, t_q] \mid [s_0, \dots, s_p, t_0, \dots, t_q] \text{ is a simplex of } X\}.$$

(24) **(Bonus)**. What can you say about links of simplices in the Tits buildings $\mathcal{T}_n(\mathbb{F})$?

The following is Quillen’s definition of a Cohen–Macaulay simplicial complex. We caution that this definition is stronger than other definitions of Cohen–Macaulay appearing in the literature.

Definition (CM complex). A d -dimensional simplicial complex X is *Cohen–Macaulay (CM)* if

(i) X is $(d - 1)$ -connected,

(ii) $\text{Link}_X(\sigma)$ is $(d - 2 - \dim(\sigma))$ -connected for every simplex σ in X .

(We may simplify the statement of the definition if we consider the empty set a (-1) -simplex, and X its link).

(25) Verify that the following simplicial complexes are CM.

(a) an n -simplex, and its barycentric subdivision

(b) the boundary of an n -simplex, and its barycentric subdivision

(c) **(Bonus)** the join of CM simplicial complexes

In Church–Ellenberg–Farb’s proof of Ash–Rudolph’s theorem, we used the following lemma. The lemma follows from Quillen’s paper “Homotopy properties of the poset of nontrivial p -subgroups of a group,” Theorem 9.1 and Corollary 9.7.

For a poset X , let $|X|$ denote its geometric realization. A map $f : X \rightarrow Y$ of posets is *strictly increasing* if $f(x_1) > f(x_2)$ for all $x_1 > x_2$.

Lemma (Quillen) Let $f : X \rightarrow Y$ be a strictly increasing map of posets. Assume that $|Y|$ is a d -dimensional CM complex. For $y \in Y$, let f_y denote the subposet

$$f_y = \{x \in X \mid f(x) \leq y\} \subseteq X$$

and further assume that $|f_y|$ is CM for all y . Then $|X|$ is CM, and $f_* : \tilde{H}_d(|X|) \rightarrow \tilde{H}_d(|Y|)$ surjects.

(26) **(Bonus)**. Prove Quillen’s lemma. *Hint*: See Quillen’s paper .

MAAZEN’S THEOREM FOR $n = 2$

Definition (Complex of partial bases). Fix a PID R . Let $B_n(R)$ denote the *complex of partial bases* in R^n . Its vertices are primitive elements v_0 of R^n , and vertices $\{v_0, \dots, v_p\}$ span a p -simplex precisely when they are a subset of a basis for R^n (possibly equal to a basis of R^n).

(27) Fix a PID R . Let $P_n(R)$ denote the poset of partial bases of R^n under inclusion. Show that the geometric realization $|P_n(R)|$ is equal to the barycentric subdivision of $B_n(R)$.

The goal of this section is to prove the following result of Maazen in the case $n = 2$.

Theorem (Maazen). Let R be a Euclidean domain. Let $P_n(R)$ denote the poset of partial bases of R^n under inclusion. Then $|P_n(R)|$ is Cohen–Macaulay.

(28) Fix a Euclidean ring R with norm $|\cdot|$, and let $n = 2$.

(a) Explain why, to prove Maazen’s theorem in the case $n = 2$, it suffices to show that the graph $|P_2(R)|$ is connected, equivalently, that $B_2(R)$ is connected. Thus given a vertex indexed by a primitive vector $\begin{bmatrix} a \\ b \end{bmatrix}$ in R^2 , it suffices to find a path to the vertex $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$.

(b) Prove the following claim: Given a primitive vector $\begin{bmatrix} a \\ b \end{bmatrix}$ in R^2 with $|b| > 0$, there is a basis

$$\left\{ \begin{bmatrix} a \\ b \end{bmatrix}, \begin{bmatrix} c \\ d \end{bmatrix} \right\}$$

of R^2 with $|d| < |b|$.

Hint: First choose any partial basis $\left\{ \begin{bmatrix} a \\ b \end{bmatrix}, \begin{bmatrix} c' \\ d' \end{bmatrix} \right\}$, and consider elements $\begin{bmatrix} c \\ d \end{bmatrix}$ of the form

$$\begin{bmatrix} c' \\ d' \end{bmatrix} - q \begin{bmatrix} a \\ b \end{bmatrix}, \quad q \in R.$$

(c) Explain why the claim completes the proof.

(29) **(Bonus)**. Can you generalize this proof strategy to $n \geq 2$?

A PROOF OF ASH–RUDOLPH

The objective of the second lecture was a simplified proof of Ash–Rudolph’s theorem following Church–Farb–Putman.

Theorem (Ash–Rudolph). Let R be a Euclidean ring and \mathbb{F} its field of fractions. Then the top homology of the associated Tits building $\tilde{H}_{n-2}(\mathcal{T}_n(\mathbb{F}))$ is generated by *integral* apartment classes.

(30) Review the main argument of the lecture: use Quillen’s lemma and Maazen’s theorem to prove Ash–Rudolph’s theorem.