

**Math 471    Fall 2009    Homework 1    due: Mon Sept 21**

0. (optional) Give a brief description of your academic background and research interests. If you work in a lab or research group, please give your supervisor's name and describe your project. One paragraph is fine.

**section 1.3, floating point number systems**

1. (a) Convert  $(2009)_{10}$  to base 2.      (b) Convert  $(110110.001)_2$  to base 10.
2. The floating point representation of a real number has the form  $\pm(0.d_1d_2\dots d_n)_\beta \cdot \beta^e$ , where  $d_1 \neq 0$  and  $-M \leq e \leq M$ . Consider a system with  $\beta = 2$ ,  $n = 4$ , and  $M = 5$ .
  - (a) Find the largest and smallest positive numbers that can be represented in this system. Give your answers in decimal form.
  - (b) Find the floating point number in this system closest to  $\sqrt{2}$ .
3. page 40, problem 6(a)

**section 1.4, finite precision arithmetic**

4. page 52, problem 13
5. Consider the equation  $x^2 + 25x + 0.1 = 0$ .
  - (a) Solve for the roots using the quadratic formula,  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ , using 4-digit arithmetic (as in the example from class). Compare with the values obtained using Matlab.
  - (b) Repeat using the alternative form of the quadratic formula,  $x = \frac{2c}{-b \pm \sqrt{b^2 - 4ac}}$ . Compare with part (a) and explain the results.
6. Let  $f(x)$  be a given function and recall the forward difference approximation of  $f'(x)$ ,

$$D_+f(x) = \frac{f(x+h) - f(x)}{h},$$

where  $h > 0$  is the step size.

- (a) Take  $f(x) = \sin x$ ,  $x = \pi/4$ ,  $h = 2^{-n}$  for  $n = 1, 2, \dots, 6$ . Following the example in class, plot the error versus  $h$  and make a table with the following format: column 1:  $h$ , column 2:  $D_+f(x)$ , column 3:  $f'(x) - D_+f(x)$ , column 4:  $(f'(x) - D_+f(x))/h$ , column 5:  $(f'(x) - D_+f(x))/h^2$ . You may modify the Matlab code given in class. Present at least eight decimal digits (type "format long" in Matlab to get the full 15 digits).
- (b) Repeat for the centered difference approximation,

$$D_0f(x) = \frac{f(x+h) - f(x-h)}{2h},$$

which also approximates  $f'(x)$ . Which approximation is more accurate? Explain why.