

## SYMMETRIC BILINEAR FORMS

Let  $k$  be a field, let  $V$  be a finite dimensional vector space over  $k$  and let  $B : V \times V \rightarrow k$  be a  $k$ -bilinear pairing. Recall that, given a basis  $v_1, v_2, \dots, v_n$  of  $V$ , we encode  $B$  in a Gram matrix  $G$  with  $G_{ij} = B(v_i, v_j)$ , and that  $B$  is symmetric if and only if  $G$  is. Changing bases of  $V$  modifies the Gram matrix by  $G \mapsto S G S^T$  for invertible  $S$ . It is natural to ask how nice we can make the matrix  $G$  by action of this kind.

**To simplify our results, assume that  $k$  does not have characteristic 2.**

- (190) Suppose that  $B$  is a symmetric bilinear form. Show that there is a basis of  $V$  for which the Gram matrix of  $B$  is diagonal. (Hint: If  $B \neq 0$ , use Problem 185 to find a vector  $v$  with  $B(v, v) \neq 0$ , then consider the decomposition  $V = kv \oplus (kv)^\perp$ .)
- (191) Let  $G$  be a symmetric matrix with entries in  $k$ . Show that there is an invertible matrix  $S$  and a diagonal matrix  $D$  such that  $G = S D S^T$ .
- (192) Let  $k = \mathbb{Q}$ . Carry out the procedure in the previous problems for
- (a)  $G = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .
  - (b)  $G = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$ .

This immediately raises the question, given two diagonal matrices  $\text{diag}(\alpha_1, \alpha_2, \dots, \alpha_n)$  and  $\text{diag}(\beta_1, \beta_2, \dots, \beta_n)$ , when are the bilinear forms  $\vec{x}^T \text{diag}(\alpha_1, \alpha_2, \dots, \alpha_n) \vec{y}$  and  $\vec{x}^T \text{diag}(\beta_1, \beta_2, \dots, \beta_n) \vec{y}$  equivalent up to a change of basis? For a general field, this is a very hard question. However, we can say some things.

- (193) Suppose that there are nonzero scalars  $\gamma_i$  in  $k$  with  $\alpha_i = \gamma_i^2 \beta_i$ . Show that the  $\vec{x}^T \text{diag}(\alpha_1, \alpha_2, \dots, \alpha_n) \vec{y}$  and  $\vec{x}^T \text{diag}(\beta_1, \beta_2, \dots, \beta_n) \vec{y}$  are equivalent.
- (194) Show that the bilinear forms  $B\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}\right) = x_1 x_2 + y_1 y_2$  and  $C\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}\right) = 5x_1 x_2 + 5y_1 y_2$  are related by a change of basis in  $\mathbb{Q}^2$ , even though 5 is not square in  $\mathbb{Q}$ .
- (195) Let  $k = \mathbb{R}$ . Show that any bilinear form over  $\mathbb{R}$  can be represented by a diagonal matrix whose entries lie in  $\{-1, 0, 1\}$ .