

4. GROUPS

Definition: A **group** G is a set with a binary operation $*$: $G \times G \rightarrow G$ obeying the properties

- (1) There is an element 1 of G such that $1 * g = g * 1 = g$ for all $g \in G$.
- (2) For all $g \in G$, there is an element g^{-1} obeying $g * g^{-1} = g^{-1} * g = 1$.
- (3) For all $g_1, g_2, g_3 \in G$, we have $(g_1 * g_2) * g_3 = g_1 * (g_2 * g_3)$.

Given a group G , a **subgroup** of G is a subset containing 1 and closed under $*$ and $g \mapsto g^{-1}$.

Depending on context, we may denote $*$ by $*$, \times , \cdot or no symbol at all, and we may denote 1 as 1 , e or Id .

Problem 4.1. Show that a group G only has one element 1 obeying the condition (1).

Problem 4.2. Let G be a group and let $g \in G$. Show that G only has one element obeying the condition (2).

Definition: Given two groups G and H , a **group homomorphism** is a map $\phi : G \rightarrow H$ obeying $\phi(g_1 * g_2) = \phi(g_1) * \phi(g_2)$. A bijective group homomorphism is called an **isomorphism** and two groups are called **isomorphic** if there is an isomorphism between them.

A group homomorphism can also be called a “map of groups” or a “group map”.


Problem 4.3. Let $\phi : G \rightarrow H$ be a group homomorphism. Show that $\phi(1) = 1$ and $\phi(g^{-1}) = \phi(g)^{-1}$.

Problem 4.4. Let $\phi : G \rightarrow H$ be a group homomorphism.

- (1) The **image** of ϕ is $\text{Im}(\phi) := \{\phi(g) : g \in G\}$. Show that $\text{Im}(\phi)$ is a subgroup of G .
- (2) The **kernel** of ϕ is $\text{Ker}(\phi) := \{g \in G : \phi(g) = 1\}$. Show that $\text{Ker}(\phi)$ is a subgroup of G .

Definition: Given two groups G and H , the **product group** is the group whose underlying set is $G \times H$, with multiplication structure $(g_1, h_1) * (g_2, h_2) = (g_1 g_2, h_1 h_2)$.

Problem 4.5. Let G and H be two groups and let π_1 and π_2 be the projections $G \times H \rightarrow G$ and $G \times H \rightarrow H$ onto the first and second factor. Show that $G \times H$ obeys the **universal property of products**, meaning that, for any group F with maps $\phi_1 : F \rightarrow G$ and $\phi_2 : F \rightarrow H$, there is a unique map $(\phi_1, \phi_2) : F \rightarrow G \times H$ such that the diagram below commutes:



We close with a few more definitions:

Definition: For $g \in G$, the **conjugacy class** of g is the set $\text{Conj}(g) := \{hgh^{-1} : h \in G\}$.

Definition: A group G is called **abelian** if $g_1 * g_2 = g_2 * g_1$ for all $g_1, g_2 \in G$.

If G is abelian, we often denote $*$ by $+$ and 1 by 0 . We **never** use these notations for a non-abelian group.