

6. NORMAL SUBGROUPS, QUOTIENT GROUPS, SHORT EXACT SEQUENCES

Problem 6.1. Let G be a group and let N be a subgroup. Show that the following are equivalent:

- (1) For all $g \in G$, we have $gNg^{-1} = N$.
- (2) N is a union of (some of the) conjugacy classes of G .
- (3) All elements of G/N have the same stabilizer, for the left action of G on G/N .
- (4) Every left coset of N in G is also a right coset.
- (5) If $g_1N = g'_1N$ and $g_2N = g'_2N$, then $g_1g_2N = g'_1g'_2N$.

Definition: A subgroup N obeying the equivalent conditions of Problem 6.1 is called a *normal subgroup* of G . We write $N \trianglelefteq G$ to indicate that N is a normal subgroup of G .

Problem 6.2. Let G be S_3 . Which of the following subgroups are normal?

- (1) The subgroup generated by (12).
- (2) The subgroup generated by (123).

Problem 6.3. Let G be a group and let N be a normal subgroup of G .

- (1) Prove or disprove: Let $\alpha : F \rightarrow G$ be a group homomorphism. Then $\alpha^{-1}(N)$ is normal in F .
- (2) Prove or disprove: Let $\beta : G \rightarrow H$ be a group homomorphism. Then $\beta(N)$ is normal in H .
- (3) At least one of the statements above is false. Find an additional hypothesis you could add to make it true.

Definition: Given a group G and a normal subgroup N , the *quotient group* G/N is the group whose underlying set is the set of cosets G/N with multiplication such that $(g_1N)(g_2N) = g_1g_2N$.

This definition makes sense by Part (4) of Problem 6.1. I won't make you check that this is a group, but do so on your own time if you have any doubt. Also, I won't make you check this, but the groups G/N and $N \setminus G$, defined in the obvious ways, are isomorphic.

Let $\phi : G \rightarrow H$ be a group homomorphism. Recall that the image and kernel of ϕ are $\text{Ker}(\phi) := \{g \in G : \phi(g) = 1\}$ and $\text{Im}(\phi) := \{\phi(g) : g \in G\}$.

Problem 6.4. Show that the kernel of ϕ is a normal subgroup of G .

Problem 6.5. Show that the "obvious" map from $G/\text{Ker}(\phi)$ to $\text{Im}(\phi)$ is an isomorphism.

We often discuss quotients using the language of short exact sequences:

Definition: A *short exact sequence* $1 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{\beta} C \rightarrow 1$ is three groups A , B and C , and two group homomorphisms $\alpha : A \rightarrow B$ and $\beta : B \rightarrow C$ such that α is injective, β is surjective, and $\text{Im}(\alpha) = \text{Ker}(\beta)$.

I will occasionally write 0 instead of 1 at one end or the other of a short exact sequence. I do this when the adjacent group (meaning A or C) is abelian and it would feel bizarre to denote the identity of that abelian group as 1.

We'll write C_n for the abelian group $\mathbb{Z}/n\mathbb{Z}$. This is called the *cyclic group* of order n .

Problem 6.6. Show that there is a short exact sequence $1 \rightarrow C_m \rightarrow C_{mn} \rightarrow C_n \rightarrow 1$.

Problem 6.7. Show that there is a short exact sequence $1 \rightarrow C_3 \rightarrow S_3 \rightarrow S_2 \rightarrow 1$.

Problem 6.8. Show that there is a short exact sequence $1 \rightarrow C_2^2 \rightarrow S_4 \rightarrow S_3 \rightarrow 1$.