

### C. SCHUR-ZASSENHAUS, THE ABELIAN CASE

The aim of the next two worksheets will be to prove:

**Theorem Schur-Zassenhaus:** Let  $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$  be a short exact sequence of finite groups where  $\text{GCD}(\#(A), \#(C)) = 1$ . Then this sequence is right split, so  $B \cong A \rtimes C$ .

This is the start of an answer to the question “how are groups assembled out of smaller groups”: When you put groups of relatively prime order together, you just get semidirect products.

Today, we’ll be proving the case where  $A$  is abelian.<sup>1</sup> Here is our main result:

**Theorem:** Let  $A$  be an abelian group,  $C$  a finite group of size  $n$ , and suppose that  $a \mapsto a^n$  is a bijection from  $A$  to  $A$ . Let  $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$  be a short exact sequence. Then this sequence is right split.

**Problem C.1.** Show that, if  $A$  is a finite abelian group and  $n$  an integer such that  $\text{GCD}(\#(A), n) = 1$ , then  $a \mapsto a^n$  is a bijection. Thus, the above Theorem does imply the Schur-Zassenhaus theorem for  $A$  abelian.

**From now on, let  $A$  be an abelian group, let  $C$  be a finite group and let  $1 \rightarrow A \rightarrow B \xrightarrow{\beta} C \rightarrow 1$  be a short exact sequence. We abbreviate  $\#(C)$  to  $n$ ; we will not introduce the hypothesis on  $a \mapsto a^n$  until later. We’ll identify  $A$  with its image in  $B$ .**

Let  $\mathcal{S}$  be the set of right inverses of  $\beta$ , meaning maps  $\sigma : C \rightarrow B$  such that  $\beta(\sigma(c)) = c$ . We emphasize that  $\sigma$  is not required to be compatible with the group multiplication in any way. Let  $B$  act on  $\mathcal{S}$  by  $(b\sigma)(c) = b\sigma(\beta(b)^{-1}c)$ .

**Problem C.2.** Check that this is an action.

Let  $\sigma_1$  and  $\sigma_2 \in \mathcal{S}$ . Set

$$d(\sigma_1, \sigma_2) = \prod_{c \in C} (\sigma_1(c)\sigma_2(c)^{-1}). \quad (*)$$

We don’t have to specify the order of the product, because every term is in  $A$ .

**Problem C.3.** Show that  $d(\sigma_1, \sigma_2)d(\sigma_2, \sigma_3) = d(\sigma_1, \sigma_3)$  and  $d(\sigma_1, \sigma_2) = d(\sigma_2, \sigma_1)^{-1}$ .

**Problem C.4.** For the action of  $B$  on  $\mathcal{S}$  described above, check that  $d(b\sigma_1, b\sigma_2) = bd(\sigma_1, \sigma_2)b^{-1}$ .

Define  $\sigma_1 \equiv \sigma_2$  if  $d(\sigma_1, \sigma_2) = 1$ .

**Problem C.5.** Check that  $\equiv$  is an equivalence relation.

Define  $\mathcal{X}$  to be the set of equivalence classes of  $\mathcal{S}$  module the relation  $\equiv$ .

**Problem C.6.** Check that the action of  $B$  on  $\mathcal{S}$  descends to an action of  $B$  on  $\mathcal{X}$ .

Now, we impose the condition that  $a \mapsto a^n$  is an automorphism of  $A$ .

**Problem C.7.** Show that the subgroup  $A$  of  $B$  acts on  $\mathcal{X}$  with a single orbit and trivial stabilizers.

The following problem was on the problem sets; check that everyone knows how to do it:

**Problem C.8.** You have now shown that  $B$  acts on  $\mathcal{X}$ , and that the restriction of this action to  $A$  has a single orbit and trivial stabilizers. Explain why this means that  $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$  is right split.

---

<sup>1</sup>This approach is closely based on that of Kurzweil and Stellmacher, *The Theory of Finite Groups*, Chapter 3.3, Springer-Verlag (2004).