

15. THE SCHUR-ZASSENHAUS THEOREM, GENERAL CASE

Today's goal is to prove:

Theorem (Schur-Zassenhaus). Let A and C be finite groups with $\text{GCD}(\#(A), \#(C)) = 1$. Then any short exact sequence $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is right split.

We introduce the following (not standard) terminology: We'll say that a pair of groups (A, C) is **straightforward** if every short exact sequence $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is right split. On the previous worksheet, we showed that (A, C) is straightforward if A is abelian and $\text{GCD}(\#(A), \#(C)) = 1$.

Problem 15.1. Suppose that (A_1, C) and (A_2, C) are straightforward and there is a short exact sequence $1 \rightarrow A_1 \rightarrow A \rightarrow A_2 \rightarrow 1$ with A_1 canonical in A . Show that (A, C) is straightforward. **Hint/Warning:** Unfortunately, I think this first problem is one of the hardest. First use that (A_2, C) is straightforward, then use that splitting to build a new sequence which we can split using that (A_1, C) is straightforward.

Problem 15.2. Let C be a finite group, let p be a prime not dividing $\#(C)$ and let P be a p -group. Show that (P, C) is straightforward.

Let p be a prime dividing $\#(A)$ and let P be a p -Sylow subgroup of A . Let $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ be a short exact sequence, with $\text{GCD}(\#(A), \#(C)) = 1$. **Assume inductively that we have shown (A', C) is straightforward whenever $\text{GCD}(\#(A'), \#(C)) = 1$ for $\#(A') < \#(A)$.**

Recall that $N_A(P) = \{a \in A : aPa^{-1} = P\}$ and likewise for $N_B(P)$.

Problem 15.3. Show that P is canonical in $N_A(P)$.

Problem 15.4. Suppose that $A = N_A(P)$. Prove that $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is right split.

So we may now assume that $N_A(P) \neq A$.

Problem 15.5. (Fratini's argument) Show that $B = AN_B(P)$. Hint: Let $b \in B$. What can you say about bPb^{-1} ?

Problem 15.6. With A, B, C, P as above, show that $1 \rightarrow N_A(P) \rightarrow N_B(P) \rightarrow C \rightarrow 1$ is exact.

Problem 15.7. Show that $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is right split.

Remark. Problem 15.1 has uses outside of finite groups. For example, let K be a compact Lie group and let U be a simply connected nilpotent Lie group. An analogous argument shows that (U, K) is straightforward.