

PROBLEM SET 5 – DUE WEDNESDAY, OCTOBER 10

See the course website for policy on collaboration.

- Let X be a Zariski closed subset of \mathbb{A}^n . Let f be a regular function on X . The open set $\{f \neq 0\}$ is denoted $D(f)$; an open set of this form is called a distinguished open. Let $Y \subset \mathbb{A}^{n+1} = \{(x, t) : x \in X, ft = 1\}$.
 - Show that Y is isomorphic to $D(f)$ by giving regular maps in each direction.
 - Conclude that every regular function on $D(f)$ is of the form $\frac{g}{f^N}$ for some regular function g on X and some nonnegative integer N .
 - Check that basic open sets are a basis for the Zariski topology on X .
- Let $B = \text{MaxSpec } A$ and let X be Zariski closed in $B \times \mathbb{P}^n$. Show that $X = Z(I)$ for some homogenous ideal $I \subset A[x_0, x_1, \dots, x_n]$.
- Let A be a commutative ring and I and J ideals. Then $[I : J]$ is defined by

$$[I : J] := \{f \in A : fj \in I \text{ for all } j \in J\}.$$

The ideal $[I : J^\infty]$, called the *saturation of I with respect to J* is defined to be

$$[I : J^\infty] = \bigcup_{n=0}^{\infty} [I : J^n].$$

Let \overline{S} denote the Zariski closure of S .

- Let I and $J \subset k[x_1, \dots, x_n]$ be radical ideals, with $I = I(X)$ and $J = I(Y)$. Show that $[I : J]$ is radical, $[I : J] = [I : J^\infty]$ and $[I : J] = I(\overline{X \setminus (X \cap Y)})$.
 - Let I and $J \subset k[x_1, \dots, x_n]$ be ideals, not necessarily radical, with $X = Z(I)$ and $Y = Z(J)$. Show that $Z([I : J^\infty]) = \overline{X \setminus (X \cap Y)}$.
- Let $U \subset \mathbb{P}^2$ be the complement of the conic $p^2 + q^2 + r^2 = 0$. In this problem, we will show that U is isomorphic to an affine variety. A similar proof shows $\mathbb{P}^N \setminus \{F = 0\}$ is affine for any homogenous polynomial F .
Embed \mathbb{P}^2 into \mathbb{P}^5 by $\phi : (p : q : r) \mapsto (p^2 : pq : pr : q^2 : qr : r^2)$. You found that $\phi(\mathbb{P}^2)$ is closed in \mathbb{P}^5 , with equations $ux = v^2$, $uy = w^2$, $xz = y^2$, $uy = vw$, $vz = wy$ and $wx = vy$.
 - Show that $\phi(U)$ lies in a linear chart of \mathbb{P}^5 , and is closed in that linear chart.
 - Give explicit generators and relations for the ring of regular functions on U .
 - In this problem, we will work with \mathbb{P}^9 and label the homogeneous coordinates as a_{ijk} for $0 \leq i, j, k$, $i + j + k = 3$. We think of (a_{ijk}) as encoding the cubic curve $\sum a_{ijk} x^i y^j z^k$ in \mathbb{P}^2 . Show that the set of cubics which factor as (linear)(quadratic) is Zariski closed in \mathbb{P}^9 .
 - I used to believe that, if $\phi : \mathbb{A}^m \rightarrow \mathbb{A}^n$ was given by homogenous polynomials all of the same degree, then $\phi(\mathbb{A}^m)$ is Zariski closed. This is false! Give a counterexample.
 - In the proof of his second statement of Noether normalization (Theorem I.5.4.10), Shavarevich implicitly assumes the following statement. Prove it.

Let $X \subsetneq \mathbb{A}^n$ be Zariski closed in \mathbb{A}^n . Let \overline{X} be the closure of X in \mathbb{P}^n . Then there is some point of $\mathbb{P}^n \setminus \mathbb{A}^n$ which is not in \overline{X} .