

MATH 116 — PRACTICE FOR EXAM 1

Generated January 17, 2023

NAME: _____

INSTRUCTOR: _____ SECTION NUMBER: _____

1. This exam has 2 questions. Note that the problems are not of equal difficulty, so you may want to skip over and return to a problem on which you are stuck.
2. Do not separate the pages of the exam. If any pages do become separated, write your name on them and point them out to your instructor when you hand in the exam.
3. Please read the instructions for each individual exercise carefully. One of the skills being tested on this exam is your ability to interpret questions, so instructors will not answer questions about exam problems during the exam.
4. Show an appropriate amount of work (including appropriate explanation) for each exercise so that the graders can see not only the answer but also how you obtained it. Include units in your answers where appropriate.
5. You may use any calculator except a TI-92 (or other calculator with a full alphanumeric keypad). However, you must show work for any calculation which we have learned how to do in this course. You are also allowed two sides of a $3'' \times 5''$ note card.
6. If you use graphs or tables to obtain an answer, be certain to include an explanation and sketch of the graph, and to write out the entries of the table that you use.
7. You must use the methods learned in this course to solve all problems.

Semester	Exam	Problem	Name	Points	Score
Fall 2016	1	1		13	
Winter 2017	1	1		12	
Total				25	

Recommended time (based on points): 23 minutes

1. [13 points] Suppose that f is a twice-differentiable, function that satisfies

$$f(0) = 1 \qquad f(2) = 2 \qquad f(4) = 4 \qquad f'(2) = 3$$

$$\int_0^2 f(x) dx = 5 \qquad \int_2^4 f(x) dx = 7.$$

Evaluate the following integrals.

a. [4 points] $\int_0^2 x f'(x) dx$

b. [4 points] $\int_{\sqrt{2}}^2 x f'(x^2) dx$

c. [5 points] $\int_0^2 x^3 f'(x^2) dx$

1. [12 points] Suppose that f is a twice differentiable function with continuous second derivative. (That is, both f and f' are differentiable, and f'' is continuous.) The following table gives some values of f and f' .

x	0	1	2	3	4	5	6	e^3
$f(x)$	7	5	-1	0	11	-3	2	9
$f'(x)$	3	-4	-2	4	-5	0	-1	2

In parts (a) through (c) below, calculate the exact numerical value of the integral. Write “NOT ENOUGH INFO” if there is not enough information to find the exact value. Be sure to show your work clearly. No partial credit will be given for estimates.

a. [4 points] $\int_1^{e^3} \frac{f'(\ln x)}{x} dx$

b. [4 points] $\int_0^4 x f''(x) dx$

c. [4 points] $\int_2^6 f'(x) [f(x)]^2 dx$