

MATH 116 — PRACTICE FOR EXAM 2

Generated November 6, 2017

NAME: _____

INSTRUCTOR: _____ SECTION NUMBER: _____

1. This exam has 14 questions. Note that the problems are not of equal difficulty, so you may want to skip over and return to a problem on which you are stuck.
2. Do not separate the pages of the exam. If any pages do become separated, write your name on them and point them out to your instructor when you hand in the exam.
3. Please read the instructions for each individual exercise carefully. One of the skills being tested on this exam is your ability to interpret questions, so instructors will not answer questions about exam problems during the exam.
4. Show an appropriate amount of work (including appropriate explanation) for each exercise so that the graders can see not only the answer but also how you obtained it. Include units in your answers where appropriate.
5. You may use any calculator except a TI-92 (or other calculator with a full alphanumeric keypad). However, you must show work for any calculation which we have learned how to do in this course. You are also allowed two sides of a 3" x 5" note card.
6. If you use graphs or tables to obtain an answer, be certain to include an explanation and sketch of the graph, and to write out the entries of the table that you use.
7. You must use the methods learned in this course to solve all problems.

Semester	Exam	Problem	Name	Points	Score
Winter 2014	3	8		12	
Fall 2014	3	6		8	
Fall 2015	3	13		10	
Fall 2016	3	1		4	
Winter 2012	3	4 (c)		4	
Fall 2012	3	2		4	
Winter 2013	2	4 (a)		8	
Fall 2013	2	6 (c)		4	
Winter 2014	2	10		12	
Fall 2014	2	7		8	
Winter 2015	2	10		15	
Winter 2016	3	10		14	
Fall 2016	2	9		10	
Winter 2017	2	5		10	
Total				119	

8. [12 points] Suppose a_n and b_n are sequences of positive numbers with the following properties.

- $\sum_{n=1}^{\infty} a_n$ converges.
- $\sum_{n=1}^{\infty} b_n$ diverges.
- $0 < b_n \leq M$ for some positive number M .

For each of the following questions, circle the correct answer. No justification is necessary.

a. [2 points] Does the series $\sum_{n=1}^{\infty} a_n b_n$ converge?

Converge

Diverge

Cannot determine

b. [2 points] Does the series $\sum_{n=1}^{\infty} (-1)^n b_n$ converge?

Converge

Diverge

Cannot determine

c. [2 points] Does the series $\sum_{n=1}^{\infty} \sqrt{b_n}$ converge?

Converge

Diverge

Cannot determine

d. [2 points] Does the series $\sum_{n=1}^{\infty} \sin(a_n)$ converge?

Converge

Diverge

Cannot determine

e. [2 points] Does the series $\sum_{n=1}^{\infty} (a_n + b_n)^2$ converge?

Converge

Diverge

Cannot determine

f. [2 points] Does the series $\sum_{n=1}^{\infty} e^{-b_n}$ converge?

Converge

Diverge

Cannot determine

6. [8 points] Suppose that $f(x)$, $g(x)$, $h(x)$ and $k(x)$ are all positive, differentiable functions. Suppose that

$$0 < f(x) < \frac{1}{x} < g(x) < \frac{1}{x^2}$$

for all $0 < x < 1$, and that

$$0 < h(x) < \frac{1}{x^2} < k(x) < \frac{1}{x}$$

for $x > 1$. Determine whether the following statements are always, sometimes or never true by circling the appropriate answer. No justification is necessary.

- a. [2 points] $\int_0^1 g(x)dx$ converges.

Always

Sometimes

Never

- b. [2 points] $\int_0^1 f(x)dx$ diverges.

Always

Sometimes

Never

- c. [2 points] $\sum_{n=1}^{\infty} h(n)$ diverges.

Always

Sometimes

Never

- d. [2 points] $\sum_{n=1}^{\infty} k(n)$ converges.

Always

Sometimes

Never

13. [10 points] Suppose a_n and b_n are sequences with the following properties.

- $\sum_{n=1}^{\infty} a_n$ converges.

- $n \leq b_n \leq e^n$.

For each of the following statements, decide whether the statement is always true, sometimes true, or never true. Circle your answer. No justification is necessary. **You only need to answer 5 of the 7 questions.** Only answer the 5 questions you want graded. If it is unclear which 5 questions are being answered, the first 5 questions you answer will be graded.

a. [2 points] The sequence $\frac{1}{b_n}$ diverges.

ALWAYS SOMETIMES NEVER

b. [2 points] The sequence a_n is bounded.

ALWAYS SOMETIMES NEVER

c. [2 points] The series $\sum_{n=1}^{\infty} \frac{1}{b_n}$ diverges.

ALWAYS SOMETIMES NEVER

d. [2 points] The series $\sum_{n=1}^{\infty} e^{-a_n}$ converges.

ALWAYS SOMETIMES NEVER

e. [2 points] The series $\sum_{n=1}^{\infty} a_n^2$ diverges.

ALWAYS SOMETIMES NEVER

f. [2 points] The series $\sum_{n=1}^{\infty} a_n b_n$ converges.

ALWAYS SOMETIMES NEVER

g. [2 points] The series $\sum_{n=1}^{\infty} \frac{b_n}{n!}$ converges.

ALWAYS SOMETIMES NEVER

1. [4 points] Suppose that the power series $\sum_{n=0}^{\infty} c_n(x-3)^n$ converges at $x = 6$ and diverges at $x = -2$. What can you say about the behavior of the power series at the following values of x ? For each part, circle the correct answer. Ambiguous responses will be marked incorrect.

a. [1 point] At $x = -3$, the power series...

CONVERGES DIVERGES CANNOT DETERMINE

b. [1 point] At $x = 0$, the power series...

CONVERGES DIVERGES CANNOT DETERMINE

c. [1 point] At $x = 8$, the power series...

CONVERGES DIVERGES CANNOT DETERMINE

d. [1 point] At $x = 2$, the power series...

CONVERGES DIVERGES CANNOT DETERMINE

2. [5 points] Determine the radius of convergence of the power series

$$\sum_{n=0}^{\infty} \frac{(2n)!}{(n!)^2} x^{2n}.$$

Justify your work carefully and write your final answer in the space provided. Limit syntax will be enforced.

Radius of convergence = _____

4. c. [4 points] Suppose $h(x)$ and $f(x)$ are continuous functions satisfying
- $0 < f(x) \leq \frac{1}{x^p}$ for $0 < x \leq 1$.
 - $\frac{1}{x^{p+\frac{1}{2}}} \leq h(x) \leq \frac{1}{x^p}$ for $x \geq 1$.

Decide whether each of the following expressions converge, diverge or if there is not enough information available to conclude.

i. If $p = \frac{1}{2}$,

(a) $\lim_{x \rightarrow \infty} h(x)$

Converges

Diverges

Not possible to conclude.

(b) $\int_1^{\infty} h(x) dx$:

Converges

Diverges

Not possible to conclude.

ii. If $p = 2$,

(a) $\int_1^{\infty} h(x) dx$:

Converges

Diverges

Not possible to conclude.

(b) $\int_0^1 f(x) dx$

Converges

Diverges

Not possible to conclude.


4. [13 points]

a. [8 points] Consider the functions $f(x)$ and $g(x)$ where

$$\frac{1}{x^2} \leq g(x) \quad \text{for} \quad 0 < x < \frac{1}{2}.$$

$$g(x) \leq \frac{1}{x^2} \quad \text{for} \quad 1 < x$$

$$\frac{1}{x^2} \leq f(x) \quad \text{for} \quad 1 < x.$$



Using the information about $f(x)$ and $g(x)$ provided above, determine which of the following integrals is convergent or divergent. Circle your answers. If there is not enough information given to determine the convergence or divergence of the integral circle NI.

i) $\int_1^{\infty} f(x) dx$ CONVERGENT DIVERGENT NI


ii) $\int_1^{\infty} g(x) dx$ CONVERGENT DIVERGENT NI

iii) $\int_0^1 f(x) dx$ CONVERGENT DIVERGENT NI

iv) $\int_0^1 g(x) dx$ CONVERGENT DIVERGENT NI

b. [5 points] Does $\int_e^{\infty} \frac{1}{x(\ln x)^2} dx$ converge or diverge? If the integral converges, compute its value. Show all your work. Use u substitution.

6. c. [4 points] Let $f(x)$ be the differentiable function shown below. Note that $f(x)$ has a horizontal asymptote at $y = 1$.



Does $\int_2^{\infty} \frac{f'(x)}{1+f(x)} dx$ converge or diverge? Circle your answer. If it converges, find its value.

CONVERGES

DIVERGES

10. [12 points] Suppose that $g(x)$ and $h(x)$ are positive continuous functions on the interval $(0, \infty)$ with the following properties:

- $\int_1^{\infty} g(x) dx$ converges.
- $\int_0^1 g(x) dx$ diverges.
- $e^{-x} \leq h(x) \leq \frac{1}{x}$ for all x in $(0, \infty)$.

For each of the following questions, circle the correct answer.

a. [2 points] Does the integral $\int_1^{\infty} h(x)^2 dx$ converge?

Converge

Diverge

Cannot determine

b. [2 points] Does the integral $\int_0^1 h(x) dx$ converge?

Converge

Diverge

Cannot determine

c. [2 points] Does the integral $\int_1^{\infty} h(1/x) dx$ converge?

Converge

Diverge

Cannot determine

d. [2 points] Does the integral $\int_0^1 g(x)h(x) dx$ converge?

Converge

Diverge

Cannot determine

e. [2 points] Does the integral $\int_1^{\infty} g(x)h(x) dx$ converge?

Converge

Diverge

Cannot determine

f. [2 points] Does the integral $\int_1^{\infty} e^x g(e^x) dx$ converge?

Converge

Diverge

Cannot determine

7. [8 points] Suppose that $f(x)$ is a differentiable function, defined for $x > 0$, which satisfies the inequalities $0 \leq f(x) \leq \frac{1}{x}$ for $x > 0$. Determine whether the following statements are always, sometimes or never true by circling the appropriate answer. No justification is necessary.

a. [2 points] $\int_1^{\infty} f(x) dx$ converges.

Always

Sometimes

Never

b. [2 points] $\int_1^{\infty} (f(x))^2 dx$ converges.

Always

Sometimes

Never

c. [2 points] $\int_0^1 f(x) dx$ converges.

Always

Sometimes

Never


d. [2 points] $\int_1^{\infty} e^{f(x)} dx$ converges.

Always

Sometimes

Never

10. [15 points] Consider the graph below depicting four functions for $x > 0$. The only point of intersection between any two of the functions is at $x = 1$. The functions $f(x)$ and $g(x)$ are both differentiable, and they each have $y = 0$ as a horizontal asymptote and $x = 0$ as a vertical asymptote.



Use the graph to determine whether the following quantities converge or diverge, and circle the appropriate answer. If there is not enough information to determine convergence or divergence, circle "not enough information". You do not need to show your work.

a. [3 points] $\int_1^{\infty} f(x) dx$

CONVERGES

DIVERGES

NOT ENOUGH INFORMATION

b. [3 points] $\int_0^1 g(x) dx$

CONVERGES

DIVERGES

NOT ENOUGH INFORMATION

c. [3 points] $\int_0^1 g'(x) e^{-g(x)} dx$

CONVERGES

DIVERGES

NOT ENOUGH INFORMATION

d. [3 points] $\int_1^{\infty} \sqrt{g(x)} dx$

CONVERGES

DIVERGES

NOT ENOUGH INFORMATION


- e. [3 points] The volume of the solid formed by rotating the region between $f(x)$ and the x -axis from $x = 1$ to $x = \infty$ about the x -axis

CONVERGES

DIVERGES

NOT ENOUGH INFORMATION


10. [14 points] A function f has domain $[0, \infty)$, and its graph is given below. The numbers A, B, C are positive constants. The shaded region has **finite area**, but it extends infinitely in the positive x -direction. The line $y = C$ is a horizontal asymptote of $f(x)$ and $f(x) > C$ for all $x \geq 0$. The point $(1, A)$ is a local maximum of f .



- a. [5 points] Determine the convergence of the improper integral below. You must give full evidence supporting your answer, showing all your work and indicating any theorems about integrals you use.

$$\int_0^1 \frac{f(x)}{x} dx$$

10. (continued) For your convenience, the graph of f is given again. The numbers A, B, C are positive constants. The shaded region has finite area, but it extends infinitely in the positive x -direction. The line $y = C$ is a horizontal asymptote of $f(x)$ and $f(x) > C$ for all $x \geq 0$. The point $(1, A)$ is a local maximum of f .



b. [3 points] Circle the correct answer. The value of the integral $\int_1^{\infty} f(x)f'(x) dx$

is $C - A$ is $\frac{C^2 - A^2}{2}$ is $B - A$ cannot be determined diverges

c. [3 points] Circle the correct answer. The value of the integral $\int_1^{\infty} f'(x) dx$

is $C - A$ is $\frac{C^2 - A^2}{2}$ is C cannot be determined diverges

d. [3 points] Determine, with justification, whether the following series converges or diverges.

$$\sum_{n=1}^{\infty} (f(n) - C)$$

9. [10 points] Suppose that f is function with the following properties:

$$f \text{ is differentiable} \quad f(x) > 0 \text{ for all } x \quad \int_1^{\infty} f(x) dx \text{ converges.}$$

For each of the following parts, determine whether the statement is always, sometimes, or never true by circling the appropriate answer. No justification is needed.

a. [2 points] $\int_{500}^{\infty} 1000f(x) dx$ converges.

ALWAYS

SOMETIMES

NEVER

b. [2 points] $\int_1^{\infty} (f(x))^{2/3} dx$ converges.

ALWAYS

SOMETIMES

NEVER

c. [2 points] $\int_1^{\infty} (f(x))^{3/2} dx$ converges.

ALWAYS

SOMETIMES

NEVER

d. [2 points] $\int_0^1 f\left(\frac{1}{x}\right) dx$ converges.

ALWAYS

SOMETIMES

NEVER

e. [2 points] $\int_1^{\infty} \frac{f'(x)}{f(x)} dx$ converges. (Note: $\frac{f'(x)}{f(x)} = \frac{d}{dx} \ln(f(x)).$)

ALWAYS

SOMETIMES

NEVER

5. [10 points] Let $f(x)$ and $g(x)$ be two functions that are differentiable on $(0, \infty)$ with continuous derivatives and which satisfy the following inequalities for all $x \geq 1$:

$$\frac{1}{x} \leq f(x) \leq \frac{1}{x^{1/2}} \quad \text{and} \quad \frac{1}{x^2} \leq g(x) \leq \frac{1}{x^{3/4}}.$$

For each of the following, determine whether the integral always, sometimes, or never converges. Indicate your answer by circling the one word that correctly fills the answer blank. No justification is necessary. No credit will be awarded for unclear markings.

a. [2 points] $\int_1^{\infty} \sqrt{f(x)} dx$ _____ converges.

Always

Sometimes

Never

b. [2 points] $\int_3^{\infty} 4000g(x) dx$ _____ converges.

Always

Sometimes

Never

c. [2 points] $\int_1^{\infty} f(x)g(x) dx$ _____ converges.

Always

Sometimes

Never

d. [2 points] $\int_5^{\infty} g'(x)e^{g(x)} dx$ _____ converges.

Always

Sometimes

Never

e. [2 points] $\int_1^{\infty} f'(x) \ln(f(x)) dx$ _____ converges.

Always

Sometimes

Never